

Topic IX

Reactions and Mechanisms

IX.1 Module 46: Ligand Substitution in Octahedral Complexes

3/8:

- Suggested reading: Chapter 12.
- Having learned about structure, isomerism, and electronic structure, let's now talk about reactivity.
- What makes transition metals unique is the reactions of the metal center (i.e., the first coordination sphere).
- **Substitution reaction:** A reaction where a ligand in the first coordination sphere is exchanged.
- Factors that affect the rate of substitution:
 - Role of the entering group.
 - Role of the leaving group.
 - Nature of the other ligands in the complex.
 - Effect of the metal center.
- Stable/unstable are thermodynamic terms, and labile/inert are kinetic terms.
 - $[\text{Co}(\text{NH}_3)_6]^{3+}$ is unstable but inert wrt. aquation (large K_{eq} , but slow to react).
 - $[\text{Ni}(\text{CN})_4]^{2-}$ is stable but labile wrt. exchange (small K_{eq} , but equilibrium is established quickly).
- **Potential energy landscape:** A thermodynamic representation of a reaction by a (potentially multidimensional) potential energy surface along its reaction coordinate.
- Thermodynamics and kinetics are two different things, but they can be related.
- Potential energy landscapes with small ΔG and large ΔE_A are slow to react.
 - Increasing ΔG (perhaps by destabilizing the reactant) can speed up the reaction.
 - Increasing ΔG often occurs at the expense of ΔE_A (the activation energy literally decreases as the free energy change increases).
- Consider the reaction $[\text{Co}(\text{NH}_3)_5\text{X}]^{2+} + \text{H}_2\text{O} \longrightarrow [\text{Co}(\text{NH}_3)_5(\text{H}_2\text{O})]^{3+} + \text{X}^-$.
 - The more bulky/less electronegative the X-type ligand, the higher the equilibrium constant and reaction rate.
- But how do we adequately compare different reactions when so many parameters are entangled together?

- One good way to compare kinetics of transition metals is with water exchange rate constants (light/heavy water being exchanged in metal coordination spheres), since this eliminates thermodynamic consternation.
 - There is massive variation among the metals.
 - The fastest (the alkali metals) run up against the **diffusion limit**.
 - Metals with larger atomic radii react more quickly (the alkali and alkaline earth metals follow this pattern nicely, but the transition metals are all over the place).
 - Group 1A: As we go down the group, the cations are getting larger and the charge density decreases, so the $M^+ - OH_2$ bond is getting weaker and more easily broken.
 - Group 2A: The charge density is larger (doubly charged) so the strength of the bond is greater so the rate of exchange is slower.
 - We define such a reaction as having a labile half life if the half life is less than 1 minute; otherwise, it is inert.
- **Diffusion limit:** At most, one molecular collision can result in one successful reaction; thus, reaction rate is bounded above by how many molecular collisions can physically occur in a given period of time at a given temperature.
- You cannot prove mechanisms; you can only disprove other plausible scenarios.
- Possible mechanisms for $ML_5X + Y \longrightarrow ML_5Y + X$ where $X = H_2O$ and $Y = \text{anion}$ or vice versa:
- Associative (A): Via a 7-coordinate intermediate.



- Corresponds to organic S_N2 .
- First step (RDS) is slow since the incoming Y causes steric hindrance.
- The transition state is either a monocapped octahedron or pentagonal bipyramidal, depending on how the Y attacks.
- Applying a steady-state approximation for $[ML_5XY]$, we have

$$\frac{d[ML_5XY]}{dt} = 0 = k_1[ML_5X][Y] - k_{-1}[ML_5XY] - k_2[ML_5XY]$$

- We can solve the above equation for $[ML_5XY]$:

$$[ML_5XY] = \frac{k_1[ML_5X][Y]}{k_{-1} + k_2}$$

- Substituting the above into the rate constant equation for the fast step will give us the rate law in terms of the reactants (where $k = \frac{k_1 k_2}{k_{-1} + k_2}$):

$$\begin{aligned} \text{Rate} &= k_2[ML_5XY] \\ &= \frac{k_1 k_2 [ML_5X][Y]}{k_{-1} + k_2} \\ &= k[ML_5X][Y] \end{aligned}$$

- Thus, the S_N2 mechanism is second-order overall, but first-order in both reactants.

- Dissociative (*D*): Via a 5-coordinate intermediate.



- Corresponds to organic $\text{S}_{\text{N}}1$.
- First step (RDS) is slow since its spontaneous elimination of a ligand.
- The transition state is generally square pyramidal, but if it is sufficiently long-lived, it can reorganize itself into the trigonal bipyramidal state.
- The dissociative mechanism predicts that the rate of the overall substitution reaction depends on only the concentration of the original complex $[\text{ML}_5\text{X}]$, and is independent of the concentration of the incoming ligand $[\text{Y}]$.
- Thus, the overall rate law is

$$\text{Rate} = k_1[\text{ML}_5\text{X}]$$

- We can also derive this with an analogous kinetic analysis to that used for the *A* mechanism, the only difference being that we simplify $\frac{k_1 k_2 [\text{ML}_5\text{X}][\text{Y}]}{k_{-1}[\text{X}] + k_2[\text{Y}]}$ to the rate law by noting that $k_2 \gg k_{-1}$ or $[\text{Y}] \gg [\text{X}]$.
- Interchange (*I*): As Y begins to bond, X begins to leave, i.e., the bond making to Y and bond breaking to X occur simultaneously.



- Corresponds to organic $\text{S}_{\text{N}}2$.
- This is how the majority of ligand substitutions occur.
- It is too simplistic to assume that a first-order rate law implies *D* and a second-order rate law implies *A*. Indeed, most substitution reactions probably involve a mechanism like this, i.e., one that is intermediate between these two extremes.
- In an interchange mechanism, the intermediate involves an association between the original ML_5X complex and the attacking Y ligand. The Y ligand remains outside the coordination sphere of ML_5X , unlike the $\text{S}_{\text{N}}2$ mechanism, so the intermediate is not seven coordinate. However, it can help weaken the M–X bond.
- Assuming high $[\text{Y}] \approx [\text{Y}]_0$, it can be shown that the rate is given by

$$\text{Rate} = \frac{k_2 K_1 [\text{M}]_0 [\text{Y}]_0}{1 + K_1 [\text{Y}]_0}$$

where $[\text{M}]_0 = [\text{ML}_5\text{X}]_0 + [\text{ML}_5\text{X} \cdot \text{Y}]$ and $[\text{Y}]_0$ are initial conditions and $K_1 = \frac{k_1}{k_{-1}}$ is the equilibrium constant for the RDS reaction.

- At high $[\text{Y}]$ and $K_1[\text{Y}]_0 \gg 1$, the rate is first-order in $[\text{M}]_0 \approx [\text{ML}_5\text{X}]$.
- At lower $[\text{Y}]$, the rate is second order.
- At some point, the kinetic analysis becomes essentially worthless. Indeed, although we speak generally about associative and dissociative reaction mechanisms, the terms *A* and *D* are reserved for situations where 7- and 5-coordinate intermediates have actually been isolated and positively identified. If no intermediates have been isolated or identified, the designations I_d and I_a are more appropriate.
 - Two minor variations on the *I* mechanism are I_d (dissociative interchange) and I_a (associative interchange).

- If breaking the M–X bond is more important, the mechanism is I_d .
- If bond formation between ML_5X and Y is significant, the mechanism is I_a .
- The difference between I_d and I_a is subtle and does not necessarily correspond to whether the observed rate law is first or second order.
- If the rates of a series of comparable substitution reactions are most sensitive to the identity of X, the leaving ligand, then the mechanism is more probably I_d , and vice versa for I_a .
- For example, the rate constants for the anation of $[Cr(NH_3)_5(H_2O)]^{3+}$ by various ligands vary very little. Thus, it is probably I_d . However, the rate constants for the anation of $[Cr(H_2O)_6]^{3+}$ by various ligands vary by three orders of magnitude. Thus, it is probably I_a .
- Solvent (e.g., water) effects in substitution reactions:
 - Many substitution reactions occurring in solvent water may have first-order kinetics regardless of whether their initial steps are primarily D or A .
 - For example, this occurs if aquation is a precursor RDS.
 - If D , Rate = $k[ML_5X]$.
 - If A , Rate = $k[ML_5X][H_2O] = k'[ML_5X]$.
 - The latter equality is valid since water has constant concentration^[1].
 - Both mechanisms lead to apparent first-order kinetics.
- Steric factors favoring D or I_d :
 - For most octahedral complexes, steric factors inhibit formation of a CN_7 intermediate, which suggests a dissociative mechanism (D or I_d) is more plausible.
 - Even cases showing second-order kinetics may not be A for this reason.
 - For example, aquation of ammine-halides is second order with a first-order dependence on $[OH^-]$.
 - If $k_2 \gg k_1$, the rate is approximately Rate = $k_2[Co(NH_3)_5X]^{2+}[OH^-]$.
 - Calls for an alternate mechanism called S_N1CB , where ligands with lower energy bond as conjugate bases and those with higher Δ_0 (hence CB).
- Kinetically analyzes S_N1CB .

IX.2 Module 47: Substitution and Ligand Field Stabilization Energy

3/10:

- Factors that affect the rate of substitution:
 1. Role of the entering group.
 2. Role of the leaving group.
 3. The nature of the other ligands in the complex.
 4. Effect of the metal center.
- Six factors that support a dissociative mechanism:
 1. The rate of reaction only slightly (within a factor of 10) changes on the incoming ligand.
 - Means that the ligand is not significantly involved in the RDS.
 2. Making the charge on complex more positive decreases the rate of substitution.

¹Is this what we meant when we said in Labalme (2020) that liquids and solids don't have active mass? In other words, the species should be included in the mass-action expression; they just simply get lumped in with the rate constant generally.

- Increasing the charge increases the bond strength, making it harder for one ligand to spontaneously break away.
- 3. Steric crowding increases reaction rate.
- 4. The volume of activation is positive and reaction rate decreases at high pressure.
 - Especially if the reactant is a gas, for example.
- 5. Reaction rate correlates with Ligand Field Activation Energy (LFAE) predictions.
- 6. Stereochemistry can give additional hints.
 - A messy subject unless you use very well crafted multidentate ligands.
- Simple changes in oxidation state can have massive effects on water exchange rate constants^[2] (e.g., Cr^{2+} to Cr^{3+} results in a shift of 15 orders of magnitude).
- Variations in complex reactivity:
 - Most first row transition metals are labile^[3], but Cr^{3+} (d^3 , $S = \frac{3}{2}$) and low-spin Co^{3+} (d^6 , $S = 0$) are usually inert.
 - d^7 - d^{10} , with filling of e_g^* levels, are labile.
 - These configurations tend to have large Jahn-Teller distortions and/or low CFSEs.
 - d^7 , d^9 , and d^{10} cases are more labile than d^8 .
 - d^8 has a $^3A_{2g}$ ground state, which is immune to Jahn-Teller distortion.
 - With strong-field ligands, d^8 may be square planar, often being inert.
 - Inert octahedral complexes tend to be those with high CFSE; viz., d^3 low-spin d^4 - d^6 .
 - ML_6 complexes of both d^3 ($^4A_{2g}$) and low-spin d^6 ($^1A_{1g}$) are immune from Jahn-Teller distortions and therefore can be perfect O_h .
 - ML_6 d^3 has $\text{CFSE} = -\frac{6}{5}\Delta_o$ and low-spin d^6 has $\text{CFSE} = -\frac{12}{5}\Delta_o + 3P$.
- The most inert complexes among the first row transition metals:
 - Co^{3+} is primarily dissociative, but with dependence on the incoming ligand I_d .
 - Cr^{3+} has a dimorphism where it can be I_d or I_a !
 - For example, it is I_d in the complex $[\text{Cr}(\text{NH}_3)_5(\text{H}_2\text{O})]^{3+}$ but I_a in the complex $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$.
- Water exchange rate constants are greatest for Jahn-Teller distorted species.
 - They are smallest for $d^{3,5,8}$.
- Variations in complex reality: Explained by the potential energy landscape for a substitution reaction $\text{ML}_6 \longrightarrow \text{ML}_5\text{Y}$:
 - O_h to C_{4v} to “ O_h .”
 - The activated complex is square pyramidal.
 - We can now use the AOM to calculate the LFSE in both the initial O_h state and the transition C_{4v} state.
 - The difference between them will be the **ligand field activation energy** or **LFAE**. The LFAE is not particularly useful because it predicts negative activation energies, which are not a thing, but it does allow us to rationalize reactivity trends.
 - If $\text{LFAE} \leq 0$, the reaction is labile; otherwise, it is moderate or slow.

²We pronounce the units of rate constants, “reciprocal seconds.”


³Werner was only able to make his conclusions about coordination chemistry because he happened to be working with the relatively inert Co^{3+} complexes.

IX.3 Module 48: Ligand Substitution in Square Planar Complexes


- Although steric factors favor *D*-type mechanisms for octahedral complexes, square planar ML_4 complexes are not so inhibited.
 - For square planar ML_4 complexes, an associative (*A*) mechanism, in which a coordination number 5 (CN5) intermediate is formed, is plausible.
- The d^8 metals are especially likely to form square planar complexes.
- Substitution of square planar complexes, such as $PtLX_3$, leads to *trans* and *cis* isomers.
- The rate law is

$$\text{Rate} = k_1[PtLX_3] + k_2[PtLX_3][Y^-]$$


- Suggests two paths, where the first term may be pseudo-first-order due to excess solvent acting as an attacking group.
- If we make L bulkier, we would expect the *D* pathway to dominate (k_1 increases, k_2 decreases).
- However, both k_1 and k_2 decrease with increasingly bulky ligands, suggesting an associative mechanism:



- S means the solvent.
 - Since $[S] \gg [ML_4]$, we see pseudo-first order kinetics in the k_1 term in the rate law.
- In square planar complexes, *d*-orbital splitting yields a $d_{x^2-y^2}$ LUMO.
 - However, it points exactly toward the ligands, so you cannot fill it without displacing a ligand.
 - Indeed, since ligands cannot attack it, they instead attack the higher-lying metal p_z orbital.
- Note that the stereochemistry stays the same under these reactions.
- An example of the *trans* effect:



- The first ligand takes any of the four spots, but the second substitutes opposite the stronger *trans*-directing ligand.
 - In the above example, chloride is a stronger *trans*-directing ligand than amine groups.
- The *trans* effect:
 - The ratio of *trans* and *cis* isomers is found to vary with the ability of L to act as a *trans*-directing ligand.
 - The increasing order of *trans*-directing ability is



- This ranking comes from a combination of the following two factors.
 - The effect is kinetic rather than thermodynamic.

- Two factors explain the trans effect (both aim at lowering ΔG^\ddagger):
 1. Weakening of the Pt–X bond *trans* to the directing ligand (σ -donor effects).
 - Destabilizes the ground state of the reactants.
 - Pt–A is influenced by *trans*-Pt–T bond because both share Pt p_x and $d_{x^2-y^2}$ orbitals. When the Pt–T bond is strong, electron density on those orbitals is shifted away from the Pt–A bond.
 - The more polarizable the L ligand, the better *trans*-director it is; e.g., $\text{I}^- > \text{Br}^- > \text{Cl}^-$.
 - The *trans*-directing ligand polarizes the metal ion, inducing a slight repulsion with the negative electron density on the leaving ligand in the *trans*-position.
 - For example, the Pt–Cl bond is longer in *cis*-[Pt(PMe₃)₂(Cl)₂] than in the *trans* form because the stronger σ -donor ability of PMe₃ weakens the opposite Pt–Cl bond in the *cis* form.
 2. Stabilization of the presumed CN5 intermediate (π -acceptor effects).
 - Stabilizes the activated complex.
 - The strongest *trans*-directors are good π -acceptor ligands.
 - Assuming an *A* mechanism, substitution involves a trigonal bipyramidal transition state.
 - The *trans* intermediate (activated complex) is more favorable for π -acceptor ligands because it permits π -delocalization in the trigonal plane.
 - When the T-ligand engages in a strong π -backbonding, charge is removed from Pt, making the metal center more electrophilic and stabilizing the TBP intermediate.
- The trans effect has applications to synthesis (synthesizing *cis* versus *trans* square planar compounds).
 - For example, *cis*-[Pt(NH₃)₂(Cl)₂] is cisplatin, the first anti-cancer drug.