

Topic VI

Bonding and Physical Properties of Metal Complexes

VI.1 Module 34: Magnetic Properties of Transition Metal Complexes

- 2/22:
- Electrons occupy the lowest energy triply degenerate orbitals in d^1 , d^2 , and d^3 configurations.
 - However, in the d^4 configuration:
 - Low spin: The fourth electron will pair up in the lower t_{2g} energy level.
 - High spin: The fourth electron will occupy a higher energy e_g orbital.
 - The pairing energy Π is made up of two parts (refer to Figure 0.9 and the associated discussion):
 1. Coulombic repulsion energy caused by having two electrons in the same orbital. Destabilization energy contribution of Π_c for each doubly occupied orbital. Has a positive sign because it increases the energy of the system.
 2. Exchange stabilization energy for each pair of electrons having the same spin and same energy. Stabilizing contribution of Π_e for each pair having same spin and same energy. Has a negative sign because it reduces the energy of the system.
 - Deciding whether the fourth electron will go into the higher energy e_g orbital at an energy cost of Δ , or be paired at an energy cost of Π .
 - Strong field ligand has big Δ so $\Pi < \Delta$; this implies a low spin configuration.
 - Weak field ligand has small Δ so $\Pi > \Delta$; this implies a high spin configuration.
 - We can experimentally discriminate between high- and low-spin compounds by measuring magnetic properties.
 - The Gouy balance can determine the magnetic susceptibility of materials.
 - A more modern way to measure magnetic properties uses a Superconducting Quantum Interference Device, or SQUID.
 - This device is just about the most sensitive machine humanity can build (can detect the magnetic field of the heart/brain).
 - Main types of magnetic behavior:
 - Diamagnetism (from electron charge).
 - Paramagnetism (spin and orbital motion of electrons on individual atoms).

- Ferromagnetism and antiferromagnetism (cooperative interaction between magnetic moments of individual atoms).
- Paramagnetism is much stronger than diamagnetism and overpowers it.
 - Ferromagnetism overpowers both.
- Theoretical background for determining magnetic spins experimentally:
 - When we place a sample in a magnetic field of magnitude H , the sample will interact with the magnetic field and magnetize. This magnetization causes the magnetic flux B in the material to differ from the magnetic flux through the space the sample occupies (were the sample not there) by an amount determined by the magnetization parameter M , which is specific to each material. These three quantities are related via the equation

$$B = H + 4\pi M$$

- If we divide the flux by the magnetic field, we obtain the magnetic susceptibility per unit volume κ of the material:

$$\frac{B}{H} = 1 + 4\pi \cdot \frac{M}{H} = 1 + 4\pi\kappa$$

- This quantity can be normalized by the molecular weight and density of the substance to give the magnetic susceptibility per mole

$$\chi_M = \kappa \cdot \frac{\text{molecular weight}}{\text{density}}$$

- Dividing χ_M by Avogadro's number gives the magnetic susceptibility per molecule χ_M^{corr} .
- **Curie's law** relates χ_M^{corr} to the magnetic moment μ by the formula

$$\chi_M^{\text{corr}} = \frac{N\mu^2 k}{3T}$$

where N is Avogadro's number, $k = 1.381 \times 10^{-23}$ J/K is the Boltzmann constant, and T is the absolute temperature of the substance.

- Note that μ is measured in units of Bohr magnetons where $1 \text{ BM} = \frac{eh}{4\pi m_e c}$. As per usual, we have $e = 1.602 \times 10^{-19}$ C is the charge of an electron, $h = 6.626 \times 10^{-34}$ Js is Planck's constant, $m_e = 9.11 \times 10^{-31}$ kg is the mass of an electron, and $c = 2.998 \times 10^8$ m/s is the speed of light.
- We can rearrange Curie's law to express the magnetic moment in terms of χ_M^{corr} as follows.

$$\mu = \sqrt{3k/N} \cdot \sqrt{\chi_M^{\text{corr}} T}$$

- Magnetic moment μ and the spin-only formula: Materials that are diamagnetic are repelled by a magnetic field, whereas paramagnetic substances are attracted into a magnetic field, i.e., show magnetic susceptibility. The unpaired electrons in paramagnetic complexes of 3d-block metal ions create a magnetic field. The magnetic moment μ is then given by the spin-only formula

$$\mu_{\text{spin-only}} = \sqrt{n(n+2)}$$

where n is the number of unpaired electrons.

- In heavier transition metals, we need to account for not just the S quantum number but also L (which accounts for some ground state relativistic effects) by using the formula

$$\mu_{S+L} = \sqrt{4S(S+1) + L(L+1)}$$