

# CHEM 20100 (Inorganic Chemistry I) Problem Sets

Steven Labalme

January 22, 2021

# Contents

0	Course Prep Problems	1
---	----------------------	---

## 0 Course Prep Problems

9/13: **2.8** The details of several steps in the particle-in-a-box model in this chapter have been omitted. Work out the details of the following steps:

- a. Show that if  $\Psi = A \sin rx + B \cos sx$  ( $A$ ,  $B$ ,  $r$ , and  $s$  are constants) is a solution to the wave equation for the one-dimensional box, then

$$r = s = \sqrt{2mE} \left( \frac{2\pi}{h} \right)$$

*Solution.*

$$\begin{aligned} \frac{-h^2}{8\pi^2m} \cdot \frac{\partial^2 \Psi(x)}{\partial x^2} &= E\Psi(x) \\ \frac{-h^2}{8\pi^2m} \cdot \frac{\partial^2}{\partial x^2} (A \sin rx + B \cos sx) &= E(A \sin rx + B \cos sx) \\ \frac{-h^2}{8\pi^2m} \cdot \frac{\partial}{\partial x} (Ar \cos rx - Bs \sin sx) &= E(A \sin rx + B \cos sx) \\ \frac{-h^2}{8\pi^2m} \cdot (-Ar^2 \sin rx - Bs^2 \cos sx) &= E(A \sin rx + B \cos sx) \\ \frac{Ar^2 h^2}{8\pi^2m} \sin rx + \frac{Bs^2 h^2}{8\pi^2m} \cos sx &= AE \sin rx + BE \cos sx \\ 0 &= \left( \frac{Ar^2 h^2}{8\pi^2m} - AE \right) \sin rx + \left( \frac{Bs^2 h^2}{8\pi^2m} - BE \right) \cos sx \end{aligned}$$

Choose  $x = 0$ .

$$\begin{aligned} &= \frac{Bs^2 h^2}{8\pi^2m} - BE \\ E &= \frac{s^2 h^2}{8\pi^2m} \\ \frac{8\pi^2mE}{h^2} &= s^2 \\ s &= \sqrt{\frac{8\pi^2mE}{h^2}} \\ \boxed{s = \sqrt{2mE} \frac{2\pi}{h}} \end{aligned}$$

With this result ...

$$\begin{aligned} 0 &= \left( \frac{Ar^2 h^2}{8\pi^2m} - AE \right) \sin rx + \left( \frac{Bs^2 h^2}{8\pi^2m} - BE \right) \cos sx \\ &= \left( \frac{Ar^2 h^2}{8\pi^2m} - AE \right) \sin rx + \left( B \left( \frac{s^2 h^2}{8\pi^2m} \right) - BE \right) \cos sx \\ &= \left( \frac{Ar^2 h^2}{8\pi^2m} - AE \right) \sin rx + (BE - BE) \cos sx \\ &= \left( \frac{Ar^2 h^2}{8\pi^2m} - AE \right) \sin rx \end{aligned}$$

Choose  $x = \frac{\pi}{2r}$ .

$$= \frac{Ar^2 h^2}{8\pi^2m} - AE$$

$$r = \sqrt{2mE} \frac{2\pi}{h}$$

□

- d. Show that substituting the value of  $r$  given in part c into  $\Psi = A \sin rx$  and applying the normalizing requirement gives  $A = \sqrt{2/a}$ .

*Solution.*

$$\begin{aligned} 1 &= \int_{\text{all space}} \Psi \Psi^* d\tau \\ &= \int_0^a \left( A \sin \frac{n\pi x}{a} \right) \left( A \sin \frac{n\pi x}{a} \right) dx \\ &= \int_0^a A^2 \sin^2 \frac{n\pi x}{a} dx \end{aligned}$$

Use  $\sin^2 u = \frac{1 - \cos 2u}{2}$ .

$$\begin{aligned} &= A^2 \int_0^a \frac{1 - \cos \frac{2n\pi x}{a}}{2} dx \\ &= \frac{A^2}{2} \left( \int_0^a dx - \int_0^a \cos \frac{2n\pi x}{a} dx \right) \\ &= \frac{A^2}{2} \left( [x]_0^a - \left[ \frac{a}{2n\pi} \sin \frac{2n\pi x}{a} \right]_0^a \right) \\ &= \frac{A^2}{2} \left( (a - 0) - \left( \frac{a}{2n\pi} \sin 2n\pi - \frac{a}{2n\pi} \sin 0 \right) \right) \\ &= \frac{A^2}{2} \left( a - \left( \frac{a}{2n\pi} \sin 2n\pi \right) \right) \end{aligned}$$

Since  $n$  is an integer,  $\sin 2n\pi = 0$ .

$$\begin{aligned} &= \frac{aA^2}{2} \\ \frac{2}{a} &= A^2 \\ A &= \sqrt{\frac{2}{a}} \end{aligned}$$

□