

Week 2

Spectrometry

2.1 Office Hours (Snyder)

- 1/17:
- Does cyclohexane only have one ^{13}C NMR signal, and only one ^1H NMR signal?
 - 1 singlet for ^{13}C .
 - 1 singlet for ^1H .
 - We don't integrate carbon.
 - We only integrate to compare things.
 - We won't have to deal with cyclohexane conformations wrt. NMR on any test.
 - What do we need to know about the Karplus correlation?
 - We won't need it for problems.
 - It's useful, but we've got other things to worry about.
 - Do chemists/when do chemists run ^{13}C NMR experiments with all carbons isotopically carbon-13?
 - Is the reason we don't integrate carbon because the placing of the carbon-13s is random? Would the proportions not still be representative?
 - For ^1H NMR, feel free to draw in the hydrogen atoms on the line-angle structure.
 - Multiplying $n + 1$ of different types of neighbors (e.g., if a hydrogen has 3 neighboring hydrogens to one side and 2 neighboring hydrogens to the other side, it has a maximum of $(3 + 1)(2 + 1) = 12$ peaks in its signal).
 - The multiplication analysis applies only to chains that are completely different.

2.2 Chapter 9: Nuclear Magnetic Resonance and Mass Spectroscopy

From Solomons et al. (2016).

- 1/18:
- **Mass spectrometry:** The formation of ions in a mass spectrometer followed by separation and detection of the ions according to mass and charge.
 - **Mass spectrum:** A graph that on the x -axis represents the formula weights of the detected ions, and on the y -axis represents the abundance of each detected ion.
 - The x -axis is labeled m/z where m is mass and z is charge.




Figure 2.1: The mass spectrum of propane.

- The examples Solomons et al. (2016) consider all have $z = +1$, so the x -axis in them effectively represents the formula weight of each detected ion.
- **Base peak:** The tallest peak in a mass spectrum.
 - Relative ion abundance on the y -axis is either expressed as a percentage of the base peak or directly as the number of detected ions.
 - Usually an easily formed fragment of the original compound.
 - The base peak in Figure 2.1 corresponds to the C_2H_5^+ ion, $m/z = 29 = 2 \cdot 12 + 5 \cdot 1$.
- **Molecular ion:** The ion with the formula weight of the original compound.
 - One of the higher value m/z peaks.
 - Usually not the base peak.
- Small peaks having m/z values 1 or 2 higher than the formula weight of the compound are due to ^{13}C and other isotopes.
- **Electron impact:** A method for ionizing molecules in a mass spectrometer by placing the sample under high vacuum and bombarding it with a beam of high-energy electrons. *Also known as EI.*
 - The energy of the electrons is in the range of 70 eV or 6.7×10^3 kJ/mol.
 - The incoming electrons ionize the molecules to molecular ions, which are radical cations since they have a +1 charge and an unshared electron.
- Note that there are ionization methods other than EI, but it is the most common.
- Localizing the radical and charge along the structure.




Figure 2.2: Molecular ions.

- The choice of where we localize the radical/charge is often arbitrary (esp. with hydrocarbons).

- However, “as we might expect, ionization potentials indicate that in [the] formation of radical cations, the nonbonding electrons of nitrogen, oxygen, and halogen atoms, and the π electrons of alkenes and aromatic molecules, are held more loosely than the electrons of carbon-carbon and carbon-hydrogen σ bonds” (Solomons et al., 2016, p. 425).
- Thus, “when a molecule contains oxygen, nitrogen, or a π bond, we place the odd electron and charge at a nitrogen, oxygen, halogen, or π bond. If resonance is possible, the radical cation may be delocalized” (Solomons et al., 2016, p. 425).
- Three important principles.
 1. The reactions that take place are all unimolecular since the pressure is kept so low.
 2. Single-barbed arrows denote the movement of single electrons.
 3. The relative ion abundances give key information about the structures of the fragments produced and their original locations in the molecule.
- Fragmentation by cleavage at a single bond.
 - When such a process happens in a molecular ion, a cation and a radical are produced, although only the cation will be detected by the positive ion mass spectrometers we’re considering.
 - Each cleavage can happen in two ways (since one fragment will take the radical and the other will take the positive charge).
 - The path that produces the more stable carbocation will occur more rapidly.
 - Notice the difference in relative ion abundance between the secondary CH_3CH_2^+ ($m/z = 29$) and the primary CH_3^+ ($m/z = 15$) in Figure 2.1.
- When drawing cleavage reactions, use brackets and delocalization; when drawing cleavage mechanisms, use localization.
- Chain branching increases the likelihood of cleavage at a branch point because a more stable carbocation can result.
- Examples of fragmentation to form resonance-stabilized cations.
 1. Alkenes ionize and frequently undergo fragmentations that yield resonance-stabilized allylic cations.




Figure 2.3: Resonance fragmentation: Alkenes.

2. Carbon-carbon bonds next to an atom with a lone pair usually break readily because the resulting carbocation is resonance stabilized.




Figure 2.4: Resonance fragmentation: Lone pairs.

3. Carbon-carbon bonds next to the carbonyl group of an aldehyde or ketone break readily because resonance-stabilized ions called **acylium ions** are produced.






Figure 2.5: Resonance fragmentation: Carbonyls.

– Note that either the C–R or the C–R' bond could break.

4. Alkyl substituted benzenes ionize by loss of a π electron and undergo loss of a hydrogen atom or methyl group to yield the relatively stable **tropylium ion**. This fragmentation gives a prominent peak (sometimes the base peak) at $m/z = 91$.



(a) Losing a hydrogen radical.



(b) Losing a methyl radical.

Figure 2.6: Resonance fragmentation: Alkyl-substituted benzene rings.

5. Monosubstituted benzenes with other than alkyl groups also ionize by loss of a π electron and then lose their substituent to yield a phenyl cation with $m/z = 77$.





Figure 2.7: Resonance fragmentation: Monosubstituted benzene rings with nonalkyl groups.

– Y is a halogen, nitro group, acyl group, nitrile group, etc.

- Fragmentation by cleavage of two bonds leads to a new radical cation and a neutral molecule.

1. Alcohols frequently show a peak at $M^+ - 18$. This corresponds to the loss of a molecule of water.

Figure 2.8: Fragmentation: Loss of H₂O.

2. Carbonyl compounds with a hydrogen on their γ carbon undergo a fragmentation called the McLafferty rearrangement.




Figure 2.9: Fragmentation: McLafferty rearrangement.

- Y may be an alkyl, hydride, ether, hydroxyl, etc.
3. There are also often peaks corresponding to the elimination of other small molecules.
- Isotope effects:
 - The presence of ^{13}C will provide a small peak at $M^+ + 1$.
 - “In the mass spectrum for a sample containing chlorine, we would expect to find peaks separated by two mass units, in an approximately 3 : 1 (75.5% : 24.5%) ratio for the molecular ion or any fragments that contain chlorine” (Solomons et al., 2016, p. 432).
 - “In the mass spectrum for a sample containing bromine, we would expect to find peaks separated by two mass units in an approximately 1 : 1 ratio (50.5% : 49.5% ^{79}Br to ^{81}Br)” (Solomons et al., 2016, p. 433).
 - In a molecule containing two bromine atoms, for example, we’ll see peaks at M^+ , $M^+ + 2$, and $M^+ + 4$ in a 1 : 2 : 1 ratio.