

Week 6

???

6.1 Examples of Group Actions

10/31:

- Today: A number of interesting group actions.
- **Left action** (of G on X): A group action of the form $g \cdot x$ (as opposed to $x \cdot g$).
- Let G be a group, and let $X = G$. Take $g \cdot x = gx$.
 - Axiom confirmation.
 1. $e \cdot x = ex = x$.
 2. $g \cdot (h \cdot x) = ghx = gh \cdot x$.
 - Let $e \in X$. Then $\text{Orb}(e) = X$. In particular, this means that the action is transitive.
 - $\text{Stab}(x) = \{g \in G \mid gx = x\} = \{e\}$ for $x \in X$ arbitrary, in general.
 - $\ker = \{e\}$. This also follows from the above. Thus, the action is faithful.
- Corollary: Let G be a finite group. Then G is isomorphic to a subgroup of S_n for some n . We may take $n = |G|$.
 - Construction: We invoke the proposition from last lecture. In particular, we know that the action $G \curvearrowright G$ implies the existence of a homomorphism $\phi : G \rightarrow S_G$ defined by $g \mapsto \psi_g$.
 - The map in the above construction has trivial kernel. By the FIT, $G/\ker \cong \text{im } \phi$. Combining these results, we obtain $G \cong G/\ker \cong \text{im } \phi \leq S_n$.
 - Applying this construction to S_3 , we deduce that $S_3 \leq S_6$.
- $\text{SO}(2) \cong \mathbb{R}/\mathbb{Z} \cong \mathbb{Q}/\mathbb{Z} \oplus \mathbb{Q}^\infty$.
 - In infinite cases, you usually want to consider some other topological things that disappear in the finite case.
- Let G be a group and take $X = G$ again. We can also consider $g \cdot x = gxg^{-1}$.
 - Axioms.
 1. $e \cdot x = exe^{-1} = x$.
 2. $g \cdot (h \cdot x) = ghxh^{-1}g^{-1} = (gh)x(gh)^{-1} = gh \cdot x$.
 - $\text{Orb}(e) = \{e\}$; not transitive if $|G| > 1$.
 - Let $x \in X$. Then $\text{Orb}(x)$ is the conjugacy class of x .
 - $\text{Stab}(x) = C_G(x)$.
 - $\ker = Z(G)$. Thus, the group action is faithful iff the center is trivial. Abelian implies not faithful.

- A nice thing about these constructions is that they cast other constructions we've encountered in the more general language of group actions.
- **Right actions** are even nastier than left cosets and right cosets, so Calegari will not mention them again.

$$- g \cdot x = x \cdot g^{-1} \text{ and } g \cdot (h \cdot x) = (x \cdot h^{-1}) \cdot g^{-1}.$$

- Let $G = G$, X be the subgroups of G . $g \cdot H = gHg^{-1}$.
 - Note that $H \leq G$ does indeed imply that $gHg^{-1} \leq G$. In particular, ...
 - H is nonempty (contains at least e), so $gHg^{-1} \supset \{geg^{-1}\}$ is nonempty;
 - $gh_1g^{-1}, gh_2g^{-1} \in gHg^{-1}$ imply that $gh_1g^{-1}gh_2g^{-1} = g(h_1h_2)g^{-1} \in gHg^{-1}$;
 - $ghg^{-1} \in gHg^{-1}$ has inverse $gh^{-1}g^{-1} \in gHg^{-1}$.
 - Axioms (entirely analogous to the last example).
 - $\text{Orb}(H)$ is the "conjugates" of H .
 - $\text{Stab}(H) = N_G(H)$.
 - $\ker = ?$. We know that $Z(G) \subset \ker$. The conclusion is that there is not a nice definition for the kernel other than the intersections of the stabilizers/normalizers.
 - ...
 - If any $H \triangleleft G$ is normal, and $x \in G$ had order 2, then $\langle x \rangle \triangleleft G$, meaning that $gxg^{-1} \in \langle x \rangle$, i.e., $x \in Z(G)$, so this rules out D_8 ??
- Fix G and $H \leq G$. Let $X = G/H$ (not assuming $H \triangleleft G$, so we know that G/H is the set of left cosets but it is not a group in general). Define $g \cdot xH = gxH$.
 - We have $g \cdot xhH = gxhH$.
 - Orbit: $\text{Orb}(eH) = X$.
 - Stabilizer: $\text{Stab}(eH) = H$.
 - $\text{Stab}(gH) = gHg^{-1}$.
 - This is because $(ghg^{-1})gH = ghH = gH$.
 - Go to the more general case $G \supset X$, $\text{Stab}(x) = H$. Then $gHg^{-1} \subset \text{Stab}(g \cdot x)$??
 - Transitive: Yes (see orbits).
 - Faithful: If H is normal, no. If H contains a normal subgroup, no. Maybe yes.
 - Kernel: If H is normal, then $\ker = H$. In general, $\ker = \bigcap_{g \in G} gHg^{-1}$ (the largest normal subgroup of H).
- Takeaway: General constructions allow us to see things we've already done.
- Next time: The most useful theorem of the course, that provides lots of information on relations between objects.