

FOURIER SERIES

- We have shown that when $v^T w = 0$, then vectors v and w are orthogonal.
- Fourier series ask us to think of continuous functions as vectors.
- Let $f(t) = \cos(t)$ and let $g(t) = \sin(t)$
- To find $f \cdot g$ of these two function “vectors” would be asking us to take a dot product that has infinitely many terms. All vectors we have used have had a finite number of components. Because these “vectors” would have infinitely many terms, it would be like asking us to take an integral!
- $f \cdot g = \int_0^{2\pi} \cos(t) \sin(t) dt$ (We limit the domain because these are sinusoidal periodic functions and thus they repeat after a period of 2π .)

$$\int_0^{2\pi} \cos(t) \sin(t) dt$$

Let $u = \cos(t)$ and $du = -\sin(t)$

$$-\int_1^1 u du$$

$$\left[-\frac{u^2}{2} \right]_1^1$$

0

OR

$$\left[-\frac{1}{2} \cos^2(t) \right]_0^{2\pi}$$

$$-\frac{1}{2} (\cos^2 2\pi - \cos^2 0)$$

$$-\frac{1}{2} (1 - 1) =$$

0

- Therefore $\cos t \cdot \sin t = 0$, therefore these function “vectors” are orthogonal, therefore they serve as basis vectors for the function space of continuous periodic functions in F^2 , therefore any continuous function also in this space can be written as a linear combination of $\cos(t)$ and $\sin(t)$!!!

$$f(t) = \frac{a_0}{2} + a_1 \cos(t) + b_1 \sin(t) + a_2 \cos(2t) + b_2 \sin(2t) + \dots$$

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nt) + b_n \sin(nt))$$

*Why divide the constant term by 2?

CALCULATING THE FOURIER COEFFICIENTS

As sine and cosine can serve as an orthogonal basis for periodic functions, consider the Fourier Series for a function

$$f(t) \text{ of period } 2\pi \text{ to be } f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nt) + b_n \sin(nt))$$

In problem 3 from your handout you already found a formula for a_0 , the constant term.

$$\frac{a_0}{2} = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) dt$$

To obtain the coefficients $a_n, n \in \mathbb{Z}$, multiply both sides of the equation by $\cos(mt)$ where $m \in \mathbb{Z}$ and $m > 0$ and integrate both sides from $-\pi$ to π since this is a periodic function of period 2π .

$$\int_{-\pi}^{\pi} f(t) \cos(mt) dt = \frac{a_0}{2} \int_{-\pi}^{\pi} \cos(mt) dt + \sum_{n=1}^{\infty} \left(a_n \int_{-\pi}^{\pi} \cos(nt) \cos(mt) dt + b_n \int_{-\pi}^{\pi} \sin(nt) \cos(mt) dt \right)$$

Simplify this result using the integrals on the right side of this equation from the handout problems 5-7. The only nonzero integral results from $\int_{-\pi}^{\pi} \cos^2(mt) dt = \pi$ in the case where $n = m$.

$$\therefore \int_{-\pi}^{\pi} \cos(mt) dt = a_m \pi$$

Since this is the case where $n = m$, replacing m with n and solving for a_n we obtain:

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \cos(nt) dt, n \in \mathbb{Z} \text{ and } n > 0$$

To obtain the coefficients $b_n, n \in \mathbb{Z}, n > 0$, multiply both sides of the equation by $\sin(mt)$ where $m \in \mathbb{Z}$ and $m > 0$ and integrate both sides from $-\pi$ to π since this is a periodic function of period 2π .

$$\int_{-\pi}^{\pi} f(t) \sin(mt) dt = \frac{a_0}{2} \int_{-\pi}^{\pi} \sin(mt) dt + \sum_{n=1}^{\infty} \left(a_n \int_{-\pi}^{\pi} \cos(nt) \sin(mt) dt + b_n \int_{-\pi}^{\pi} \sin(nt) \sin(mt) dt \right)$$

Simplify this result using the integrals on the right side of this equation using the handout problems you did for 8. The only nonzero integral results from $\int_{-\pi}^{\pi} b_m \sin^2(mt) dt = b_m \pi$ in the case where $n = m$. Relabeling m as n and solving for b_n we obtain:

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) \sin(nt) dt, n \in \mathbb{Z} \text{ and } n > 0$$