

Complex Linear Independence: Decomplexification

4/7:

- When given a complex system of equations, it is necessary to **decomplexify** it.
- **Decomplexify:** To model a complex system of equations with a strictly real system for the purpose of applying the tenets of real linear algebra to it.
- Consider the following complex system of equations.

$$\begin{aligned}(2+i)x_1 + (1+i)x_2 &= 3+6i \\ (3-i)x_1 + (2-2i)x_2 &= 7-i\end{aligned}$$

– The solutions will be complex numbers: $x_1 = a_1 + ib_1$ and $x_2 = a_2 + ib_2$, where $a_1, a_2, b_1, b_2 \in \mathbb{R}$.

- Transform it into a matrix system of equations. Separate the real and complex parts, and factor out all instances of the imaginary number i so that it is a coefficient to any complex matrix.

$$\begin{aligned}\begin{bmatrix} 2+i & 1+i \\ 3-i & 2-2i \end{bmatrix} \begin{bmatrix} a_1+ib_1 \\ a_2+ib_2 \end{bmatrix} &= \begin{bmatrix} 3+6i \\ 7-i \end{bmatrix} \\ \left(\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} + \begin{bmatrix} i & i \\ -i & -2i \end{bmatrix} \right) \left(\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} + \begin{bmatrix} ib_1 \\ ib_2 \end{bmatrix} \right) &= \left(\begin{bmatrix} 3 \\ 7 \end{bmatrix} + \begin{bmatrix} 6i \\ -i \end{bmatrix} \right) \\ \underbrace{\left(\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} + i \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \right)}_A \underbrace{\left(\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} + i \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \right)}_x &= \underbrace{\left(\begin{bmatrix} 3 \\ 7 \end{bmatrix} + i \begin{bmatrix} 6 \\ -1 \end{bmatrix} \right)}_b\end{aligned}$$

- Foil the left side of the above equation^[1].

$$\left(\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \right) + i \left(\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \right) = \begin{bmatrix} 3 \\ 7 \end{bmatrix} + i \begin{bmatrix} 6 \\ -1 \end{bmatrix}$$

- Split the above system of equations into a real system of equations and a complex system of equations by setting equal to each other the real components of each side and the imaginary components of each side.

$$\begin{aligned}\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} &= \begin{bmatrix} 3 \\ 7 \end{bmatrix} \\ \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} &= \begin{bmatrix} 6 \\ -1 \end{bmatrix}\end{aligned}$$

- Multiply out the matrices above to yield a system of four equations.

$$\begin{aligned}2a_1 + a_2 - b_1 - b_2 &= 3 \\ 3a_1 + 2a_2 + b_1 + 2b_2 &= 7 \\ a_1 + a_2 + 2b_1 + b_2 &= 6 \\ -a_1 - 2a_2 + 3b_1 + 2b_2 &= -1\end{aligned}$$

- Condense the above system of equations into a single matrix system of equations.

$$\begin{bmatrix} 2 & 1 & -1 & -1 \\ 3 & 2 & 1 & 2 \\ 1 & 1 & 2 & 1 \\ -1 & -2 & 3 & 2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 6 \\ -1 \end{bmatrix}$$

¹Note that the minus sign appears in the real component because, when multiplying the two “last” parts, $i^2 = -1$. Note that the minus sign appears in the real component because, when multiplying the two “last” parts, $i^2 = -1$.

- Solve for a_1 , a_2 , b_1 , and b_2 using an augmented matrix and Gauss-Jordan elimination.

$$\left[\begin{array}{cccc|c} 2 & 1 & -1 & -1 & 3 \\ 3 & 2 & 1 & 2 & 7 \\ 1 & 1 & 2 & 1 & 6 \\ -1 & -2 & 3 & 2 & -1 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right]$$

$$\begin{bmatrix} a_1 \\ a_2 \\ b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 2 \\ -1 \end{bmatrix}$$

- From these four values, the original solutions $x_1 = a_1 + ib_1$ and $x_2 = a_2 + ib_2$ can be found.

$$x_1 = 1 + 2i$$

$$x_2 = 2 - i$$