

## Script 3

# Introducing a Continuum

### 3.1 Journal

10/20: **Axiom 1.** *A continuum is a nonempty set  $C$ .*

**Definition 3.1.** Let  $X$  be a set. An **ordering** on the set  $X$  is a subset  $<$  of  $X \times X$  with elements  $(x, y) \in <$  written as  $x < y$ , satisfying the following properties:

- a) (*Trichotomy*) For all  $x, y \in X$ , exactly one of the following holds:  $x < y$ ,  $y < x$ , or  $x = y$ .
- b) (*Transitivity*) For all  $x, y, z \in X$ , if  $x < y$  and  $y < z$ , then  $x < z$ .

**Remark 3.2.**

- a) In mathematics, “or” is understood to be inclusive unless stated otherwise. So in Definition 3.1a above, the word “exactly” is needed.
- b)  $x < y$  may also be written as  $y > x$ .
- c) By  $x \leq y$ , we mean  $x < y$  or  $x = y$ ; similarly for  $x \geq y$ .
- d) We often refer to elements of a continuum  $C$  as **points**.

**Axiom 2.** *A continuum  $C$  has an ordering  $<$ .*

**Definition 3.3.** If  $A \subset C$ , then a point  $a \in A$  is a **first** point of  $A$  if for every element  $x \in A$ , either  $a < x$  or  $a = x$ . Similarly, a point  $b \in A$  is called a **last** point of  $A$  if, for every  $x \in A$ , either  $x < b$  or  $x = b$ .

**Lemma 3.4.** *If  $A$  is a nonempty, finite subset of a continuum  $C$ , then  $A$  has a first and last point.*

**Lemma.** *Let  $A$  be a nonempty, finite set (i.e.,  $|A| = n$  for some  $n \in \mathbb{N}$ ), let  $a$  be any element of  $A$ , and let the set  $B = A \setminus \{a\}$ . Then  $|B| = n - 1$ .*

*Proof.* We first prove that  $|\{a\}| = 1$ . By Definition 1.33, to do so, it will suffice to find a bijection  $f : \{a\} \rightarrow [1]$ . Since  $[1] = \{1\}$  by Definition 1.29,  $f : \{a\} \rightarrow \{1\}$  defined by  $f(a) = 1$  is clearly such a bijection. We now demonstrate that  $B \cap \{a\} = (A \setminus \{a\}) \cap \{a\} = \emptyset$ . The previous two results combined with the fact that  $B \cup \{a\} = (A \setminus \{a\}) \cup \{a\} = A$  imply by Theorem 1.34b that  $|A| = |B| + |\{a\}|$ . It follows that  $n = |B| + 1$ , so  $|B| = n - 1$ .  $\square$

*Proof of Lemma 3.4.* We consider first points herein (the proof is symmetric for last points). If  $A$  is a finite set, then by Definition 1.30,  $|A| = n$  for some  $n \in \mathbb{N}$ . Thus, if we prove the claim for each  $n \in \mathbb{N}$  individually, we will have proven the claim. To prove a property pertaining to any natural number, we induct on  $n$ .

For the base case  $n = 1$ , there is only one element (which we may call  $a$ ) in  $A$ . Since  $a = a$ , i.e., “for every  $x \in A$ , either  $a < x$  or  $a = x$ ” is a true statement, it follows by Definition 3.3 that  $A$  has a first point. Now suppose inductively that we have proven the claim for  $n$ , i.e., we know that if  $A$  is a nonempty,

finite subset of a continuum  $C$  with  $|A| = n$ , then  $A$  has a first point. We wish to prove the same claim if  $|A| = n + 1$ . Let  $a$  be an arbitrary element of  $A$ , and consider the set  $B = A \setminus \{a\}$ . By the lemma,  $|B| = n$ . Consequently, the induction hypothesis applies and asserts that  $B$  has a first point  $a_0$ . Clearly,  $a_0$  is also an element of  $A$ , but it may or may not be the first point of  $A$  (the first point may now be  $a$ ). Since  $C$  has an ordering  $<$  (see Axiom 2), Definition 3.1 asserts that either  $a < a_0$ ,  $a_0 < a$ , or  $a = a_0$ . We now divide into three cases. If  $a < a_0$ , then since  $a_0 \leq x$  for all  $x \in A$  by Definition 3.3, Definition 3.1 implies that  $a \leq x$  for all  $x \in A$ . Thus, by Definition 3.3,  $a$  is the first point in  $A$ , and we have proven the claim for  $|A| = n + 1$  in this case. If  $a_0 < a$ , then it is still true that  $a_0 \leq x$  for all  $x \in A$ . This means by Definition 3.3 that  $a_0$  is still the first point in  $A$ , proving the claim for  $|A| = n + 1$  in this case. If  $a = a_0$ , then  $a \in B$ , contradicting the fact that  $B = A \setminus \{a\}$ , so we need not consider this final case. This closes the induction.  $\square$

**Theorem 3.5.** *Suppose that  $A$  is a set of  $n$  distinct points in a continuum  $C$ , or in other words,  $A \subset C$  has cardinality  $n$ . Then the symbols  $a_1, \dots, a_n$  may be assigned to each point of  $A$  so that  $a_1 < a_2 < \dots < a_n$ , i.e.,  $a_i < a_{i+1}$  for all  $1 \leq i \leq n - 1$ .*

*Proof.* We divide into two cases ( $|A| = 0$  and  $|A| \in \mathbb{N}$ ).

If  $|A| = 0$ , then the statements “the symbols  $a_1, \dots, a_n$  may be assigned to each point of  $A$ ” and “ $a_i < a_{i+1}$  for all  $1 \leq i \leq n - 1 = -1$ ” are both vacuously true.

If  $|A| \in \mathbb{N}$ , we induct on  $|A| = n$ . For the base case  $n = 1$ , denote the single element of  $A$  by  $a_1$ . Since  $a_i < a_{i+1}$  for all  $1 \leq i \leq n - 1 = 0$  is vacuously true, the base case holds. Now suppose inductively that we have proven the claim for  $n$ , i.e., for a set  $A \subset C$  satisfying  $|A| = n$ , the symbols  $a_1, \dots, a_n$  may be assigned to each point of  $A$  so that  $a_1 < a_2 < \dots < a_n$ . We now wish to prove the claim with regards to a set  $A \subset C$  with  $|A| = n + 1$ . By Lemma 3.4, there is a last point  $a_{n+1} \in A$ , which may be denoted as such (we will rigorously confirm this later). Since the set  $A \setminus \{a_{n+1}\}$  has cardinality  $n$  (see the lemma from Lemma 3.4), we have by the induction hypothesis that its  $n$  elements can be named  $a_1, \dots, a_n$  and ordered  $a_1 < a_2 < \dots < a_n$ . Clearly these  $n$  elements are elements of  $A$  and all that's left to do is determine where  $a_{n+1}$  fits into the established order. But by Definition 3.3,  $x \leq a_{n+1}$  for all  $x \in A$ , i.e.,  $x < a_{n+1}$  for all  $x \in A \setminus \{a_{n+1}\}$ . Consequently, as its name would suggest, it is true that  $a_1 < a_2 < \dots < a_n < a_{n+1}$ , as desired.  $\square$

**Definition 3.6.** If  $x, y, z \in C$  and either (i) both  $x < y$  and  $y < z$  or (ii) both  $z < y$  and  $y < x$ , then we say that  $y$  is **between**  $x$  and  $z$ .

**Corollary 3.7.** *Of three distinct points in a continuum, one must be between the other two.*

*Proof.* Let  $A$  be a subset of the described continuum containing the three distinct points. It follows by Theorem 3.5 that the symbols  $a_1, a_2, a_3$  may be assigned to each point of  $A$  so that  $a_1 < a_2 < a_3$ . Thus,  $a_1 < a_2$  and  $a_2 < a_3$ , so  $a_2$  is between  $a_1$  and  $a_3$  by Definition 3.6.  $\square$

**Axiom 3.** *A continuum  $C$  has no first or last point.*

**Definition 3.8.** We define an ordering on  $\mathbb{Z}$  by  $m < n$  if  $n = m + c$  for some  $c \in \mathbb{N}$ .

**Exercise 3.9.**

- a) Prove that with this ordering  $\mathbb{Z}$  satisfies Axioms 1-3.

*Proof.* Clearly,  $\mathbb{Z}$  is a nonempty set, so Axiom 1 is immediately satisfied.

Axiom 2 asserts that  $\mathbb{Z}$  must have an ordering  $<$ . As such, we must verify that the ordering given by Definition 3.8 satisfies the stipulations of Definition 3.1. To prove that  $<$  satisfies the trichotomy, it will suffice to show that for all  $x, y \in \mathbb{Z}$ , exactly one of the following holds:  $x < y$ ,  $y < x$ , or  $x = y$ . Let  $x, y$  be arbitrary elements of  $\mathbb{Z}$ . We divide into three cases. First, suppose for the sake of contradiction that  $x < y$  and  $y < x$ . By Definition 3.8, this implies that  $y = x + c$  and  $x = y + c'$  for some  $c, c' \in \mathbb{N}$ . Substituting, we have  $y = y + c' + c$ , or  $0 = c' + c$  by the cancellation law of addition. But since  $c', c \in \mathbb{N}$ , the closure of addition on  $\mathbb{N}$  implies that  $(c' + c) \in \mathbb{N}$ . Therefore,  $c' + c \neq 0$ , a contradiction. Second, suppose for the sake of contradiction that  $x < y$  and  $x = y$ . By Definition 3.8, this implies that  $y = x + c$  for some  $c \in \mathbb{N}$ . Substituting, we have  $y = y + c$ , or  $0 = c$  by the cancellation law of

addition. But since  $c \in \mathbb{N}$ ,  $c \neq 0$ , a contradiction. The proof of the third case ( $y < x$  and  $x = y$ ) is symmetric to that of the second case. To prove that  $<$  is transitive, it will suffice to show that for all  $x, y, z \in \mathbb{Z}$ , if  $x < y$  and  $y < z$ , then  $x < z$ . Let  $x, y, z$  be arbitrary elements of  $\mathbb{Z}$  for which it is true that  $x < y$  and  $y < z$ . By Definition 3.8, we have  $y = x + c$  and  $z = y + c'$  for some  $c, c' \in \mathbb{N}$ . Substituting, we have  $z = x + c + c'$ . Since  $(c + c') \in \mathbb{N}$  by the closure of addition on  $\mathbb{N}$ , Definition 3.8 implies that  $x < z$ .

Axiom 3 asserts that  $\mathbb{Z}$  must have no first or last point. Suppose for the sake of contradiction that  $\mathbb{Z}$  has some first point  $a$ . Then by Definition 3.3,  $a \leq x$  for every  $x \in \mathbb{Z}$ . However, under the closure of subtraction on  $\mathbb{Z}$ ,  $(a - 1) \in \mathbb{Z}$ . Since  $(a - 1) + 1 = a$ , Definition 3.8 asserts that  $a - 1 < a$ , a contradiction. The proof is symmetric for the last point.  $\square$

- b) Show that for any  $p = [\frac{a}{b}] \in \mathbb{Q}$ , there is some  $(a_1, b_1) \in p$  with  $0 < b_1$ .

*Proof.* Let  $[\frac{a}{b}]$  be an arbitrary element of  $\mathbb{Q}$ . It follows by Definition 2.5 that  $(a, b) \in X$ . Since we also have  $(a, b) \sim (a, b)$  by Exercise 2.2e, Definition 2.5 implies that  $(a, b) \in [\frac{a}{b}]$ . By the trichotomy (see Definition 3.1) of the ordering defined by Definition 3.8 on  $\mathbb{Z}$  which Exercise 3.9a just proved is valid, we have  $0 < b$ ,  $b < 0$ , or  $0 = b$ . We divide into three cases. First, suppose that  $0 < b$ . Then  $(a, b)$  is an element  $(a_1, b_1) \in [\frac{a}{b}]$  for which  $0 < b_1$ , and we are done. Second, suppose that  $b < 0$ . Since  $(-a)(b) = (-b)(a)$ , we have by the definition of  $\sim$  that  $(-a, -b) \sim (a, b)$ . Additionally, we have by the closure of integer multiplication that  $-a, -b \in \mathbb{Z}$ , and since  $b \neq 0$  by Exercise 2.2e,  $-b \neq 0$ . Thus, by Exercise 2.2e,  $(-a, -b) \in X$ . This coupled with the previously proven fact that  $(-a, -b) \sim (a, b)$  implies by Definition 2.5 that  $(-a, -b) \in [\frac{a}{b}]$ . Now recall that  $b < 0$  by hypothesis, so we may use Definition 3.8 to see that  $b + c = 0$  for some  $c \in \mathbb{N}$ . It follows that  $-(b + c) = 0$ , i.e.,  $-b - c = 0$ , i.e.,  $-b = 0 + c$ , meaning that  $0 < -b$  by Definition 3.8. Thus,  $(-a, -b)$  is an element  $(a_1, b_1) \in [\frac{a}{b}]$  for which  $0 < b_1$ . Third, suppose that  $b = 0$ . But this contradicts Exercise 2.2e which asserts that  $b \neq 0$ , so we need not consider this case.  $\square$

- c) Define an ordering  $<_{\mathbb{Q}}$  on  $\mathbb{Q}$  as follows. For  $p, q \in \mathbb{Q}$ , let  $(a_1, b_1) \in p$  be such that  $0 < b_1$  and let  $(a_2, b_2) \in q$  be such that  $0 < b_2$ . Then we define  $p <_{\mathbb{Q}} q$  if  $a_1 b_2 < a_2 b_1$ . Show that  $<_{\mathbb{Q}}$  is a well-defined relation on  $\mathbb{Q}$ .

*Proof.* For the relation  $<_{\mathbb{Q}}$  to be well-defined, Definition 3.1 tells us that it must satisfy the trichotomy and be transitive.

To prove that  $<_{\mathbb{Q}}$  satisfies the trichotomy, it will suffice to show that for all  $p, q \in \mathbb{Q}$ , exactly one of the following holds:  $p <_{\mathbb{Q}} q$ ,  $q <_{\mathbb{Q}} p$ , or  $p = q$ . Let  $p, q$  be arbitrary elements of  $\mathbb{Q}$ , let  $(a, b) \in p$  be such that  $0 < b$  (we know that such an element exists by Exercise 3.9b), and let  $(c, d) \in q$  be such that  $0 < d$ . We divide into three cases. First, suppose for the sake of contradiction that  $p <_{\mathbb{Q}} q$  and  $q <_{\mathbb{Q}} p$ . Then  $ad < bc$  and  $cb < da$  by the definition of  $<_{\mathbb{Q}}$ . But this violates the trichotomy known to hold for the ordering  $<$  on the integers by Exercise 3.9a, a contradiction. Second, suppose for the sake of contradiction that  $p <_{\mathbb{Q}} q$  and  $p = q$ . By the definition of  $<_{\mathbb{Q}}$ ,  $ad < bc$ . Additionally, by Exercise 2.6,  $(a, b) \sim (c, d)$ , implying by Exercise 2.2e that  $ad = bc$ . But once again, the simultaneous results that  $ad < bc$  and  $ad = bc$  violate the trichotomy of the integers, a contradiction. The proof of the third case is symmetric to that of the second.

To prove that  $<_{\mathbb{Q}}$  is transitive, it will suffice to show that for all  $p, q, r \in \mathbb{Q}$ , if  $p <_{\mathbb{Q}} q$  and  $q <_{\mathbb{Q}} r$ , then  $p <_{\mathbb{Q}} r$ . Let  $p, q, r$  be arbitrary elements of  $\mathbb{Q}$  for which it is true that  $p <_{\mathbb{Q}} q$  and  $q <_{\mathbb{Q}} r$ , let  $(a, b) \in p$  be such that  $0 < b$ , let  $(c, d) \in q$  be such that  $0 < d$ , and let  $(e, f) \in r$  such that  $0 < f$ . By the definition of  $<_{\mathbb{Q}}$ , we have  $ad < bc$  and  $cf < de$ . By Definition 3.8, this implies that  $bc = ad + g$  and  $de = cf + g'$  for some  $g, g' \in \mathbb{N}$ . Thus, by the multiplicative property of equality,

$$bcf = adf + gf \qquad bde = bcf + bg'$$

so that by the additive property of equality

$$\begin{aligned} bcf + bde &= adf + gf + bcf + bg' \\ bde &= adf + gf + bg' \end{aligned}$$

Since  $0 < b$  and  $0 < f$  for  $b, f \in \mathbb{Z}$ , we know that  $b, f \in \mathbb{N}$ . Since  $g, g' \in \mathbb{N}$  as well by hypothesis, the closures of multiplication and subsequently of addition for  $\mathbb{N}$  guarantee that  $(gf + bg') \in \mathbb{N}$ . Thus, by Definition 3.8,  $adf < bde$ . Now suppose for the sake of contradiction that  $af \not< be$ . By Definition 3.1, this implies that either  $af = be$  or  $be < af$ . We divide into two cases. First, suppose that  $af = be$ . Then by the multiplicative property of equality,  $adf = bde$ , contradicting the fact that  $adf < bde$ . Second, suppose that  $be < af$ . Then by Definition 3.8,  $af = be + g$  for some  $g \in \mathbb{N}$ . Thus,  $adf = bde + gd$  where  $gd \in \mathbb{N}$  by the closure of multiplication for the natural numbers ( $d \in \mathbb{N}$  since  $d \in \mathbb{Z}$  and  $0 < d$  by hypothesis). Thus,  $bcd < adf$  by Definition 3.8, contradicting the fact that  $adf < bde$ . Therefore, it must be true that  $af < be$ . It follows by the definition of  $<_{\mathbb{Q}}$  that  $p <_{\mathbb{Q}} r$ .  $\square$

d) Show that  $\mathbb{Q}$  with the ordering  $<_{\mathbb{Q}}$  satisfies Axioms 1-3.

*Proof.* Clearly,  $\mathbb{Q}$  is a nonempty set, so Axiom 1 is immediately satisfied.

By Exercise 3.9c,  $\mathbb{Q}$  has an ordering, so Axiom 2 is satisfied.

Axiom 3 asserts that  $\mathbb{Q}$  must have no first or last point. Suppose for the sake of contradiction that  $\mathbb{Q}$  has some first point  $p$ . Then by Definition 3.3,  $p <_{\mathbb{Q}} x$  or  $p = x$  for all  $x \in \mathbb{Q}$ . Let  $(a, b) \in p$  be such that  $0 < b$  (see Exercise 3.9b). Under the closure of integer subtraction,  $(a - 1) \in \mathbb{Z}$ , so  $[\frac{a-1}{b}] \in \mathbb{Q}$ . Since  $ba = ba - b + b = b(a - 1) + b$  where  $b \in \mathbb{N}$  since  $b \in \mathbb{Z}$  and  $0 < b$ , Definition 3.8 implies that  $(a - 1)b < ba$ . It follows by the definition of  $<_{\mathbb{Q}}$  that  $[\frac{a-1}{b}] <_{\mathbb{Q}} [\frac{a}{b}] = p$ , a contradiction. The argument is symmetric for the last point.  $\square$

**Definition 3.10.** If  $a, b \in C$  and  $a < b$ , then the set of points between  $a$  and  $b$  is called a **region** and denoted by  $\underline{ab}$ .

**Remark 3.11.** One often sees the notation  $(a, b)$  for regions. We will reserve the notation  $(a, b)$  for ordered pairs in a product  $A \times B$ . These are very different things.

**Theorem 3.12.** If  $x$  is a point of a continuum  $C$ , then there exists a region  $\underline{ab}$  such that  $x \in \underline{ab}$ .

*Proof.* Let  $x$  be an arbitrary point in a continuum  $C$ . By Axiom 2,  $C$  has an ordering  $<$ , which we will frequently make use of throughout the remainder of this proof. By Axiom 3,  $C$  has no first or last points, so it cannot be true that  $x \leq y$  for all  $y \in C$ , nor can it be true that  $x \geq y$  for all  $y \in C$ . This implies that there exists an  $a \in C$  such that  $a < x$  and that there exists a  $b \in C$  such that  $b > x$ . Since  $a < x$  and  $x < b$ , Definition 3.6 implies that  $x$  is between  $a$  and  $b$ . Note also that by Definition 3.1 (specifically transitivity),  $a < b$ . Therefore, since  $a, b \in C$ ,  $a < b$ , and  $x$  is between  $a$  and  $b$ , Definition 3.10 implies that  $x \in \underline{ab}$ .  $\square$

**Definition 3.13.** Let  $A$  be a subset of a continuum  $C$ . A point  $p$  of  $C$  is called a **limit point** of  $A$  if every region  $R$  containing  $p$  has nonempty intersection with  $A \setminus \{p\}$ . Explicitly, this means:

$$\text{for every region } R \text{ with } p \in R, \text{ we have } R \cap (A \setminus \{p\}) \neq \emptyset.$$

Notice that we do not require that a limit point  $p$  of  $A$  be an element of  $A$ . We will use the notation  $LP(A)$  to denote the set of limit points of  $A$ .

**Theorem 3.14.** If  $p$  is a limit point of  $A$  and  $A \subset B$ , then  $p$  is a limit point of  $B$ .

**Lemma.** Let  $A, B, C$  be sets such that  $A \subset B$ . Then  $A \cap C \subset B \cap C$ .

*Proof.* Let  $x$  be an arbitrary element of  $A \cap C$ . By Definition 1.6, this implies that  $x \in A$  and  $x \in C$ . Since  $x \in A$  and  $A \subset B$ , Definition 1.3 implies that  $x \in B$ . Thus,  $x \in B$  and  $x \in C$ , so  $x \in B \cap C$  by Definition 1.6.  $\square$

*Proof.* To prove that a limit point  $p$  of  $A \subset B$  is a limit point of  $B$ , Definition 3.13 tells us that it will suffice to show that for every region  $R$  with  $p \in R$ , we have  $R \cap (B \setminus \{p\}) \neq \emptyset$ . Let  $p$  be a limit point of  $A$ , and let  $R$  be an arbitrary region with  $p \in R$ . Then by Definition 3.13, we have  $R \cap (A \setminus \{p\}) \neq \emptyset$ . Thus, by Definition 1.8, there is an element  $x \in R \cap (A \setminus \{p\})$ . Since  $A \setminus \{p\} \subset B \setminus \{p\}$  (because  $A \subset B$  and  $\{p\} = \{p\}$ ), it follows by the lemma that  $R \cap (A \setminus \{p\}) \subset R \cap (B \setminus \{p\})$ . Consequently, by Definition 1.3, the previously referenced object  $x \in R \cap (A \setminus \{p\})$  is also an element of  $R \cap (B \setminus \{p\})$ . Thus, by Definition 1.8,  $R \cap (B \setminus \{p\}) \neq \emptyset$ , as desired.  $\square$