

Chapter 1

Multilinear Algebra

1.1 Notes

3/28: • Motivation for the course and an overview of Guillemin and Haine (2018).

3/30: • Plan:

– More (multi)linear algebra.

• Dual spaces.

• Let V be an n -dimensional real vector space.

• **Hom** (V, \mathbb{R}): The set of all homomorphisms (i.e., linear maps) from V to \mathbb{R} . *Also known as V^* .*

• **Dual basis** (for V^*): The set of linear transformations from V to \mathbb{R} defined by

$$e_j \mapsto \begin{cases} 1 & j = i \\ 0 & j \neq i \end{cases}$$

where e_1, \dots, e_n is a basis of V . *Denoted by e_1^*, \dots, e_n^* .*

• Check: e_1^*, \dots, e_n^* are a basis for V^* .

– Are they linearly independent? Let $c_1 e_1^* + \dots + c_n e_n^* = 0 \in \text{Hom}(V, \mathbb{R})$. Then

$$c_i = (c_1 e_1^* + \dots + c_n e_n^*)(e_i) = 0 \in \mathbb{R}$$

as desired.

– Span? Let $\varphi \in \text{Hom}(V, \mathbb{R})$. Then we can verify that

$$\varphi(e_1) e_1^* + \dots + \varphi(e_n) e_n^* = \varphi$$

■ We prove this by verifying the previous statement on the basis of V (if two linear transformations have the same action on the basis of a vector space, they are equal).

• With a choice of basis for V , we obtain an isomorphism $\varepsilon : V \rightarrow V^*$ with the mapping $e_i \mapsto e_i^*$ for all i .

• The dual space is known as such because $(V^*)^* \cong V$, where \cong is **canonical** (no choice of basis is needed).

• One more property of dual spaces: **functoriality**.

- Given a linear transformation $A : V \rightarrow W$, we know that $A^* : W^* \rightarrow V^*$ where A^* is the transpose of A . In particular, if $\varphi \in W^*$, then $\varphi \circ A : V \rightarrow \mathbb{R}$.
- Claim: A^* is linear.
- **Functoriality:** If $A : V \rightarrow W$ and $B : W \rightarrow U$, then $B^* : U^* \rightarrow W^*$ and $A^* : W^* \rightarrow V^*$. The functoriality statement is that $(B \circ A)^* = A^* \circ B^*$.
- A^* is the **pullback** (or transpose) of A .
- Let v_1, \dots, v_n be a basis for V and w_1, \dots, w_m be a basis for W . Then $[A]_{v_1, \dots, v_n}^{w_1, \dots, w_m} = A$ is the matrix of the linear transformation A with respect to these bases. Then if v_1^*, \dots, v_n^* and w_1^*, \dots, w_m^* are the corresponding dual bases, then $[A^*]_{v_1^*, \dots, v_n^*}^{w_1^*, \dots, w_m^*} = A^T$. We can and should verify this for ourselves.
- This is over the real numbers, so A^* is just the transpose because there are no complex numbers of which to take the conjugate!
- A generalization: Tensors.
- **k -tensor:** A **multilinear** map

$$T : \underbrace{V \times \cdots \times V}_{k \text{ times}} \rightarrow \mathbb{R}$$

- **Multilinear** (map T): A function T such that

$$\begin{aligned} T(v_1, \dots, v_i^1 + v_i^2, \dots, v_k) &= T(v_1, \dots, v_i^1, \dots, v_k) + T(v_1, \dots, v_i^2, \dots, v_k) \\ T(v_1, \dots, \lambda v_i, \dots, v_k) &= \lambda T(v_1, \dots, v_i, \dots, v_k) \end{aligned}$$

for all $(v_1, \dots, v_k) \in V^k$.

- The determinant is an n -tensor!
- 1-tensors are just covectors.
- $L^k(V)$: The vector space of all k -tensors on V .
- Calculating $\dim L^k(V)$. (Answer not given in this class.)
- Let $A : V \rightarrow W$. Then $A^* : L^k(W) \rightarrow L^k(V)$.
 - Check $(A \circ B)^* = B^* \circ A^*$.
- **Multi-index of n of length k :** A k -tuple (i_1, \dots, i_k) where each $i_j \in \mathbb{N}$ satisfies $1 \leq i_j \leq n$ ($j = 1, \dots, k$). Denoted by I .
- Let e_1, \dots, e_n be a basis for V .
- **Tensor product** (of $T_1 \in L^k(V)$, $T_2 \in L^l(V)$): The function from V^{k+l} to \mathbb{R} defined by

$$(v_1, \dots, v_{k+l}) \mapsto T_1(v_1, \dots, v_k) T_2(v_{k+1}, \dots, v_{k+l})$$

Denoted by $T_1 \otimes T_2$.

- Claims:
 1. $T_1 \otimes T_2 \in L^{k+l}(V)$.
 2. $A^*(T_1 \otimes T_2) = A^*(T_1) \otimes A^*(T_2)$.
- e_I^* : The function $e_{i_1}^* \otimes \cdots \otimes e_{i_k}^*$, where $I = (i_1, \dots, i_k)$ is a multi-index of n of length k .
- Claim: Letting I range over all n^k multi-indices of n of length k , the e_I^* are a basis for $L^k(V)$.
- If $V = \mathbb{R}$, then $V = \mathbb{R}e_1$. If $V = \mathbb{R}^2$, then $V = \mathbb{R}e_1 \oplus \mathbb{R}e_2$.
- We know that $L^1(V) = V^* = \mathbb{R}e_1^*$. Thus, $e_1^* \otimes e_2^* : V \times V \rightarrow \mathbb{R}$. Thus, for example,

$$(e_1^* \otimes e_2^*)((1, 2), (3, 4)) = e_1^*(1, 2) \cdot e_2^*(3, 4) = 1 \cdot 4 = 4$$

1.2 Chapter 1: Multilinear Algebra

From Guillemin and Haine (2018).

- 3/31:
- Guillemin and Haine (2018) defines real vector spaces, the operations on them, their basic properties, and the zero vector.
 - **Linearly independent** (vectors v_1, \dots, v_k): A finite set of vectors $v_1, \dots, v_k \in V$ such that the map from \mathbb{R}^k to V defined by $(c_1, \dots, c_k) \mapsto c_1 v_1 + \dots + c_k v_k$ is injective.
 - **Spanning** (vectors v_1, \dots, v_k): We require that the above map is surjective.
 - Guillemin and Haine (2018) defines basis, finite-dimensional vector space, dimension, subspace, linear map, and kernel.
 - **Image** (of $A : V \rightarrow W$): The range space of A , a subspace of W . Also known as $\mathbf{im}(A)$.
 - Guillemin and Haine (2018) defines the matrix of a linear map.
 - **Inner product** (on V): A map $B : V \times V \rightarrow \mathbb{R}$ with the following three properties.

– *Bilinearity*: For vectors $v, v_1, v_2, w \in V$ and $\lambda \in \mathbb{R}$, we have

$$B(v_1 + v_2, w) = B(v_1, w) + B(v_2, w)$$

and

$$B(\lambda v, w) = \lambda B(v, w)$$

– *Symmetry*: For vectors $v, w \in V$, we have $B(v, w) = B(w, v)$.

– *Positivity*: For every vector $v \in V$, we have $B(v, v) \geq 0$. Moreover, if $v \neq 0$, then $B(v, v) > 0$.

- **W-coset**: A set of the form $\{v + w \mid w \in W\}$, where W is a subspace V and $v \in V$. Denoted by $v + W$.
 - If $v_1 - v_2 \in W$, then $v_1 + W = v_2 + W$.
 - It follows that the distinct W -cosets decompose V into a disjoint collection of subsets of V .
- **Quotient space** (of V by W): The set of distinct W -cosets in V , along with the following definitions of vector addition and scalar multiplication.

$$(v_1 + W) + (v_2 + W) = (v_1 + v_2) + W$$

$$\lambda(v + W) = (\lambda v) + W$$

Denoted by V/W .

- **Quotient map**: The linear map $\pi : V \rightarrow V/W$ defined by

$$\pi(v) = v + W$$

– π is surjective.

– Note that $\ker(\pi) = W$ since for all $w \in W$, $\pi(w) = w + W = 0 + W$, which is the zero vector in V/W .

- If V, W are finite dimensional, then

$$\dim(V/W) = \dim(V) - \dim(W)$$

- Proposition 1.2.9: Let $A : V \rightarrow U$ be a linear map. If $W \subset \ker(A)$, then there exists a unique linear map $A^\sharp : V/W \rightarrow U$ with the property that $A = A^\sharp \circ \pi$, where $\pi : V \rightarrow V/W$ is the quotient map.
 - This proposition rephrases in terms of quotient spaces the fact that if $w \in W$, then $A(v+w) = Av$.

- **Dual space** (of V): The set of all linear functions $\ell : V \rightarrow \mathbb{R}$, along with the following definitions of vector addition and scalar multiplication.

$$(\ell_1 + \ell_2)(v) = \ell_1(v) + \ell_2(v) \qquad (\lambda\ell)(v) = \lambda \cdot \ell(v)$$

Denoted by V^* .

- **Dual basis** (of e_1, \dots, e_n a basis of V): The basis of V^* consisting of the n functions that take every $v = c_1e_1 + \dots + c_ne_n$ to one of the c_i . Denoted by e_1^*, \dots, e_n^* . Given by

$$e_i^*(v) = c_i$$

for all $v \in V$.

- Claim 1.2.12: If V is an n -dimensional vector space with basis e_1, \dots, e_n , then e_1^*, \dots, e_n^* is a basis of V^* .

Proof. We will first prove that e_1^*, \dots, e_n^* spans V^* . Let $\ell \in V^*$ be arbitrary. Set $\lambda_i = \ell(e_i)$ for all $i \in [n]$. Define $\ell' = \sum_{i=1}^n \lambda_i e_i^*$. Then

$$\ell'(e_j) = \sum_{i=1}^n \lambda_i e_i^*(e_j) = \lambda_j \cdot 1 = \ell(e_j)$$

for all $j \in [n]$. Therefore, since ℓ, ℓ' take identical values on the basis of V , $\ell = \ell'$, as desired.

We now prove that e_1^*, \dots, e_n^* spans V^* . Let $\sum_{i=1}^n \lambda_i e_i^* = 0$. Then for all $j \in [n]$,

$$\lambda_j = \left(\sum_{i=1}^n \lambda_i e_i^* \right)(e_j) = 0$$

as desired. □

- **Transpose** (of A): The map from W^* to V^* defined by $\ell \mapsto A^*\ell = \ell \circ A$ for all $\ell \in W^*$.
- Claim 1.2.15: If e_1, \dots, e_n is a basis of V , f_1, \dots, f_m is a basis of W , e_1^*, \dots, e_n^* and f_1^*, \dots, f_m^* are the corresponding dual bases, and $[a_{i,j}]$ is the $m \times n$ matrix of A with respect to $\{e_i\}, \{f_i\}$, then the linear map A^* is defined in terms of $\{f_i^*\}, \{e_i^*\}$ by the transpose matrix $(a_{j,i})$.