

## Chapter 3

# Determinants

- 9/29:
- The determinant, geometrically, is the volume of the object (in  $\mathbb{R}^3$ ) you get when you take linear combinations of the vectors.
  - In 2D:
    - Let  $v_1, v_2$  be two vectors. Put tail to tail and forming a parallelogram, the determinant of the matrix  $(v_1, v_2)$  is the area of said parallelogram.
    - Linearity 1:  $D(av_1, v_2, \dots, v_n) = aD(v_1, \dots, v_n)$  is the same as saying that if you stretch one vector by  $a$ , you scale up the area by that much, too.
    - Linearity 2:  $D(v_1, \dots, v_{k+} + v_{k-}, \dots, v_n) = D(-) + D(+)$ .
    - Antisymmetry:  $D(v_1, \dots, v_k, \dots, v_j, \dots, v_n) = -D(v_1, \dots, v_j, \dots, v_k, \dots, v_n)$ . Interchanging columns flips the sign of the determinant.
    - Basis:  $D(e_1, \dots, e_n) = 1$ .
  - Determinant: Denoted by  $D(v_1, \dots, v_n)$ , where  $(v_1, \dots, v_n)$  is an  $n \times n$  matrix.
- 10/1:
- Consider an  $n \times n$  matrix  $A$  consisting of  $n$  columns containing vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n \in \mathbb{R}^n$ .
    - $D(A)$  is the volume of the solid  $V = \sum_{i=1}^n \alpha_i v_i$ .
    - $D(\mathbf{e}_1, \dots, \mathbf{e}_n) = 1$ .




Figure 3.1: Visualizing properties of determinants.

- Basic properties of the determinant.
  - If  $A$  has a zero column, then  $\det A = 0$ : Scalar property.
  - If  $A$  has two equal columns, then  $\det A = 0$ : Multiply one by minus and add.

- If  $A$  has a column which is a multiple of another, then  $\det A = 0$ : Pull out the multiple and then you have the previous one.
- If columns are linearly dependent, then  $\det A = 0$ : Decompose it into sums, split, add back up with previous properties.
- The determinant is preserved under column reduction.
- $\det A^T = \det A$ : Put everything in rref.
- If  $A$  is not invertible, then  $\det A = 0$  (not invertible implies linearly dependent columns, implies  $\det A = 0$ ).
- $\det(AB) = \det A \det B$ .
- Determinant of...
  - A diagonal matrix: The product of the diagonal entries (pull out the terms, and then note that the remaining identity matrix has determinant 1).
  - An upper triangular matrix: The product of the diagonal entries (column reduction to make it into a diagonal matrix, and then the property above).