

## Chapter 3

# Numerical Sequences and Series

### 3.1 Notes

- 11/8:      • Any bounded sequence in  $\mathbb{R}^k$  has a convergent subsequence.
- 11/10:     • Read and understand the section about Cauchy sequences converging and the sup/inf.

### 3.2 Chapter 3: Numerical Sequences and Series

*From Rudin (1976).*

- 11/7:      • Convergence of sequences is relative.
- For example, the sequence  $1/n$  for  $n = 1, 2, \dots$  converges in  $\mathbb{R}$ , but not in  $(0, \infty)$ .
- **Range** (of  $\{p_n\}$ ): The set of all points  $p_n$ .
- This definition squares nicely with the formal definition of a sequence as a function  $p$  defined on  $\mathbb{N}$ .
- Theorem 3.6a: If  $\{p_n\}$  is a sequence in a compact metric space  $X$ , then some subsequence of  $\{p_n\}$  converges to a point of  $X$ .
- Theorem 3.7: The subsequential limits of a sequence  $\{p_n\}$  in a metric space  $X$  form a closed subset of  $X$ .
- **Diameter** (of  $E$ ): The supremum of the set

$$S = \{d(p, q) : p, q \in E\}$$

where  $E$  is a nonempty subset of a metric space  $X$ . Denoted by **diam**  $E$ .

- Theorem 3.10:
- (a) If  $\bar{E}$  is the closure of a set  $E$  in a metric space  $X$ , then
$$\text{diam } \bar{E} = \text{diam } E$$
  - (b) If  $K_n$  is a sequence of compact sets in  $X$  such that  $K_n \supset K_{n+1}$  ( $n = 1, 2, 3, \dots$ ) and if  $\lim_{n \rightarrow \infty} \text{diam } K_n = 0$ , then  $\bigcap_1^\infty K_n$  consists of exactly one point.
- **Complete** (metric space): A metric space in which every Cauchy sequence converges.
- All compact metric spaces and all Euclidean spaces are complete.

- The metric space  $(\mathbb{Q}, |x - y|)$  is not complete.
- **Monotonically increasing** (sequence  $\{s_n\}$ ): A sequence  $\{s_n\}$  of real numbers such that  $s_n \leq s_{n+1}$  for each  $n \in \mathbb{N}$ .
- **Monotonically decreasing** (sequence  $\{s_n\}$ ): A sequence  $\{s_n\}$  of real numbers such that  $s_n \geq s_{n+1}$  for each  $n \in \mathbb{N}$ .
- **Monotonic sequences**: The class of all sequences that are either monotonically increasing or monotonically decreasing.
- **Upper limit** (of  $\{s_n\}$ ): The supremum of the set  $E$  of all subsequential limits of  $\{s_n\}$ . *Denoted by  $s^*$ ,  $\limsup_{n \rightarrow \infty} s_n$ .*
- **Lower limit** (of  $\{s_n\}$ ): The infimum of the set  $E$  of all subsequential limits of  $\{s_n\}$ . *Denoted by  $s_*$ ,  $\liminf_{n \rightarrow \infty} s_n$ .*
- Theorem 3.17: Let  $\{s_n\}$  be a sequence of real numbers. Then  $s^*$  has (and is the only number to have both of) the following two properties.
  - (a)  $s^* \in E$ .
  - (b) If  $x > s^*$ , then there is an integer  $N$  such that  $n \geq N$  implies  $s_n < x$ .

An analogous result holds for  $s_*$ .

11/8:

- Series are defined in terms of sequences. Moreover, sequences can be defined in terms of series: Let  $a_1 = s_1$ ,  $a_n = s_n - s_{n-1}$  ( $n \in \mathbb{N} + 1$ ). Thus, every theorem about sequences can be stated in terms of series and vice versa, but it is nevertheless useful to consider both concepts (Rudin, 1976, p. 59).
- Theorem 3.27: Suppose  $\{a_n\}$  is a monotonically decreasing sequence of nonnegative terms. Then the series  $\sum_{n=1}^{\infty} a_n$  converges if and only if the series

$$\sum_{k=0}^{\infty} 2^k a_{2^k} = a_1 + 2a_2 + 4a_4 + 8a_8 + \cdots$$

converges.

- Theorem 3.29: If  $p > 1$ ,

$$\sum_{n=2}^{\infty} \frac{1}{n(\log n)^p}$$

converges; if  $p \leq 1$ , the series diverges.

- Note that  $\log n = \ln n$ .
- Note that we sum from  $n = 2$  since  $\log 1 = 0$ .
- **e**: The number

$$e = \sum_{n=0}^{\infty} \frac{1}{n!}$$

- Theorem 3.31:  $\lim_{n \rightarrow \infty} (1 + 1/n)^n = e$ .
- Theorem 3.32:  $e$  is irrational.
- Theorem 3.39: Given the power series  $\sum c_n z^n$ , put

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|} \qquad R = \frac{1}{\alpha}$$

(If  $\alpha = 0$ , let  $R = +\infty$ ; if  $\alpha = +\infty$ , let  $R = 0$ .) Then  $\sum c_n z^n$  converges if  $|z| < R$  and diverges if  $|z| > R$ .

- **Radius of convergence** (of a power series): The number  $R$  defined by Theorem 3.39.
- Theorem 3.41 (partial summation formula): Given two sequence  $\{a_n\}, \{b_n\}$ , put

$$A_n = \begin{cases} \sum_{k=0}^n a_k & n \geq 0 \\ 0 & n = -1 \end{cases}$$

Then if  $0 \leq p \leq q$ , we have

$$\sum_{n=p}^q a_n b_n = \sum_{n=p}^{q-1} A_n (b_n - b_{n+1}) + A_q b_q - A_{p-1} b_p$$

- **Product** (of  $\sum a_n, \sum b_n$ ): The series  $\sum c_n$  defined by

$$c_n = \sum_{k=0}^n a_k b_{n-k}$$

for each  $n = 0, 1, 2, \dots$

- We motivate this definition by noting that if  $\sum c_n$  is the product of  $\sum a_n, \sum b_n$ , then

$$\sum_{n=0}^{\infty} a_n z^n \cdot \sum_{n=0}^{\infty} b_n z^n = \sum_{n=0}^{\infty} c_n z^n$$

- Setting  $z = 1$  then yields the given definition.

- The product of two convergent series may diverge. However...
- Theorem 3.50 (by Mertens): Suppose (a)  $\sum_{n=0}^{\infty} a_n$  converges absolutely, (b)  $\sum_{n=0}^{\infty} a_n = A$ , (c)  $\sum_{n=0}^{\infty} b_n = B$ , and (d)  $\sum_{n=0}^{\infty} c_n$  is the product of  $\sum_{n=0}^{\infty} a_n$  and  $\sum_{n=0}^{\infty} b_n$ . Then

$$\sum_{n=0}^{\infty} c_n = AB$$

- Theorem 3.51 (by Abel): If  $\sum a_n, \sum b_n, \sum c_n$  converge to  $A, B, C$ , respectively, and  $\sum c_n$  is the product of  $\sum a_n, \sum b_n$ , then  $C = AB$ .
- **Rearrangement** (of  $\sum a_n$ ): A series  $\sum a'_n$  defined by  $a'_n = a_{k_n}$  for each  $n \in \mathbb{N}$ , where  $\{k_n\}$  is a sequence in which every positive integer appears once and only once (that is,  $\{k_n\}$  is a 1-1 function from  $\mathbb{N}$  onto  $\mathbb{N}$ ).
- Theorem 3.54: Let  $\sum a_n$  be a series of real number which converges, but not absolutely. Suppose  $-\infty \leq \alpha \leq \beta \leq \infty$ . Then there exists a rearrangement  $\sum a'_n$  with partial sums  $s'_n$  such that

$$\liminf_{n \rightarrow \infty} s'_n = \alpha \qquad \limsup_{n \rightarrow \infty} s'_n = \beta$$

- Theorem 3.55: If  $\sum a_n$  is a series of complex numbers which converges absolutely, then every rearrangement of  $\sum a_n$  converges, and they all converge to the same sum.