

Week 8

???

8.1 Linear Algebra Review and Rational Canonical Form

2/20:

- Nori's change of heart.
 - We've all seen linear algebra; thus, we'll speedrun it and then do exterior algebra and determinants. That's where we'll finish.
- The following is part 1 of a linear algebra course.
- Let F be a field.
- **Vector space:** An F -module.
- **Linearly independent** (subset $S \subset V$): Same definition we're familiar with.
- **Spanning** (subset $S \subset V$): A subset S of V that is a set of generators of V .
- S is a **basis** implies that S generates V and is linearly independent.
- Every linearly independent subset of V can be extended to a basis.
- Every spanning set S contains a basis.
 - Any maximal linearly independent subset of S is a basis.
- S_1, S_2 are bases for V implies that $|S_1| = |S_2|$.
 - The replacement theorem in Dummit and Foote (2004) is a good way to prove this.
- We are now done with part 1; this is part 2 of a linear algebra course.
- Let $T : V \rightarrow V$ be a linear transformation.
- Let A be a ring. What is an $A[X]$ -module M ?
 - It is an abelian group $(M, +)$ and a ring homomorphism $\rho : A[X] \rightarrow \text{End}(M, +)$.
 - Since $A \hookrightarrow A[X]$, $\rho|_A$ turns M into an A -module.
 - Since $aX = Xa$, $\rho(a)\rho(X) = \rho(X)\rho(a)$.
 - But since we consider M to be a module, we write $a := \rho(a)$: Thus, $a\rho(X)m = \rho(X)am$ for all $m \in M$.
 - Note that $\rho(X) \in \text{End}_A(M)$ (which is the set of all A -module homomorphisms).
 - Additionally, $\rho(X) : M \rightarrow M$ is an A -module homomorphism.

- Put $\rho(X) = T$. Thus, an $A[X]$ -module is a pair (M, T) , where M is an A -module and $T \in \text{End}_A(M)$.
- Conversely, such (M, T) gives rise to an $A[X]$ -module.
 - In particular, the action is

$$\left(\sum_{n=0}^{\ell} a_n X^n \right) m = \sum_{n=0}^{\ell} a_n T^n m$$

- Take $A = F$ a field concerned with (V, T) where V is any F -vector space and $T : V \rightarrow V$ is a linear transformation.
 - This induces a module over $F[X]$.
- V finite dimensional induces $\rho : F[X] \rightarrow \text{End}_F(V) \cong M_n(F)$ defined by $X \mapsto T$.
 - $\rho(X) = T$ and $\rho(c) = c$ for all $c \in F$.
- Let n^2 be the dimension of the F -vector space??
- Then $\ker(\rho) = (f)$ for some f be monic of degree $d \leq n^2$.

$$\begin{array}{ccc} F[X] & \xrightarrow{\rho} & \text{End}_F(V) \\ \downarrow & \nearrow \bar{\rho} & \\ F[X]/(f) & & \end{array}$$

Figure 8.1: $F[X]$ -module actions.

- We have the constraint on the degree of f by the isomorphism from Lecture 3.1.
- **Minimal polynomial** (of T): The polynomial f that generates $\ker(\rho)$.
 - In particular, V is a torsion $F[X]$ -module $(f \cdot g)$.
- **Cyclic vector**: A vector $v \in V$ belonging to (V, T) such that v, Tv, T^2v, \dots spans V .
- Assume $v, Tv, T^2v, \dots, T^{k-1}v$ are linearly independent, but $v, Tv, \dots, T^k v$ are not.
 - Then

$$T^k v = a_0 v + a_1 Tv + \dots + a_{k-1} T^{k-1} v$$
 where all $a_i \in F$ and not all $a_i = 0$.
 - It follows that $T^m k \in \langle v, Tv, \dots, T^{k-1}v \rangle = W$ a vector space.
 - Let $g(X) = X^k - (a_{k-1}X^{k-1} + \dots + a_1X + a_0)$. Then $g(T)v = 0$. This implies that g is the minimal polynomial of T .
 - It follows that $T^h g(T)v = 0$. Thus, $g(T)T^h v = 0$ for all h .
 - Lastly, it follows that $g(T)w = 0$ for all $w \in W$.
 - Assume v is a cyclic vector. Then $W = V$. It follows that $g(T)v = 0$ for all $v \in V$.
 - The original assumption posits that no polynomial of degree less than or equal to $k - 1$ can annihilate v .
- Consider $V = F[X]/(f)$. Let $\deg(f) = d$, let $T : V \rightarrow V$, and let T be the “multiply by X ” linear transformation. It follows that if $v_i = \bar{X}^{i-1}$ ($i = 1, \dots, d$), then

$$Tv_i = v_{i+1}$$

for $i = 1, \dots, d - 1$ and

$$Tv_d = -(a_0 v_1 + a_1 v_2 + \dots + a_{d-1} v_d)$$

- If $d = 3$, then we have

$$M(T) = \begin{pmatrix} 0 & 0 & -a_0 \\ 1 & 0 & -a_1 \\ 0 & 1 & -a_2 \end{pmatrix}$$

- The above matrix is called the **companion matrix** of f (monic of degree 3).

- **Rational canonical form:** The form (V, T) given by

$$F[X]/(f_1) \oplus \cdots \oplus F[X]/(f_s)$$

where $f_2 \mid f_1, \dots, f_s \mid f_{s-1}$ and $\deg(f_s) > 0$.

- When $V = 0$, then $s = 0$. In this case, f_1 is the minimal polynomial of T .
- The form consisting of a block diagonal matrix of companion matrices.

- **Jordan canonical form:**

- Has to do with p -primary components!

- There's one more canonical form, too.
- Since no one knows what canonical forms are and we very much need them for what Nori was planning to do, Nori will change his plans. No tensors in the last week, either.
- p -primary components: When $p = X - a$, $a \in F$.
- (V, T) is **p -primary** if there exists an n such that $(T - a)^n v = 0$ for all $v \in V$.
- $1_V : V \rightarrow V$ is the identity.
- $a \cdot 1_V = a_V : V \rightarrow V$.
- $(T - a_v)^n = 0 \in \text{End}_F(V)$.
- We're now doing generalized eigenspaces ?? lol.
- The p -primary component is as the generalized a -eigenspace.
 - $(T - a)v = 0$, i.e., $Tv = av$ is the a -eigenspace; the eigenspaces are components of the generalized eigenspaces.
- Let $V = F[X]/(X - a)^n$. Let $v_1 = 1$, $v_2 = \overline{X - a}$, \dots , $v_n = \overline{(X - a)^{n-1}}$.
 - We know that $X(X - a)^r = (X - a + a)(X - a)^r = (X - a)^{r+1} + a(X - a)^r$.
 - Nori writes Jordan blocks as

$$\begin{pmatrix} a & 0 & 0 & 0 \\ 1 & a & 0 & 0 \\ 0 & 1 & a & 0 \\ 0 & 0 & 1 & a \end{pmatrix}$$

not with 1's in the superdiagonal.

- Thus, the *last* generalized eigenvector is an eigenvector here, instead of the *first*.

8.2 Office Hours (Nori)

- Midterm: We never covered the universal property of a quotient in class, did we?
 - That's the special lemma from last office hours.
- PSet 7: 7.3 and 7.4 typos.
- Wednesday lecture?
 - Seventh week summary will suffice.
- What do you need us to know about the rational canonical form? Should I still read Dummit and Foote (2004), Section 12.2 or is that no longer necessary?
 - Nori will probably push ahead with 12.2. Thus I should read it. He's not sure what he'll do beyond that, though, since he doesn't want to jam tensors into the last week.
 - I will need tensor products for representation theory, regardless, so if I want to take it, I should self-study it.
 - No chance tensor products will be covered next quarter.
 - Serre is a terrific mathematician whose wife is a super chemist, and that's why he wrote his book on representation theory (and wrote it in a less terse manner than usual).
 - No tensors means no exterior algebra, too.
 - Nori hasn't read any of Dummit and Foote (2004).
 - The transfer theory of groups arises in a later chapter, and that's important for representation theory, though.
- Nori doesn't think any teacher pays attention to what courses are supposed to cover as stated in the course catalog.
 - We will never do modules, multilinear and quadratic forms.
 - p -adic field and Galois theory.
 - Nori thinks the proof of Theorem 12.4 is very difficult to follow for a first-timer.
 - Solvable groups were supposed to be a MATH 25700 topic, but got cut because of 9-week quarters.
 - Sycotomic fields have applications to the representation theory of finite groups; there are theorems of representation theory that you need sycotomic fields to prove.
 - Emil Artin: Galois Theory is worth looking up.
 - Gauss and constructions of 17-gons also needs sycotomic fields.

8.3 Office Hours (Nori)

- 2/21:
- Lecture 6.1: Proposition proof?
 - Lecture 6.1: $(2) \subsetneq \mathbb{Z}$ example?
 - Lecture 6.1: The end of the theorem proof.
 - Lecture 6.2: Does the first theorem you proved not appear in the book until Chapter 12?
 - Lecture 6.2: What is A in the proof?
 - Resources for the proofs in Week 6?
 - Lecture 7.1: Quotient stuff.

- Lecture 7.2: Why does $\text{Ann}(v) = (p^k)$, why not just $(p^k) \subset \text{Ann}(v)$? Additionally, how does $p^k w' = 0$ imply that $p^k \in \text{Ann}(w)$?
 - R is a PID!
 - $\text{Ann}(w)$ should be $\text{Ann}(w')$ in the centered line.
 - We don't need to know the theorem from the book for a while (second year of graduate school at least).
 - It's good to know the proofs from class just for going forward in math, but we probably will not be asked to reproduce them on an exam.
- Lecture 7.3: RCF proof?
 - It's not $m_i, 1$, it's $m_{i,1}$!
 - Rewrite the proof when I'm awake enough to understand it.

8.4 Chapter 12: Modules over Principal Ideal Domains

From Dummit and Foote (2004).

Section 12.2: The Rational Canonical Form

- As stated previously, we apply the results of Section 12.1 to $F[X]$ -modules herein.
- Let V be a finite dimensional vector space over F of dimension N . Let (V, T) be an $F[X]$ -module.
- Since V is finite dimensional, it is finitely generated as an F -module and hence also as an $F[X]$ -module.
- If V were free, it would be isomorphic to a direct sum of copies of $F[X]$ (by Theorem 12.5(1)) and hence be infinite dimensional.
 - Thus, V is a torsion $F[X]$ -module.
 - Theorem 12.5(3): V is isomorphic to the direct sum of cyclic, torsion $F[X]$ -modules.
 - This decomposition will allow us to choose a basis for V with respect to which the matrix representation for the linear transformation T is in a specific simple form.
- **Rational canonical form** (of a matrix): The form obtained when we use the invariant factor decomposition of the relevant vector space.
- **Jordan canonical form** (of a matrix): The form obtained when we use the elementary divisor decomposition (and when F contains all the eigenvalues of T).
- Theorem 12.9 ensures that the RCF and JCF are unique, justifying the labeling of them as *canonical*.
- An application of canonical forms: Classifying distinct linear transformations.
 - Two matrices that represent the same linear transformation (hence are similar) have the same RCF and JCF.
 - This is another instance of the structure of the space being acted upon (e.g., the invariant factor decomposition of V) providing information on the algebraic objects (e.g., linear transformations) which are acting.
- **Representation Theory of Groups**: The special case of algebraic objects acting on spaces concerning groups acting on vector spaces.
- **Eigenvalues, eigenvectors, eigenspaces**, and the **determinant** are defined for linear transformations and analogously for matrices.

- Properties of eigenvalues.

Proposition 12.12. TFAE.

1. λ is an eigenvalue of T .
2. $\lambda I - T$ is a singular linear transformation of V .
3. $\det(\lambda I - T) = 0$.

Proof. Given. □

- **Characteristic polynomial** (of a linear transformation): The polynomial defined as follows, where T is the linear transformation in question. *Denoted by $c_T(X)$. Given by*

$$c_T(X) = \det(XI - T)$$

- Defined similarly for matrices A .
- A monic polynomial of degree $\dim V$.
- The eigenvalues are the roots.
- **Minimal polynomial** (of a linear transformation): The unique monic polynomial which generates the ideal $\text{Ann}(V)$ in $F[X]$. *Denoted by $m_T(X)$.*
 - Defined similarly for matrices A .
 - We know that such a polynomial exists by Theorem 12.5(3).
 - Exercise 12.2.5: The degree of the minimal polynomial is at most n^2 .
- **Cayley-Hamilton Theorem:** The minimal polynomial for T is a divisor of the characteristic polynomial for T .
 - Thus, the degree of the minimal polynomial is at most n .
- We now build up to the **rational canonical form**.
- Introduction.

- Theorem 12.5: There exists an isomorphism

$$V \cong F[X]/(a_1(X)) \oplus \cdots \oplus F[X]/(a_m(X)) \quad (12.1)$$

- The invariant factors a_i are only determined up to units, but since $F[X]^\times = F - \{0\}$, we can make the a_i unique by requiring them to be monic.
- Theorem 12.5(3) asserts that $(a_m(X)) = \text{Ann}(V)$.
- The minimal polynomial and the invariant factors.

Proposition 12.13. The minimal polynomial $m_T(X)$ is the largest invariant factor of V . All of the invariant factors of V divide $m_T(X)$.

- We now build up to calculating the minimal polynomial of T and the other invariant factors.
- Choosing a basis for each of the summands in Equation 12.1.
 - Recall that the action of T on V is equivalent to the action of X on each summand.
 - Recall also (from the Example following Proposition 11.1) that $1, \bar{X}, \bar{X}^2, \dots, \bar{X}^{k-1}$ gives a basis of $F[X]/(a(X))$, where $a(X) = X^k + b_{k-1}X^{k-1} + \cdots + b_0$.

- With respect to this basis, the linear transformation $T = l_X$ acts via

$$\begin{aligned}
 1 &\mapsto \bar{X} \\
 \bar{X} &\mapsto \bar{X}^2 \\
 \bar{X}^2 &\mapsto \bar{X}^3 \\
 &\vdots \\
 \bar{X}^{k-2} &\mapsto \bar{X}^{k-1} \\
 \bar{X}^{k-1} &\mapsto \bar{X}^k = -b_0 - b_1\bar{X} - \cdots - b_{k-1}\bar{X}^{k-1}
 \end{aligned}$$

- The last equality holds since $a(\bar{X}) = 0$ in $F[X]/(a(X))$.
- With respect to this basis, the matrix for multiplication by X is called the **companion matrix** of $a(X)$.
- Applying this procedure to each of the cyclic modules on the right side of Equation 12.1 under an appropriate basis yields the **direct sum** of the companion matrices for the invariant factors as the matrix of T .
- Note that this matrix is uniquely determined by the invariant factors of the $F[X]$ -module V . These invariant factors, in turn, uniquely determine V up to isomorphism by Theorem 12.9.
- **First subdiagonal**: The set of entries in a matrix which lie directly below a diagonal entry. *Also known as subdiagonal.*
- **Companion matrix** (of a polynomial): The $k \times k$ matrix, pertaining to the polynomial $a(X) = X^k + b_{k-1}X^{k-1} + \cdots + b_0$, which consists of 1's down the first subdiagonal, $-b_0, \dots, -b_{k-1}$ down the last column, and zeros elsewhere. *Denoted by $\mathcal{C}_{a(X)}$. Given by*

$$\mathcal{C}_{a(X)} = \begin{pmatrix} 0 & 0 & \cdots & \cdots & \cdots & -b_0 \\ 1 & 0 & \cdots & \cdots & \cdots & -b_1 \\ 0 & 1 & \cdots & \cdots & \cdots & -b_2 \\ 0 & 0 & \ddots & & & \vdots \\ \vdots & \vdots & & \ddots & & \vdots \\ 0 & 0 & \cdots & \cdots & 1 & -b_{k-1} \end{pmatrix}$$

- **Direct sum** (of matrices): The block diagonal matrix consisting of the component matrices.
 - See the RCF example below.
- **Rational canonical form** (of a matrix): A matrix that is the direct sum of companion matrices for monic polynomials $a_1(X), \dots, a_m(X)$ of degree at least one with $a_1(X) \mid a_2(X) \mid \cdots \mid a_m(X)$. *Also known as RCF. Given by*

$$\begin{pmatrix} \mathcal{C}_{a_1(X)} & & & \\ & \mathcal{C}_{a_2(X)} & & \\ & & \ddots & \\ & & & \mathcal{C}_{a_m(X)} \end{pmatrix}$$

- **Invariant factors** (of the RCF): The polynomials a_i in the above definition.
- Definition of a **block diagonal** matrix.
- **Rational canonical form** (of a linear transformation): The matrix representing T which is in rational canonical form.
- Dummit and Foote (2004) proves that the rational canonical form is unique by means of running the generation process in reverse.

Theorem 12.14 (Rational Canonical Form for Linear Transformations). Let V be a finite dimensional vector space over the field F , and let T be a linear transformation of V .

1. There is a basis for V with respect to which the matrix for T is in rational canonical form, i.e., is a block diagonal matrix whose diagonal blocks are the companion matrices for monic polynomials $a_1(X), \dots, a_m(X)$ of degree at least one with $a_1(X) \mid a_2(X) \mid \dots \mid a_m(X)$.
 2. The rational canonical form for T is unique.
- Why the *rational* canonical form?
 - “Rational” refers to the fact that this canonical form is calculated entirely within the field F and exists for any linear transformation T .
 - This is not the case for the JCF, which only exists if the field F contains the eigenvalues for T .
 - Similar matrices, modules, and the RCF.

Theorem 12.15. Let S and T be linear transformations of V . Then TFAE.

1. S and T are similar linear transformations.
2. The $F[X]$ -modules obtained from V via S and via T are isomorphic $F[X]$ -modules.
3. S and T have the same rational canonical form.

Proof. Given. □

- Observation: Any $n \times n$ matrix A with entries in F arises as the matrix for some linear transformation T of an n -dimensional vector space.
- This observation allows us to restate Theorems 12.14-12.15 in the language of matrices.

Theorem 12.16 (Rational Canonical Form for Matrices). Let A be an $n \times n$ matrix over the field F .

1. The matrix A is similar to a matrix in rational canonical form, i.e., there is an invertible $n \times n$ matrix P over F such that $P^{-1}AP$ is a block diagonal matrix whose diagonal blocks are the companion matrices for monic polynomials $a_1(X), \dots, a_m(X)$ of degree at least one with $a_1(X) \mid a_2(X) \mid \dots \mid a_m(X)$.
2. The rational canonical form for A is unique.

Theorem 12.17. Let A, B be $n \times n$ matrices over the field F . Then A, B are similar iff A, B have the same RCF.

- **Invariant factors** (of a matrix): The invariant factors of the matrix’s RCF.
- RCF and similarity questions for A do not depend on which field contains the entries of A .

Corollary 12.18. Let A, B be two $n \times n$ matrices over a field F , and suppose F is a subfield of the field K .

1. The rational canonical form of A is the same whether it is computed over K or over F . The minimal and characteristic polynomials and the invariant factors of A are the same whether A is considered as a matrix over F or as a matrix over K .
2. The matrices A, B are similar over K iff they are similar over F , i.e., there exists an invertible $n \times n$ matrix P with entries from K such that $B = P^{-1}AP$ iff there exists an (in general different) invertible $n \times n$ matrix Q with entries from F such that $B = Q^{-1}AQ$.

Proof. Given. □

- Takeaways from Corollary 12.18.
 - The RCF for A is an $n \times n$ matrix with entries in the smallest field containing the entries of A .
 - Further explanation of the word *rational*: The RCF is the same matrix even if we allow conjugation of A by nonsingular matrices whose entries come from larger fields.
- Characteristic polynomials and invariant factors.

Lemma 12.19. Let $a(X) \in F[X]$ be any monic polynomial.

1. The characteristic polynomial of the companion matrix of $a(X)$ is $a(X)$.
2. If M is the block diagonal matrix

$$M = \begin{pmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_k \end{pmatrix}$$

given by the direct sum of matrices A_1, \dots, A_k , then the characteristic polynomial of M is the product of the characteristic polynomials of A_1, \dots, A_k .

Proof. See the exercises. □

Proposition 12.20. Let A be an $n \times n$ matrix over the field F .

1. The characteristic polynomial of A is the product of all the invariant factors of A .
2. (The Cayley-Hamilton Theorem) The minimal polynomial of A divides the characteristic polynomial of A .
3. The characteristic polynomial of A divides some power of the minimal polynomial of A . In particular, these polynomials have the same roots, not counting multiplicities.

Proof. Given. □

- The relations in Proposition 12.20 are frequently useful in determining the invariant factors of A , particularly for $\deg(A)$ small.
- **Elementary row and column operations:** The following three operations, where A is an $n \times n$ matrix over the field F and $XI - A$ is an $n \times n$ matrix with entries in $F[X]$. *Given by*
 - (i) Interchanging two rows or columns.
 - (ii) Adding a multiple (in $F[X]$) of one row or column to another.
 - (iii) Multiplying any row or column by a unit in $F[X]$, i.e., by a nonzero element in F .
- **Smith Normal Form** (of a matrix): The following form of the $n \times n$ matrix $XI - A$ with entries from $F[X]$, where a_1, \dots, a_m are the invariant factors of A . *Given by*

$$\begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & a_1(X) & & \\ & & & & a_2(X) & \\ & & & & & \ddots \\ & & & & & & a_m(X) \end{pmatrix}$$

- Computing the invariant factors in general.

Theorem 12.21. Let A be an $n \times n$ matrix over the field F . Using the three elementary row and column operations above, the $n \times n$ matrix $XI - A$ with entries in $F[X]$ can be put into Smith Normal Form.

- Dummit and Foote (2004) provides algorithms for computing the invariant factor decomposition and the RCF. Return to later.