

Making the Case for Skeletal Muscle Myopathy and Its Contribution to Exercise Intolerance in Heart Failure With Preserved Ejection Fraction

See Article by Weiss et al

Dalane W. Kitzman, MD
Mark J. Haykowsky, PhD
Corey R. Tomczak, PhD

Thirty-five years ago, Robert Luchi provided the first description of heart failure (HF) with preserved ejection fraction (HFpEF). HFpEF is now the most common form of HF in older adults, particularly women, and its prevalence is increasing and its prognosis is worsening. The primary symptom in chronic HFpEF, even when patients are well compensated and nonedematous, is severe exercise intolerance, characterized by exertional fatigue and dyspnea, associated with reduced quality of life.¹ Thus, understanding the pathophysiology of exercise intolerance in HFpEF is critical for improving patient-centered outcomes.

Exercise intolerance can be objectively and reproducibly measured as reduced peak exercise oxygen consumption (VO_2) by expired gas analysis. Using this technique, we and others have shown that the reduction in peak VO_2 in HFpEF is at least as severe as in age-matched people with HF with severely reduced ejection fraction (HFrEF; mean ejection fraction, 30%).¹ By the Fick equation, reduced peak VO_2 must be because of either reduced cardiac output (CO), reduced arteriovenous oxygen content difference ($\text{A-VO}_2\text{Diff}$), or a combination of these factors.² It has conventionally been assumed that reduced exercise CO is the sole driver of exercise intolerance in HFpEF. However, we showed that reduced exercise $\text{A-VO}_2\text{Diff}$ accounts for at least 50% of the reduction in peak VO_2 and is a stronger independent predictor of peak VO_2 than exercise CO,³ results confirmed by several others.⁴

Reduced $\text{A-VO}_2\text{Diff}$ during exercise can be caused by reduced convective and diffusive oxygen delivery to exercising skeletal muscle, impaired oxygen utilization by the exercising skeletal muscle, or a combination of these factors.² Indeed, it is now known that, as has been previously established in HFrEF, there are multiple skeletal muscle abnormalities in HFpEF that impair oxygen utilization and seem to contribute to reduced peak VO_2 (Table).^{2,5-11} Among these, growing evidence indicates that impaired mitochondrial function may be among the most consequential. As the sole mechanism for using oxygen and fuel substrate to produce energy, mitochondrial health is obviously a critical determinant of peak VO_2 . Multiple reports support that muscle mitochondrial function is impaired in HFpEF and is a significant contributor to reduced peak $\text{A-VO}_2\text{Diff}$ and consequently peak VO_2 . Our group showed that HFpEF patients have a downward-shifted relationship of peak VO_2 to percent lean leg muscle mass, indicative of impaired oxygen utilization during exercise.⁵ Using skeletal muscle biopsy, we showed that HFpEF patients have 3 separate findings relatively specific for mitochondrial dysfunction and that were significant independent predictors of their reduced peak VO_2 : reduced type 1 (oxidative) muscle fibers;⁶ reduced mitochondrial density;⁷ and reduced citrate synthase, a key enzyme regulating oxidative metabolism.⁷ We also found evidence of impaired mitochondrial fusion that was associated with reduced peak VO_2 .⁷ In an animal model of HFpEF, Bowen et al⁹ found multiple abnormalities, including reduced in situ mitochondrial respiratory reserve capacity, a key measure of

The opinions expressed in this editorial are not necessarily those of the editors or of the American Heart Association.

Correspondence to: Dalane W. Kitzman, MD, Kermit Glenn Phillips II Chair in Cardiovascular Medicine, Professor of Internal Medicine: Section on Cardiovascular Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27012. E-mail dkitzman@wakehealth.edu

Key Words: Editorials
■ exercise intolerance ■ heart failure ■ magnetic resonance spectroscopy ■ mitochondrial function ■ skeletal muscle

© 2017 American Heart Association, Inc.

Table. Skeletal Muscle Abnormalities in HFpEF

Reduced % lean muscle mass
Fiber atrophy
Increased intermuscular adipose
Increased ratio of thigh intermuscular adipose/muscle areas
Reduced capillary density (capillary/fiber ratio)
Downward shifted relationship between %leg lean mass/peak VO ₂ (impaired oxygen utilization)
Reduced % type 1 (oxidative) muscle fibers
Shift in type 1/type 2 fiber ratio
Reduced citrate synthase activity
Reduced mitochondrial density
Impaired mitochondrial fusion
Reduced mitochondrial respiratory reserve capacity (maximal respiratory control ratio)
Accelerated high-energy phosphate depletion during exercise
Delayed high-energy phosphate repletion after exercise
Increased fatigability
Increased oxidative stress

Most of these abnormalities have been correlated with reduced exercise capacity. HFpEF indicates heart failure with preserved ejection fraction.

skeletal muscle oxidative phosphorylation that correlates well with peak VO₂ in humans.

However, the strongest proof that a specific abnormality contributes to exercise intolerance is provided when the factor is measured during exercise. This is enabled by phosphorous magnetic resonance spectroscopy, a technique that allows continuous assessments of ATP and creatine phosphate (PCr) concentrations and turnover rates during and after exercise. The rates of breakdown and resynthesis of high-energy phosphates during and after exercise are fundamental determinants of whole-body VO₂ during exercise and recovery, and PCr kinetics serve as an excellent indirect measure of mitochondrial oxidative capacity. Bhella et al⁸ were the first to use magnetic resonance spectroscopy to report abnormal PCr kinetics during and after exercise in HFpEF; however, only 2 patients were studied.

In this issue of the journal, Weiss et al¹¹ report an elegantly designed study with sophisticated magnetic resonance spectroscopy measurements that markedly extends prior literature. Weiss et al performed serial magnetic resonance spectroscopy measurements of PCr during calf extensor exercise to exhaustion and during recovery in HFpEF patients compared with HFrEF patients and healthy controls. Calf extensor exercise was an important feature because the small mass of exercising muscle excludes reduced CO as a cause of exercise limitation because even a severely weakened heart would have ample capacity for the work involved. Weiss et al found that compared with normal subjects, HFpEF patients had severe exercise intolerance, and this was associated with rapid depletion of high-energy

phosphate, which was observed early during exercise, further excluding reduced CO and muscle blood flow reserve as a cause. Furthermore, HFpEF patients had markedly delayed repletion of high-energy phosphate during recovery. These abnormalities were significantly worse in HFpEF than in HFrEF.¹¹ These data provide the strongest proof to date that HFpEF patients have significantly impaired skeletal muscle bioenergetics that contribute to their severe exercise intolerance.

Weiss et al also found markedly increased intermuscular adipose tissue in HFpEF, a finding we originally reported and showed to be correlated with reduced peak VO₂.¹⁰ Excess intermuscular adipose tissue is inversely related to mitochondrial density and seems to suppress mitochondrial biogenesis.¹⁰ Surprisingly, Weiss found no significant correlation between intermuscular adipose and PCr kinetics.

These findings of abnormal skeletal muscle mitochondrial function in HFpEF and their contribution to exercise intolerance should not be a surprise because HFrEF patients have the same abnormalities that significantly contribute to their severe exercise intolerance, highlighting that in HF patients, in general, skeletal muscle dysfunction is a major contributor to exercise intolerance.^{2,12}

Together with other abnormalities previously reported in skeletal muscle in HFpEF patients,^{2,5-11} these data make a strong case for a skeletal muscle myopathy in HFpEF, similar to that described in HFrEF.^{2,12} Importantly, these abnormalities are not merely secondary to deconditioning since (1) they develop even when physical activity is forcibly maintained during the development of HFrEF¹²; and (2) the pattern of abnormalities differs from deconditioning, particularly the fiber-type shift, which is the exact opposite from deconditioning. Further, multiple lines of evidence strongly support that these abnormalities are also not due merely to reduced CO because (1) they persist even when CO is relatively preserved and when CO is normalized with inotropes or cardiac transplant¹²; and (2) their improvement does not correlate with changes in CO.¹²

Thus, the skeletal muscle abnormalities are likely intrinsic to the HFpEF syndrome and not a secondary consequence or an epiphenomenon. The intrinsic nature of skeletal muscle dysfunction is consistent with the current paradigm of HFpEF as a systemic syndrome, likely triggered by circulating factors, such as inflammation or other as yet undiscovered factors, that then cause dysfunction in multiple organ systems.¹³ If it were not so, then why would the circulating triggering factors only damage myocardial muscle, while sparing skeletal muscle, which shares many fundamental characteristics with cardiac muscle? Indeed, infusion of blood from an old animal into a young animal not only creates HFpEF-like changes in the heart but also in skeletal muscle. This also is not necessarily unique to HFpEF because in many ways, HFrEF is a systemic syndrome as well.

These data have potentially important therapeutic implications. Exercise training is the only intervention definitively proven to improve peak VO_2 in HFpEF.¹⁴ We previously showed that >90% of the improvement in peak VO_2 with exercise training is because of improved A- VO_2 Diff, and a meta-analysis of 6 trials indicates that training improves peak VO_2 in HFpEF without significantly altering resting systolic or diastolic function.¹⁴ Bowen et al⁹ showed that the impaired mitochondrial dysfunction in their HFpEF animal model was prevented by exercise training. Thus, it is possible that improvement in skeletal muscle mitochondrial function is a significant contributor to training-related improvements in peak VO_2 in human HFpEF, which is known to be the case for HFrEF.¹² Furthermore, caloric restriction, which can improve mitochondrial function, improves peak VO_2 in obese HFpEF.¹⁵

How can exercise training (and caloric restriction) improve peak VO_2 when nearly all of the pharmacological agents tested in clinical trials to date, spanning several drug classes, have failed? Pharmacological trials have targeted cardiac and vascular mechanisms. However, myocardium is terminally differentiated, with limited capacity for improvement. Arterial stiffness, the most consistently observed vascular abnormality in HFpEF, develops over decades, is associated with medial calcification, and has not been modifiable in HFpEF, even with prolonged therapy and novel agents. In contrast, skeletal muscle, including mitochondrial biogenesis, has robust capacity for rapid rejuvenation and repair, with detectable improvements within 3 days after initiating exercise training. This makes a strong case for targeting skeletal muscle, including mitochondrial dysfunction, to improve exercise intolerance in HFpEF.

Further supporting the value of targeting skeletal muscle abnormalities in HFpEF is that, compared with myocardium, obtaining objective measurements of skeletal muscle mass and function, including mitochondrial function, is relatively easy and safe. Moreover, there are now multiple agents in phase 2 clinical trials, primarily of older patients with physical disability associated with sarcopenia, targeting a variety of skeletal muscle abnormalities, including mitochondrial dysfunction. There are also several agents shown in animal models to improve muscle mitochondrial dysfunction. Of particular note, a recently launched study (PANACHE [Partial Adenosine A1 Receptor Agonist in Chronic Heart Failure With Preserved Ejection Fraction], NCT No. 03098979) may be the first trial to test whether a pharmacological agent (neladenoson bialanate, BAY1067197) targeted specifically to skeletal muscle and myocardial mitochondrial dysfunction can improve exercise intolerance in HFpEF.

Thus, while continuing the quest for agents that address the cardiac and arterial abnormalities contributing to exercise intolerance in HFpEF, there is a compelling case to target the skeletal muscle abnormalities that contribute at least as strongly to exercise intoler-

ance and that are more plastic and may be more easily remediable. While treating HFpEF with agents aimed at skeletal muscle may seem counterintuitive, HFpEF patients will be grateful for their improved exercise tolerance and quality of life, regardless of from which muscle the improvement derives.

SOURCES OF FUNDING

Dr Kitman is supported by National Institutes of Health (NIH) grants R01AG18915, R01AG045551, R01HL107257, P30AG021332, R15NR016826, The Kermit Glenn Phillips II Chair in Cardiovascular Medicine, Wake Forest School of Medicine, Dr Haykowsky is supported by the Moritz Chair in Geriatrics in the College of Nursing and Health Innovation at the University of Texas at Arlington, and Dr Tomczak the Saskatchewan Health Research Foundation and the Heart and Stroke Foundation of Canada.

DISCLOSURES

Dr Kitman declares the following relationships: consultant for Abbvie, Relypsa, Corvia Medical, and Bayer; research grant funding from Novartis, Bayer, and St. Luke's Hospital of Kansas City; and stock ownership in Gilead Sciences. The other authors report no conflicts.

AFFILIATIONS

From the Department of Cardiovascular Medicine and Section on Geriatrics and Gerontology, Wake Forest School of Medicine, Winston-Salem, NC (D.W.K.); College of Nursing and Health Innovation, University of Texas at Arlington (M.J.H.); and Integrative Cardiovascular Physiology Lab, College of Kinesiology, University of Saskatchewan, Saskatoon, Canada (C.R.T.).

FOOTNOTES

Circ Heart Fail is available at <http://circheartfailure.ahajournals.org>.

REFERENCES

1. Kitman DW, Little WC, Brubaker PH, Anderson RT, Hundley WG, Marburger CT, Brosnihan B, Morgan TM, Stewart KP. Pathophysiological characterization of isolated diastolic heart failure in comparison to systolic heart failure. *JAMA*. 2002;288:2144–2150.
2. Haykowsky MJ, Tomczak CR, Scott JM, Paterson DI, Kitman DW. Determinants of exercise intolerance in patients with heart failure and reduced or preserved ejection fraction. *J Appl Physiol* (1985). 2015;119:739–744. doi: 10.1152/jappphysiol.00049.2015.
3. Haykowsky MJ, Brubaker PH, John JM, Stewart KP, Morgan TM, Kitman DW. Determinants of exercise intolerance in elderly heart failure patients with preserved ejection fraction. *J Am Coll Cardiol*. 2011;58:265–274. doi: 10.1016/j.jacc.2011.02.055.
4. Dhakal BP, Malhotra R, Murphy RM, Pappagianopoulos PP, Bagdish AL, Weiner RB, Houstis NE, Eisman AS, Hough SS, Lewis GD. Mechanisms of exercise intolerance in heart failure with preserved ejection fraction: the role of abnormal peripheral oxygen extraction. *Circ Heart Fail*. 2015;8:286–294. doi: 10.1161/CIRCHEARTFAILURE.114.001825.

5. Haykowsky MJ, Brubaker PH, Morgan TM, Kritchevsky S, Eggebeen J, Kitzman DW. Impaired aerobic capacity and physical functional performance in older heart failure patients with preserved ejection fraction: role of lean body mass. *J Gerontol A Biol Sci Med Sci*. 2013;68:968–975. doi: 10.1093/gerona/glt011.
6. Kitzman DW, Nicklas B, Kraus WE, Lyles MF, Eggebeen J, Morgan TM, Haykowsky M. Skeletal muscle abnormalities and exercise intolerance in older patients with heart failure and preserved ejection fraction. *Am J Physiol Heart Circ Physiol*. 2014;306:H1364–H1370. doi: 10.1152/ajp-heart.00004.2014.
7. Molina AJ, Bharadwaj MS, Van Horn C, Nicklas BJ, Lyles MF, Eggebeen J, Haykowsky MJ, Brubaker PH, Kitzman DW. Skeletal muscle mitochondrial content, oxidative capacity, and Mfn2 expression are reduced in older patients with heart failure and preserved ejection fraction and are related to exercise intolerance. *JACC Heart Fail*. 2016;4:636–645. doi: 10.1016/j.jchf.2016.03.011.
8. Bhella PS, Prasad A, Heinicke K, Hastings JL, Arbab-Zadeh A, Adams-Huet B, Pacini EL, Shibata S, Palmer MD, Newcomer BR, Levine BD. Abnormal haemodynamic response to exercise in heart failure with preserved ejection fraction. *Eur J Heart Fail*. 2011;13:1296–1304. doi: 10.1093/eurjhf/hfr133.
9. Bowen TS, Rolim NP, Fischer T, Baekkerud FH, Medeiros A, Werner S, Brønstad E, Rognmo O, Mangner N, Linke A, Schuler G, Silva GJ, Wisløff U, Adams V, Optimex Study Group. Heart failure with preserved ejection fraction induces molecular, mitochondrial, histological, and functional alterations in rat respiratory and limb skeletal muscle. *Eur J Heart Fail*. 2015;17:263–272. doi: 10.1002/ejhf.239.
10. Haykowsky MJ, Kouba EJ, Brubaker PH, Nicklas BJ, Eggebeen J, Kitzman DW. Skeletal muscle composition and its relation to exercise intolerance in older patients with heart failure and preserved ejection fraction. *Am J Cardiol*. 2014;113:1211–1216. doi: 10.1016/j.amjcard.2013.12.031.
11. Weiss K, Schar M, Panjraht GS, Zhang Y, Sharma K, Bottomley PA, Golozar A, Steinberg A, Gerstenblith G, Russel SD, Weiss RG. Fatigability, exercise intolerance, and abnormal skeletal muscle energetics in heart failure. *Circ Heart Fail*. 2017;10:e004129. doi: 10.1161/CIRCHEARTFAILURE.117.004129.
12. Middlekauff HR. Making the case for skeletal myopathy as the major limitation of exercise capacity in heart failure. *Circ Heart Fail*. 2010;3:537–546. doi: 10.1161/CIRCHEARTFAILURE.109.903773.
13. Shah SJ, Kitzman DW, Borlaug BA, van Heerebeek L, Zile MR, Kass DA, Paulus WJ. Phenotype-specific treatment of heart failure with preserved ejection fraction: a multiorgan roadmap. *Circulation*. 2016;134:73–90. doi: 10.1161/CIRCULATIONAHA.116.021884.
14. Pandey A, Parashar A, Kumbhani DJ, Agarwal S, Garg J, Kitzman D, Levine BD, Drazner M, Berry JD. Exercise training in patients with heart failure and preserved ejection fraction: meta-analysis of randomized control trials. *Circ Heart Fail*. 2015;8:33–40. doi: 10.1161/CIRCHEARTFAILURE.114.001615.
15. Kitzman DW, Brubaker P, Morgan T, Haykowsky M, Hundley G, Kraus WE, Eggebeen J, Nicklas BJ. Effect of caloric restriction or aerobic exercise training on peak oxygen consumption and quality of life in obese older patients with heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA*. 2016;315:36–46. doi: 10.1001/jama.2015.17346.