

Logistic Regression

Quiz, 5 questions

1
point

1.

Suppose that you have trained a logistic regression classifier, and it outputs on a new example x a prediction $h_{\theta}(x) = 0.4$. This means (check all that apply):

- ☐ Our estimate for $P(y = 1|x; \theta)$ is 0.6.
 - ☐ Our estimate for $P(y = 0|x; \theta)$ is 0.6.
 - ☐ Our estimate for $P(y = 0|x; \theta)$ is 0.4.
 - ☐ Our estimate for $P(y = 1|x; \theta)$ is 0.4.
-

1
point


2.

Logistic Regression

Quiz, 5 questions

Suppose you have the following training set, and fit a logistic regression classifier $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$.

x_1	x_2	y
1	0.5	0
1	1.5	0
2	1	1
3	1	0



Which of the following are true? Check all that apply.

- ☐ $J(\theta)$ will be a convex function, so gradient descent should converge to the global minimum.
- ☐ Adding polynomial features (e.g., instead using $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_1 x_2 + \theta_5 x_2^2)$) could increase how well we can fit the training data.
- ☐ The positive and negative examples cannot be separated using a straight line. So, gradient descent will fail to converge.
- ☐ Because the positive and negative examples cannot be separated using a straight line, linear regression will perform as well as logistic regression on this data.

3. Logistic Regression

Quiz, 5 questions

For logistic regression, the gradient is given by

$\frac{\partial}{\partial \theta_j} J(\theta) = \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x_j^{(i)}$. Which of these is a correct gradient descent update for logistic regression with a learning rate of α ? Check all that apply.

- ☐ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x_j^{(i)}$ (simultaneously update for all j).
 - ☐ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m \left(\frac{1}{1+e^{-\theta^T x^{(i)}}} - y^{(i)} \right) x_j^{(i)}$ (simultaneously update for all j).
 - ☐ $\theta := \theta - \alpha \frac{1}{m} \sum_{i=1}^m (\theta^T x - y^{(i)})x^{(i)}$.
 - ☐ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x_j^{(i)}$ (simultaneously update for all j).
-

1
point

4.

Which of the following statements are true? Check all that apply.

- ☐ The sigmoid function $g(z) = \frac{1}{1+e^{-z}}$ is never greater than one (> 1).
 - ☐ Linear regression always works well for classification if you classify by using a threshold on the prediction made by linear regression.
 - ☐ For logistic regression, sometimes gradient descent will converge to a local minimum (and fail to find the global minimum). This is the reason we prefer more advanced optimization algorithms such as fminunc (conjugate gradient/BFGS/L-BFGS/etc).
 - ☐ The cost function $J(\theta)$ for logistic regression trained with $m \geq 1$ examples is always greater than or equal to zero.
-

Logistic Regression


Quiz, 5 questions

1 point


5.

Suppose you train a logistic classifier $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$. Suppose $\theta_0 = 6, \theta_1 = -1, \theta_2 = 0$. Which of the following figures represents the decision boundary found by your classifier?


☐ Figure:



☐ Figure:




☐ Figure:



☐ Figure:

Logistic Regression

Quiz, 5 questions

☐

I, **George Wolberg**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account. Learn more about Coursera's Honor Code

Submit Quiz

