1. The probability of a leap year selected at random contain 53 Sunday is:
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at
random. The probability of drawing a black ball is:
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5
3. The probability that it will rain tomorrow is 0.85. What is the
probability that it will not rain tomorrow
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these
4. What is the probability that a number selected from the numbers
(1, 2, 3,,15) is a multiple of 4?
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3
5. What are the total outcomes when we throw three coins?
(a) 4 (b) 5 (c) 8 (d) 7
6. The probability that a prime number selected at random from the
numbers (1,2,3,35) is :
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these
7. The sum of the probability of an event and non event is :
(a) 2 (b) 1 (c) 0 (d) none of these.
8. The following probabilities are given; choose the correct answer
The following probabilities and given, should all our aniens.
for that which is not possible.
for that which is not possible.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is:
for that which is not possible. (a) 0.15 (b) $2/7$ (c) $7/5$ (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) $1/4$ (b) $3/8$ (c) $1/2$ (d) $1/8$
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has:
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number. (A) 2/3 (B) 1 (C) 5/6 (D) 1/2 12. Two coins are thrown at the same time. Find the probability of
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number. (A) 2/3 (B) 1 (C) 5/6 (D) 1/2 12. Two coins are thrown at the same time. Find the probability of getting both heads.
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number. (A) 2/3 (B) 1 (C) 5/6 (D) 1/2 12. Two coins are thrown at the same time. Find the probability of

13. Two dice are thrown simultaneously. The probability of getting a sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	9	
	ds are numbere ime number.	ed from 1 to 10	00. Find the	probability of	
(A) 3/4		(C) 1/4	(D)	29/100	
_	a blue ball is do n a bag is:			f the probability n the number of	
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50					
mixed thoro	narked with num oughly. One car lity that the num (B) 1/10	d is drawn fro mber on card i	m this box ras s a perfect s	andomly, then equare.	
18. What is (A) 1/7	the probability (B) 53/366	•	•	• •	
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$					
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12 1/2					
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4					

22. Riya and Ka same birthday i	-	•	t both will hav	ve the
(A) 364/365		•	(D) ?	/133225
23. A number 2 2. Then the pro (A) 1/5 (B) 2	bability that x2	< 2 is?	e numbers -2,	-1, 0 , 1,
24. A jar contai a marble is draw red is 2/3, then (A) 10 (B) 6	wn at random fr the number of	om the jar, the white marbles	probability th	
25. A number is Then the proba (A) 7/50 (B) 4	bility that it is a	multiple of 3 a		umbers.
26. Consider a with n dots sho showing 4 dots	wing up is prop is?	ortional to n. T	he probability	
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$	
27. Runs score 93, and 20. The	-		tches are 50,	70, 82,
	b) 25.49		d) 25.69	9
28. Find median	ys 15, 11, 9, 5,	18, 4, 18, 1 3, 17		The median is the middle value when a data is ordered Mode is the number that occurs most often
a) 13, 15	b) 13, 18	C) 18, 15		d) 13, 16
29. A coin is to	-	s. The probabil	ity that tails t	=
3 cases is a) $\frac{1}{2}$	_· b) ¹ / ₃	c) $^{1}/_{4}$		THTT, HTTT, TTHT, TTTH $d) \frac{1}{6}$
30. X is a variat	, J	1	of E(X²) is	. 0
a) 8 b)		c) 27	d) 9	
31.The random respectively. Le				.5
2			25 Va	ar(x) - 4Var(y)

25 * 0.2 + 4 * 0.5

3

a) 3

b) 4

c) 5

d) 7

32.Out of the following values, which one is not possible in probability?

a) P(x) = 1

b) $\sum x P(x) = 3$

c) P(x) = 0.5

d) P(x) = -0.5

33.If E(x) = 2 and E(z) = 4, then E(z - x) = ?

a) 2

b) 6

c) 0

d) Insufficient data

34. The covariance of two independent random variable is

a) 1

b) 0

c) - 1

d) Undefined

35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is?

a) 0

b) 1

c) 3

d) Insufficient data

36.If P(x) = 0.5 and x = 4, then E(x) = ?

a) 1

b) 0.5

c) 4

d) 2

37.In a discrete probability distribution, the sum of all probabilities is always?

a) 0

b) Infinite

c) 1

d) Undefined

38.If the probability of hitting the target is 0.4, find mean and variance.

M = 0.4V = 0.4(1-0.4)

a) 0.4, 0.24

b) 0.6, 0.24

c) 0.4, 0.16

d) 0.6, 0.16

39. If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance?

a) 0.6, 0.24

b) 6, 2.4

c) 0.4, 0.16

d) 4, 1.6

M = 10 * 0.6 V = 10 * 0.6 * (1-0.6)

40. Find the mean of tossing 8 coins.

8 * 1/2 = 4

a) 2

b) 4

c) 8

d) 1

41. What is the mean and variance for standard normal distribution?

a) Mean is 0 a	ınd variance is	b) Mean is 1 and	variance	is 0
c) Mean is 0 a	ınd variance is	∞ d) Mean is ∞ an	d variand	ce is 0
40.14				
		ariable X is given by $_{ extsf{-}}$		•
a) E(X)	b) E(X2)	c) $E(X2) - (E(X))^2$	2	d) (E(X))2
43.Mean of a	random varia	ble X is given by		
a) E(X)	b) E(X2)	c) $E(X2) - (E(X))2$		d) (E(X))2
44.Mean of a	constant 'a' is	·		
a) 0	b) a	c) a/2	d) 1	
45 Variance	f a constant '	o' ic		

46. Find the mean and variance of X?

b) a

f(x) 1/9 2/9 3/9 2/9 1/9 [(2)^2 *3/9] + [(3)^2 *2/9] + [(4)^2 *1/9	Х	0	1	2	3	4	Mean = 0*1/9 + 1*2/9 + 2*3/9 + 3*2/9 + 4*1/9 = 2
	f(x)	1/9	2/9	3/9	2/9	1/9	variance = [(0)^2 *1/9] + [(1)^2 *2/9] + [(2)^2 *3/9] + [(3)^2 *2/9] + [(4)^2 *1/9] = 16/3

c) a/2

a) 2, 4/3

a) 0


b) 3, 4/3

c) 2, 2/3

d) 3, 2/3

d) 1

47. Find the expectation of a random variable X?



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2
- d) \sqrt{npq}