
A Program comparisons tested per cell and per library
-log10 P, libraries as replicates
-log10 P, cells as replicates
Interferon-stimulated
Lipid-associated
Matrix remodeling
Inflammatory monocyte
Angiogenic
Complement
Resident/tissue
B Macrophage pseudobulk from raw counts, ~ sex + site + group
(19 libraries; 0 genes at FDR < 0.05)
-log10 P
log2 fold change (non-healing / healing)
CXCL5
FN1
CXCL14
CTHRC1
MIR3945HG
C Macrophage-state composition per library
(mean and 95% CI)
% of the library's macrophages
Resident
adj. OR 3.26
(1.07-9.89)
Inflammatory
adj. OR 0.40
(0.10-1.60)
Lipid-associated
adj. OR 0.50
(0.20-1.25)
Healing
Non-healing
D Resident-state proportion under alternative models
(model-based P / stratified permutation P)
M0 unadjusted (19)
M1 + sex (19)
M2 + sex + site, primary (19)
M3 + depth, %mito (19)
S1 foot only (14)
S2 putative donors (12)
S3 excl. G2, G13 (17)
S4 excl. opposite-sex cells (19)
P = 0.001 / 0.033
P = 0.039 / 0.037
P = 0.038 / 0.072
P = 0.022 / 0.038
P = 0.382 / 0.208
P = 0.398 / 0.345
P = 0.281 / 0.127
P = 0.054 / 0.049
odds ratio, resident state, non-healing vs healing (95% CI)