
A GSE134431 ulcers (7 vs 6), DESeq2: 50 genes at FDR < 0.05
-log10 P
log2 FC (non-healing / healing)
GK
PTPRN
COL22A1
KRT33B
PLA2G7
FCGR3B
B Gene sets over-represented among genes nominally higher in healing ulcers
Neutrophil extracellular trap formation
Systemic lupus erythematosus
neutrophil activation involved in...
neutrophil degranulation
neutrophil mediated immunity
inflammatory response
regulation of immune response
Leishmaniasis
Phagosome
Staphylococcus aureus infection
-log10 FDR
C Bulk vs single-cell (raw-count pseudobulk)
rho 0.07 (0.05-0.08); nominal in both:
rho 0.35 (0.11-0.56), n = 60
macrophage pseudobulk log2 FC
bulk log2 FC
D Selected immune, complement and lipid genes in bulk ulcers
FABP5
S100A9
CD9
GPNMB
S100A8
LPL
LYVE1
IL1B
PLIN2
FOLR2
CD163
VCAN
C1QC
APOE
TREM2
C1QA
MRC1
C1QB
CXCL8
VSIG4
LYZ
FCN1
AIF1
APOC1
log2 FC (95% CI)
myeloid activation
resident / complement
lipid metabolism