



My Courses (<https://learningcenter.hfsa.org/Users/Home.aspx>)
/ **2026 AHFTC Board Certification Review - OnDemand**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=4f%2fOexzNsorKAF1lCFotvA%3d%3d>)
/ **2026 AHFTC Board Certification Question Bank**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=YSpCYks4EUa7JCy700Rkkw%3d%3d>)
/ **Practice Exam - 1**

Correct Answer

Next Q

Practice Exam - 1

Question 34 of 117



A 43-year-old female with a history of peripartum cardiomyopathy progresses to end-stage heart failure and undergoes heart transplantation. She is concerned about side effects of immunosuppression.

Which of the following statements is correct?

- A. As compared to tacrolimus, cyclosporine use is associated with increased rates of hirsutism, hypertension and diabetes
- ☒ B. The major toxicities of calcineurin inhibitors include kidney dysfunction, hypertension, dyslipidemia, and neurotoxicity
- C. Mycophenolate mofetil inhibits T but not B lymphocyte proliferation
- D. Routine monitoring of mycophenolate levels is recommended in post transplant immunosuppression management

Commentary

References

Testing Point: The Board-certified AHFTC specialist should understand the basic principles of immunosuppression after heart transplant.

Correct Answer: B. The major toxicities of calcineurin inhibitors include kidney dysfunction, hypertension, dyslipidemia, and neurotoxicity

Rationale: The correct answer is B which correctly identifies toxicities common to calcineurin inhibitor therapies.

Answer A is incorrect as cyclosporine is associated with decreased rather than increased rates of diabetes compared to tacrolimus.

Answer C is incorrect as MMF inhibits both T and B lymphocyte proliferation.

Answer D is incorrect as routine monitoring of mycophenolate levels is not recommended post transplantation.

Powered by Oasis.



(<https://www.oasis-lms.com/>).