



(<https://learningcenter.hfsa.org/>).



**My Courses** (<https://learningcenter.hfsa.org/Users/Home.aspx>)  
/ **2026 AHFTC Board Certification Review - OnDemand**  
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=4f%2fOexzNsorKAF1lCFotvA%3d%3d>)  
/ **2026 AHFTC Board Certification Question Bank**  
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=YSpCYks4EUa7JCy700Rkkw%3d%3d>)  
/ **Practice Exam - 1**

Correct Answer

Next Q

## Practice Exam - 1

### Question 89 of 117



A 54-year-old man with chronic HFrEF and a non-ischemic cardiomyopathy (EF 20%) presents to the emergency department with acute decompensated heart failure and cardiogenic shock. He is stabilized with milrinone and his end-organ function returns to normal. A repeat echocardiogram confirms severe LV dysfunction with moderate mitral insufficiency. His RV is enlarged and hypocontractile with a TAPSE of 1.16 cm. Right heart catheterization is performed demonstrating an RA=18, PA= 58/26, and PCWP=24 with a cardiac index=1.8 l/min/m<sup>2</sup> and an SVO<sub>2</sub>=52. The decision is made to list the patient for transplant but he becomes hemodynamically unstable requiring more inotropic support and begins having sustained VT successfully treated with ICD shocks. He is 6'4" and weighs 260# and is a blood type O.

Which of the following is the best single durable mechanical circulatory support intervention for this patient?

- A.** Intra-aortic balloon pump
- B.** Impella CP
- C.** Implantable LVAD
- D.** Total artificial heart

Powered by Oasis.



(<https://www.oasis-lms.com/>).