



(<https://learningcenter.hfsa.org/>).



My Courses (<https://learningcenter.hfsa.org/Users/Home.aspx>)
/ **2026 AHFTC Board Certification Review - OnDemand**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=4f%2fOexzNsorKAF1lCFotvA%3d%3d>)
/ **2026 AHFTC Board Certification Question Bank**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=YSpCYks4EUa7JCy700Rkkw%3d%3d>)
/ **Practice Exam - 1**

Correct Answer

Next C

Practice Exam - 1

Question 23 of 117



Which of the following is not a criterion for LVAD therapy?

- ☒ **A.** An EF of 30% with NYHA class II symptoms on stable GDMT
- ☐ **B.** NYHA Class IV symptoms
- ☐ **C.** Current treatment with inotropic support
- ☐ **D.** Cardiac index of < 2.2 L/min/m² on GDMT for at least 45 of the past 60 days

Commentary

Correct Answer: A. An EF of 30% with NYHA class II symptoms on stable GDMT

Rationale: Features supporting consideration for durable Left Ventricular Assist Device therapy include advanced Stage D heart failure with persistent severe symptoms despite guideline-directed medical therapy, inotrope dependence, cardiac index < 2.2 L/min/m² despite GDMT for at least 45 of the previous 60 days, or dependence on temporary mechanical circulatory support such as an intra-aortic balloon pump or similar device.

Of the listed choices, answer A is not an accurate LVAD criterion. Reduced ejection fraction alone, particularly in a patient with only NYHA class II symptoms on stable GDMT, is not sufficient to justify LVAD therapy.

