



(<https://learningcenter.hfsa.org/>).



My Courses (<https://learningcenter.hfsa.org/Users/Home.aspx>)
/ **2026 AHFTC Board Certification Review - OnDemand**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=4f%2fOexzNsorKAF1lCFotvA%3d%3d>)
/ **2026 AHFTC Board Certification Question Bank**
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=YSpCYks4EUa7JCy700Rkkw%3d%3d>)
/ **Practice Exam - 1**

Correct Answer

Next Q

Practice Exam - 1

Question 84 of 117



Please select the correct condition.

Which condition is associated with a blunted Y descent in the CVP tracing?

- ☒ **A.** Tamponade
- ☐ **B.** Constriction
- ☐ **C.** Restriction
- ☐ **D.** Heart Transplantation
- ☐ **E.** RV infarction

Commentary

References

A rapid Y descent is a unique characteristic of the CVP waveform that indicates rapid early filling of the RV due to poor RV compliance and generally reflective of RV diastolic dysfunction. All of the choices are characterized by RV diastolic dysfunction/poor RV compliance except tamponade, which is commonly associated with loss of the Y descent due to early (as well as late) restriction of diastolic filling from the pan-diastolic elevation in pericardial pressure from pericardial fluid. Loss of the Y descent would also be expected in tricuspid stenosis. The presence of a rapid Y descent is almost always pathologic and its presence should be investigated unless consistent with a previously known diagnosis.

