



(<https://learningcenter.hfsa.org/>).



**My Courses** (<https://learningcenter.hfsa.org/Users/Home.aspx>)  
/ **2026 AHFTC Board Certification Review - OnDemand**  
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=4f%2fOexzNsorKAF1lCFotvA%3d%3d>)  
/ **2026 AHFTC Board Certification Question Bank**  
(<https://learningcenter.hfsa.org/Users/UserOnlineCourse.aspx?LearningActivityID=YSpCYks4EUa7JCy700Rkkw%3d%3d>)  
/ **Practice Exam - 1**

Correct  
Answer

Show

Next Q

## Practice Exam - 1

### Question 91 of 117



The patient is a 35 year old man status post LVAD implantation one year ago who presents to the emergency room with hematuria and epistaxis. He has currently been on warfarin 5mg daily with a goal INR of 2-2.5. His INR in the ED is 10. A urine toxicology drug screen shows the following:

- Amphet/Methamphetamine: Positive
- Barbiturates, Urine: Negative
- Benzodiazepines, Urine: Negative
- Cannabinoids, Urine: Positive
- Cocaine, Urine: Positive
- Cotinine: >20 ng/ml
- Opiates, urine: Positive
- Phencyclidine, urine: Negative

Which of the following could be responsible for his elevated INR?

**A.** Methamphetamine

**B.** Cannabis

**C.** Cocaine

**D.** Opiates

## References

1. Page RL2, Allen LA, Kloner RA, et al. Medical Marijuana, Recreational Cannabis, and Cardiovascular Health: A Scientific Statement from the American Heart Association. Circulation 2020; 142(10): e131-e152.

2. Hollbrook AM, Pereira JA, Labiris R, et al. Systematic overview of warfarin and its drug and food interactions. Arch Intern Med. 2005;165:1095-1106.

**Powered** by Oasis.



(<https://www.oasis-lms.com/>).