

A VERIFICATION OF SIERPIŃSKI'S CONJECTURE TO 10^{11} AND COMPLETE EXCEPTION SETS FOR $m/n = 1/x + 1/y + 1/z$, $4 \leq m \leq 19$

JAIDEV SHAH

ABSTRACT. Call $n \geq 2$ an *exception* for m if m/n is not a sum of three unit fractions. Erdős and Straus conjectured that $m = 4$ has no exceptions, Sierpiński conjectured the same for $m = 5$, and Schinzel conjectured that every $m \geq 4$ has only finitely many. Pomerance and Weingartner recently proved that exceptions must exist for every sufficiently large m , with numerical evidence that every $m \geq 20$ has a prime exception in $(m^2, 2m^2)$. We contribute the complementary small- m data. First, Sierpiński's conjecture holds for all $2 \leq n \leq 10^{11}$, extending the classical bound 1,057,438,801 by a factor of 94.6. Second, we determine the complete exception sets for every $6 \leq m \leq 19$ over all $n \leq 10^9$, primes and composites alike: 243 exceptions in total, the largest being 83,449 (for $m = 16$), giving empirical values of Schinzel's threshold n_m . Third, we replicate the Pomerance–Weingartner computation for $20 \leq m \leq 60$ and determine the exact boundary of their window phenomenon: among $4 \leq m \leq 60$, the interval $(m^2, 2m^2)$ contains no exception precisely for $m \in \{4, 5, 6, 7, 8, 9, 11, 19\}$. All exceptional values were certified non-representable by two independently written programs, and representability certificates were re-verified in exact rational arithmetic. Code and data accompany the paper.

1. INTRODUCTION

For an integer $m \geq 4$, consider the Diophantine equation

$$(1) \quad \frac{m}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}, \quad x, y, z \in \mathbb{N}.$$

We say $n \geq 2$ is an *exception* for m if (1) has no solution. Erdős and Straus [2] conjectured in 1948 that $m = 4$ has no exceptions; the conjecture has been verified for $n \leq 10^{17}$ by Salez [8], with a recent preprint reporting 10^{18} [5]. Sierpiński [9] conjectured the same for $m = 5$, and Schinzel (see [9]) conjectured that for every $m \geq 4$ there is a threshold n_m such that no $n \geq n_m$ is an exception. Vaughan [10] proved that for fixed m the number of exceptions up to N is $O(N \exp(-c(\log N)^{2/3}))$, and Elsholtz and Tao [1] obtained detailed statistics for the number of solutions of (1) when $m = 4$ and n is prime.

The landscape changed recently: Pomerance and Weingartner [7] proved that if n_m exists then $n_m \geq \exp(m^{1/3+o(1)})$ — exceptions *must* occur for every sufficiently large m — and, in explicit form, that for $m \geq 6.52 \times 10^9$ some prime $p \in (m^2, 2m^2)$ is an exception for m . Their numerical computations support the much stronger assertion that every $m \geq 20$ has a prime exception in this window.

The map of problem (1) thus has three regions: a conjecturally exception-free zone ($m = 4, 5$, and, as our data suggests, 6), a zone of nonempty but apparently finite exception sets ($7 \leq m \leq 19$), and a zone of guaranteed exceptions ($m \geq 20$, provably for large m). This note charts the middle region and pushes the verification frontier of the first. Our contributions:

2020 *Mathematics Subject Classification*. 11D68 (primary), 11Y50, 11D72 (secondary).

Key words and phrases. Egyptian fractions, unit fractions, Erdős–Straus conjecture, Sierpiński conjecture, Schinzel conjecture.

- (1) Sierpiński's conjecture holds for all $2 \leq n \leq 10^{11}$ (Theorem 4.1). The best bounds we could locate are the classical $n \leq 1,057,438,801$ (mid-1960s; cited in Graham's survey [4]) and, for the residual class $n \equiv 1 \pmod{5}$, roughly 10^{10} in the recent preprint [3].
- (2) The complete exception sets for $6 \leq m \leq 19$ over all $n \leq 10^9$ (Theorem 4.2), yielding empirical values of n_m that we could not locate in the literature or in the OEIS.
- (3) An independent replication of the Pomerance–Weingartner window computation for $20 \leq m \leq 60$, together with the exact boundary of the phenomenon: among $4 \leq m \leq 60$ the window $(m^2, 2m^2)$ is exception-free precisely for $m \in \{4, 5, 6, 7, 8, 9, 11, 19\}$ (Section 5).

Remark 1.1. We were unable to consult the numerical section of [7] beyond its abstract while preparing this note. Should it or its references already contain exception sets for $m \leq 19$, priority for the overlap of course belongs there; the 10^9 -exhaustiveness (including composite n), the n_m table, and the Sierpiński bound appear in any case to be new.

2. THE DECISION CRITERION

All computations rest on the following elementary and well-known reduction, which we state with proof to make the note self-contained. The essential point is that it is a *decision procedure*: it can certify non-representability, not merely fail to find solutions.

Lemma 2.1. *Let $m \geq 4$ and $n \geq 2$ with $m \nmid n$. Then (1) is solvable if and only if there exist an integer x with $n/m < x \leq 3n/m$ and a divisor pair $d \cdot d' = b^2$, where $a = mx - n$ and $b = nx$, such that*

$$d \equiv d' \equiv -b \pmod{a}.$$

In that case $y = (b + d)/a$ and $z = (b + d')/a$ solve (1) together with x . If $m \mid n$, then (1) is solvable, e.g. $m/n = 1/(2q) + 1/(3q) + 1/(6q)$ with $q = n/m$.

Proof. Suppose (1) holds with $x \leq y \leq z$. Then $1/x < m/n \leq 3/x$, so $n/m < x \leq 3n/m$, and subtracting $1/x$ gives $1/y + 1/z = a/b$ with $a = mx - n \geq 1$, $b = nx$. From $ayz = b(y + z)$ we get $(ay - b)(az - b) = b^2$; put $d = ay - b$ and $d' = az - b$, both positive and congruent to $-b$ modulo a , with $dd' = b^2$.

Conversely, given such a pair, $y = (b + d)/a$ and $z = (b + d')/a$ are positive integers, and

$$\frac{1}{y} + \frac{1}{z} = \frac{a(2b + d + d')}{(b + d)(b + d')} = \frac{a(2b + d + d')}{b(2b + d + d')} = \frac{a}{b},$$

using $(b + d)(b + d') = b^2 + b(d + d') + dd' = b(2b + d + d')$. Adding $1/x$ recovers (1). $\square$

Enumerating all x in the stated range and, for each x , all divisors $d \leq b$ of b^2 (obtained from the full factorization of $b = nx$) therefore decides (1) for any given pair (m, n) .

Two standard reductions structure the search over n . First, if (1) is solvable for n then it is solvable for every multiple of n (scale the denominators); hence a composite exception must have *all* of its divisors > 1 exceptional, and in particular all of its prime factors. Second, $n < m/3$ is trivially an exception since then $m/n > 3$; we therefore only report exceptions with $n \geq m/3$, in line with the convention of [7].

3. COMPUTATIONS

3.1. Production engine. A C program (`mfrac.c` in the accompanying repository) combines a segmented sieve of Eratosthenes with a first-hit certificate search for each prime p : candidate least denominators $x = \lfloor p/m \rfloor + 1, \lfloor p/m \rfloor + 2, \dots$ are tried in order, and for each x the divisor

condition of Lemma 2.1 is tested over divisors $d = p^j f$ ($j \in \{0, 1, 2\}$, $f \mid x^2$) built from a partial factorization of x , with all congruence arithmetic exact in 64/128-bit integers. A prime not certified within 96 candidate values of x escalates to 20,000 candidates; anything still uncertified is flagged as a suspect and *never* silently classified. Suspects are then settled by the full decision procedure of Section 2. Across all runs reported here (4.6 billion primes) every suspect was resolved; for $m = 12$, for instance, the eleven suspects raised by the scan are precisely the eleven exceptions 433, $\dots$, 12241 of Theorem 4.2.

For every m the range $n \leq 10^6$ (primes *and* composites) was settled by the complete decision procedure directly. The composite gap above 10^6 was closed by the divisor reduction: all 4,769,723 products of exceptional primes lying in $(10^6, 10^9]$ across $7 \leq m \leq 19$ were decided individually; none is an exception. For $m = 5$ the composite case is immediate: every composite $n \leq 10^{11}$ has a prime factor below $\sqrt{10^{11}} < 10^6$, and no $n \leq 10^6$ is an exception for $m = 5$.

3.2. Independent verification. Because a claimed exception is a negative statement, we re-verified every one with a second program (`check2.c`) written independently around a different algorithm: for each admissible x it iterates y over the full interval $(b/a, 2b/a]$ and tests integrality of $z = by/(ay - b)$, with no divisor enumeration, no factorization, and no modular shortcuts. All 243 claimed exceptions were re-certified, with representable neighbours as positive controls. In addition, a brute-force checker over exact rationals (Python `fractions`) was run against both C programs for all $m \leq 14$, $n < 400$ (perfect agreement), and 200 randomly sampled prime certificates with $m \in [5, 19]$, $p \leq 10^9$ were re-verified in exact rational arithmetic (200/200 valid). As a further consistency check, the number of primes processed in the Sierpiński run, 4,117,976,315, equals $\pi(10^{11}) - \pi(10^6)$ exactly.

The total computational cost of all results in this note is under one CPU-hour on four cores of a commodity cloud container.

4. RESULTS

Theorem 4.1. *For every integer $2 \leq n \leq 10^{11}$ there exist positive integers x, y, z with $5/n = 1/x + 1/y + 1/z$.*

Theorem 4.2. *For $6 \leq m \leq 19$, the exceptions $n \leq 10^9$ (with $n \geq m/3$) of equation (1) are exactly the following; there are none for $m = 6$.*

m	exceptions $n \leq 10^9$
6	none
7	2
8	2, 3, 11, 17, 131, 241
9	2, 5, 11, 19
10	2, 3, 7, 11, 43, 61, 67, 181
11	2, 3, 4, 37
12	2, 3, 5, 7, 13, 25, 29, 31, 37, 73, 97, 193, 433, 577, 1129, 1657, 1873, 2521, 2593, 3433, 10369, 12049, 12241
13	2, 3, 4, 5, 7, 14, 53, 61, 67, 79, 211, 281
14	2, 3, 4, 5, 17, 19, 29, 59, 257, 353, 841
15	2, 3, 4, 8, 16, 17, 19, 23, 31, 34, 47, 53, 61, 79, 113, 122, 137, 151, 197, 226, 233, 271, 541, 1103, 1171, 1367, 4201, 6301, 12601, 16831, 20521
16	2, 3, 4, 5, 6, 7, 9, 11, 13, 17, 22, 23, 33, 34, 37, 73, 97, 113, 121, 131, 167, 193, 241, 257, 262, 421, 482, 577, 593, 641, 769, 1201, 1489, 2113, 2521, 2689, 3169, 3361, 4801, 4993, 5281, 8161, 8641, 33601, 36529, 78721, 83449
17	2, 3, 4, 5, 6, 7, 9, 13, 19, 23, 41, 43, 53, 71, 73, 157, 281, 421, 1123, 2081
18	2, 3, 4, 5, 7, 10, 11, 13, 19, 22, 23, 29, 31, 37, 38, 41, 47, 59, 61, 73, 109, 113, 131, 137, 149, 181, 193, 223, 239, 281, 379, 389, 397, 433, 457, 541, 599, 613, 661, 761, 811, 821, 911, 1009, 1297, 1381, 2269, 2819, 9461, 16561, 17389, 28081, 35281
19	2, 3, 4, 5, 6, 7, 8, 10, 11, 13, 23, 26, 29, 41, 43, 46, 82, 97, 137, 181, 193, 229, 353

Under Schinzel’s conjecture that each exception set is finite, and on the evidence that each set above terminates more than four orders of magnitude below the search bound, the resulting empirical thresholds are

m	6	7	8	9	10	11	12	13	14	15	16	17	18	19
n_m	2	3	242	20	182	38	12242	282	842	20522	83450	2082	35282	354

These values are exact if and only if no exception exceeds 10^9 ; Pomerance–Weingartner’s lower bound $n_m \geq \exp(m^{1/3+o(1)})$ concerns large m and does not by itself preclude further exceptions for $m \leq 19$, so the values above should be read as conjectural.

Observation 4.3. (i) Composite exceptions occur only for $m \geq 11$, and in exactly three cases a prime and its square are both exceptional for the same m : (5, 25) for $m = 12$, (29, 841) for $m = 14$, and (11, 121) for $m = 16$.

(ii) The largest exceptions concentrate in the residue class $n \equiv 1 \pmod{m}$. All eleven exceptions above 400 for $m = 12$ are $\equiv 1 \pmod{24}$ — the classes in which the Erdős–Straus equation resists all polynomial identities — and the four largest exceptions for $m = 18$, as well as the three largest for $m = 15$, are all $\equiv 1 \pmod{m}$. The largest exception overall, 83449 for $m = 16$, satisfies $83449 \equiv 9 \pmod{16}$, again a square class. This is consistent with the quadratic-residue obstruction of Schinzel in the $m = 4$ theory (see [6, 1]).

5. THE WINDOW BOUNDARY

Running the complete decision procedure over all primes $p \in (m^2, 2m^2)$ for $20 \leq m \leq 60$ reproduces the numerical claim of [7]: every such m has at least one prime exception in its window, 1494 in total. (The counts grow roughly linearly, from 6 at $m = 20$ to 120 at $m = 60$.) Since Theorem 4.2 determines the small- m exception sets completely, the boundary of the phenomenon can be stated exactly.

Theorem 5.1. *Among $4 \leq m \leq 60$, the interval $(m^2, 2m^2)$ contains no exception for m precisely when*

$$m \in \{4, 5, 6, 7, 8, 9, 11, 19\}.$$

In particular the window phenomenon described in [7] for $m \geq 20$ in fact begins at $m = 10$, with exactly two later gaps at $m = 11$ and $m = 19$. The data suggests:

Conjecture 5.2. *For every $m \geq 4$ outside $\{4, 5, 6, 7, 8, 9, 11, 19\}$, the interval $(m^2, 2m^2)$ contains a prime exception for m .*

For $m \geq 20$ this is supported by [7] (and proved there for $m \geq 6.52 \times 10^9$); the content added by our data is that the eight listed values are the *only* failures.

6. DATA AND CODE AVAILABILITY

The two C programs (`mfrac.c`, `check2.c`), the complete exception lists, the window-exception lists for $20 \leq m \leq 60$, and a run-by-run verification summary are available at

<https://github.com/shahjaidev/explainers/tree/main/unit-fractions>

Every claimed representation can be re-verified from the published witnesses in exact arithmetic, and every claimed exception is reproducible with either program in seconds.

ACKNOWLEDGMENTS

The computations, code, and a draft of this note were produced with the assistance of an AI system (Claude, Anthropic), working under the author's direction; the author reviewed and takes responsibility for all claims. We thank the authors of [7], whose theorem motivated this computation.

REFERENCES

- [1] C. Elsholtz and T. Tao, *Counting the number of solutions to the Erdős–Straus equation on unit fractions*, J. Aust. Math. Soc. **94** (2013), 50–105.
- [2] P. Erdős, *Az $1/x_1 + 1/x_2 + \dots + 1/x_n = a/b$ egyenlet egész számú megoldásairól*, Mat. Lapok **1** (1950), 192–210.
- [3] B. Ghermoul, *Almost a complete proof of the generalized Erdős–Straus conjecture: $5/a = 1/b + 1/c + 1/d$* , arXiv:2508.07367 (2025).
- [4] R. L. Graham, *Paul Erdős and Egyptian fractions*, in: Erdős Centennial, Bolyai Soc. Math. Stud. **25**, Springer, 2013, 289–309.
- [5] S. Mihnea and D. C. Bogdan, *Further verification and empirical evidence for the Erdős–Straus conjecture*, arXiv:2509.00128 (2025).
- [6] L. J. Mordell, *Diophantine Equations*, Academic Press, London, 1969.
- [7] C. Pomerance and A. Weingartner, *Exceptions to the Erdős–Straus–Schinzel conjecture*, Ramanujan J. **69** (2026); arXiv:2511.16817.
- [8] S. E. Salez, *The Erdős–Straus conjecture: new modular equations and checking up to $N = 10^{17}$* , arXiv:1406.6307 (2014).
- [9] W. Sierpiński, *Sur les décompositions de nombres rationnels en fractions primaires*, Mathesis **65** (1956), 16–32.
- [10] R. C. Vaughan, *On a problem of Erdős, Straus and Schinzel*, Mathematika **17** (1970), 193–198.

Email address: shahjaidevn@gmail.com