



Advanced Classical Mechanics

Lecture 6: Conservation Laws

Readings: Morin - Chapter 5, Marion Chapter 2

Amir Shahmoradi

Joshua Alexander Osborne

Department of Physics / College of Science

The University of Texas at Arlington

Arlington, Texas

Newtonian Dynamics

Problem 1. (Force components in polar coordinates):

Recall the definition of Polar Coordinates from Question 6 of our lecture 1 (ACM1). The Cartesian Coordinates can be written in terms of polar coordinates as,

$$x = r \cos \theta ; y = r \sin \theta .$$

We would like to express the equation of motion in Polar Coordinates as opposed to Cartesian Coordinates ($\mathbf{F} = m\mathbf{a} \equiv m\ddot{\mathbf{r}}$).



Using the representation of the Polar bases as illustrated in the figure, show that the Polar equation of motion can be written as,

$$\mathbf{F} \equiv F_r \hat{\mathbf{r}} + F_\theta \hat{\boldsymbol{\theta}} = m\ddot{\mathbf{r}},$$


where,

$$F_r = m(\ddot{r} - r\dot{\theta}^2),$$

$$F_\theta = m(r\ddot{\theta} + 2\dot{r}\dot{\theta}).$$




Problem 2. (Circular pendulum): A mass hangs from a massless string of length l . Conditions have been set up so that the mass swings around in a horizontal circle, with the string making a constant angle β with the vertical. What is the angular frequency, ω , of this motion?



Newtonian Dynamics

Problem 3. (Driving on tilted ground):

A plane tilts down at an angle θ below the horizontal. On this plane, a projectile is fired with speed v at an angle θ above the horizontal, as shown in the figure. What is the distance, d , along the plane that the projectile travels? What is d in the limit $\theta \rightarrow 90^\circ$? What θ yields the maximum horizontal distance?



Problem 4. (A force $F_\theta = 2m\dot{r}\dot{\theta}$): Consider a particle that feels an angular force only, of the form $F_\theta = 2m\dot{r}\dot{\theta}$. Show that $r = Ae^\theta + B^{-\theta}$, where A and B are constants of integration, determined by the initial conditions.

Conservation laws of Physics

- What we mean by the “conservation of a physical quantity” is the fact that its value must remain constant over time.
- Three main conservation laws of physics.
 - Conservation of energy (and mass).
 - Conservation of linear momentum.
 - Conservation of angular momentum.
- These conservation laws are not axioms (i.e., accepted without proof), rather can be derived from the Newtonian Dynamics.
- Historically, there was confusion about conservation of energy vs. momentum.
 - During 1676-1689, Gottfried Leibniz realized (using the works of the contemporary physicist Christiaan Huygens) that the quantity mv^2 appears to be constant when there is no interaction (e.g., no friction). He called this conserved quantity ***vis viva*** (meaning the ***living force***).
 - The contemporary physicist Isaac Newton contested this idea and had suggested that it is another quantity that is conserved and is of the form mv .
 - It took about half a century for other researchers to realize that both Newton and Leibniz were right and that their discoveries referred to two independent conservation laws of Physics.
- We define a force on an object as **conservative force** if and only if it depends on the location of the object (as opposed to velocity or other physical quantities).
- We define **work** for a given force as the energy transferred **to or from** an object as the object is displaced along the direction of the force,

$$w = \mathbf{F} \cdot d\mathbf{x}$$



Gottfried Leibniz



Isaac Newton

Conservation laws of Physics

Problem 5. (Derivation of Conservation of Energy from Newtonian Dynamics):

Consider a force, in just one dimension, that **depends only on position**: $F = F(x)$. If the force acts on a particle of mass m with initial location x_0 and velocity v_0 , show that the integral of the force $F(x)$ over distance traveled leads to the law of conservation of energy as we know it in the form,

$$\frac{1}{2}mv^2 - \int_{x_0}^x F(x')dx' = E - V = \text{const.}$$

The integral in the above equation has a specific name in Physics: The work done by the force $F(x)$ over the distance traveled from x_0 to x . In other words, the above expression states that **the change in the kinetic energy of a system equals the work done by the force on it during the displacement**.

Problem 6. (Gravitational potential energy): Consider two-point masses, M and m , separated by a distance r . Newton's law of gravitation says that the force between them is attractive and has magnitude GMm/r^2 . Show that the potential energy of the system with respect to a separation distance r_0 of infinity is given by,

$$V(r) = -\frac{GMm}{r}$$

Problem 7. (Gravity near the earth): Show that the gravitational potential energy of a mass m at height y , relative to the ground, is given by,

$$V(R + y) - V(R) = mgy, y \ll R$$

where g is the gravity acceleration and R is the earth radius.

Problem 8. (Unwinding string): A mass is connected to one end of a massless string, the other end of which is connected to a very thin frictionless vertical pole. The string is initially wound completely around the pole, in a very large number of tiny horizontal circles, so that the mass touches the pole. The mass is released, and the string gradually unwinds. What angle does the string make with the pole at the moment it becomes completely unwound?

Conservation laws of Physics

Problem 9. (Raising a book): Assume that you lift a book up at constant speed, so there is no change in kinetic energy. However, we know that the potential energy of the book is increasing. Explain how this is consistent with the law of conservation of energy.

Problem 10. (Conservation of energy in three dimensions) Consider the same situation as in Problem 5 but in three dimensions. Show that the Newton's law leads to the same form of the law of conservation of energy but only expressed in terms of the 3D position vector of the particle $\mathbf{r}$.

