

Question Bank

Q1. Which of the following is a polynomial?

- A. $x^2 + 5/x + 2$
- B. $3x^2 - 4x + 7$
- C. $\sqrt{x} + 3$
- D. $1/(x^2 + 1)$

Q2. What is the degree of the polynomial $p(x) = 4x^3 - 2x^2 + 5x - 7$?

- A. 1
- B. 2
- C. 3
- D. 0

Q3. A polynomial of degree 2 is known as:

- A. Linear polynomial
- B. Quadratic polynomial
- C. Cubic polynomial
- D. Biquadratic polynomial

Q4. If $p(x) = x^2 - 3x + 2$, what is the value of $p(2)$?

- A. 0
- B. 1
- C. 2
- D. 4

Question Bank

Q5. A linear polynomial can have at most how many zeroes?

- A. 0
- B. 1
- C. 2
- D. 3

Q6. Consider the following table for a polynomial $p(x) = ax + b$:

x	-1	0	1
$p(x)$	-3	-1	1

What are the values of a and b ?

- A. $a=1$, $b=-1$
- B. $a=2$, $b=-1$
- C. $a=-1$, $b=1$
- D. $a=2$, $b=1$

Q7. If one zero of the polynomial $p(x) = x^2 - kx + 6$ is 3, then the value of k is:

- A. 2
- B. 5
- C. 6
- D. 9

Q8. Observe the graph of the polynomial $y = x^2 - 4$. Where does the graph intersect the x -axis?


- A. (2, 0) and (-2, 0)

Question Bank

- B. (4, 0) and (-4, 0)
- C. (0, 2) and (0, -2)
- D. (0, 4) and (0, -4)

Q9. For a quadratic polynomial $ax^2 + bx + c$, the sum of zeroes is given by:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$



Q10. If the zeroes of the quadratic polynomial $x^2 + (a+1)x + b$ are 2 and -3, then:

- A. $a = -7, b = -1$
- B. $a = 5, b = -1$
- C. $a = 2, b = -6$
- D. $a = 0, b = -6$

Question Bank

Q11. Look at the table for a polynomial $p(x)$:

x	-2	-1	0	1	2
$p(x)$	0	3	4	3	0

Based on the table, what are the zeroes of the polynomial?

- A. -1 and 1
- B. -2 and 2
- C. 0 and 4
- D. -2 and 0

Q12. What is the product of the zeroes of the polynomial $2x^2 - 7x + 3$?

- A. $\frac{7}{2}$
- B. $\frac{3}{2}$
- C. $-\frac{3}{2}$
- D. $-\frac{7}{2}$

Q13. A cubic polynomial can have at most how many zeroes?

- A. 1
- B. 2
- C. 3
- D. 4

Q14. If the graph of a polynomial $y = p(x)$ intersects the x -axis at exactly three points, then the degree of the polynomial is at least:

Question Bank

- A. 1
- B. 2
- C. 3
- D. 4

Q15. The zeroes of the polynomial $x^2 - 2x - 8$ are:

- A. 4, -2
- B. -4, 2
- C. 4, 2
- D. -4, -2

Q16. Identify the quadratic polynomial whose sum and product of zeroes are -3 and 2 respectively:

- A. $x^2 + 3x + 2$
- B. $x^2 - 3x + 2$
- C. $x^2 + 3x - 2$
- D. $x^2 - 3x - 2$

Q17. Which of the following is a cubic polynomial?

- A. $x^2 - 1$
- B. $x^3 - x$

Q18. Which of the following is not a polynomial?

- A. $x^2 + 3x + 2$

Question Bank

B. $x^2 + \sqrt{2}x + 5$

C. $x^2 + 1/x + 2$

D. $5x^3 - 4x + \sqrt{3}$

Q19. The degree of the polynomial $p(x) = 4x^3 - 3x^2 + 7x - 9$ is:

A. 1

B. 2

C. 3

D. 4

Q20. If $p(x) = x^2 - 2x - 3$, then the value of $p(3)$ is:

A. 0

B. 3

C. 6

D. -3

Q21. A quadratic polynomial can have at most how many zeroes?

A. 0

B. 1

C. 2

D. 3

Q22. For the polynomial $p(x) = x^2 - 2x - 8$, the zeroes are:

A. 4, -2

B. -4, 2

C. 4, 2

D. -4, -2

Question Bank

Q23. If alpha and beta are zeroes of the polynomial $p(x) = x^2 - 5x + 6$, then the value of $\alpha + \beta$ is:

- A. 5
- B. -5
- C. 6
- D. -6

Q24. If alpha and beta are zeroes of the polynomial $p(x) = x^2 - 5x + 6$, then the value of $\alpha * \beta$ is:

- A. 5
- B. -5
- C. 6
- D. -6

Q25. Observe the table for the polynomial $p(x) = x^2 - 3x - 4$:

x	-2	-1	0	1	2	3	4	5
---	----	----	---	---	---	---	---	---

p(x)	6	0	-4	-6	-6	-4	0	6
------	---	---	----	----	----	----	---	---

Based on the table, the zeroes of the polynomial are:

- A. -2 and 5
- B. -1 and 4
- C. 0 and -4
- D. 1 and 2

Q26. A quadratic polynomial whose sum of zeroes is -3 and product of zeroes is 2 is:

- A. $x^2 - 3x + 2$
- B. $x^2 + 3x + 2$
- C. $x^2 + 2x - 3$
- D. $x^2 - 2x + 3$


Question Bank

Q27. Consider the graph of the polynomial $y = x^2 - 4$. The points where the graph intersects the x-axis are:

- A. (0, 2) and (0, -2)
- B. (2, 0) and (-2, 0)
- C. (4, 0) and (-4, 0)
- D. (0, 4) and (0, -4)

Q28. The relationship between the zeroes and coefficients of a quadratic polynomial $ax^2 + bx + c$ is:

- A. Sum = c/a , Product = $-b/a$
- B. Sum = $-b/a$, Product = c/a
- C. Sum = b/a , Product = c/a
- D. Sum = $-b/a$, Product = $-c/a$



Q29. If one zero of the polynomial $p(x) = x^2 - kx + 6$ is 3, then the value of k is:

- A. 2
- B. 5
- C. 6
- D. 3

Question Bank

Q30. The number of zeroes of a linear polynomial is:

- A. 0
- B. 1
- C. 2
- D. 3

Q31. If the graph of a polynomial intersects the x-axis at 3 points, it is a:

- A. Linear polynomial
- B. Quadratic polynomial
- C. Cubic polynomial
- D. Biquadratic polynomial

Q32. Consider the following table for a polynomial $p(x)$:

x	$p(x)$
-1	0
0	-3
1	-4
2	-3
3	0

The zeroes of the polynomial are:

- A. -3 and 0
- B. -1 and 3
- C. 0 and 2
- D. -1 and 2

Q33. If α , β , and γ are the zeroes of a cubic polynomial $ax^3 + bx^2 + cx + d$, then $\alpha + \beta + \gamma$ is equal to:

Question Bank

Q34. If $p(x) = ax + b$, a is not equal to 0, is a linear polynomial, then its zero is given by:

- A. a/b
- B. $-b/a$
- C. b/a
- D. $-a/b$

Q35. The graph of the polynomial $y = ax + b$ is always a:

- A. Parabola
- B. Circle
- C. Straight line
- D. Hyperbola

Q36. A quadratic polynomial can have at most how many zeroes?

- A. 0
- B. 1
- C. 2
- D. 3

Q37. If the graph of a polynomial intersects the x -axis at exactly two points, then the number of zeroes of the polynomial is:

- A. 0
- B. 1

Question Bank

C. 2

D. 3

Q38. Consider the following table for a polynomial $p(x) = x^2 - 3x + 2$:

x	-1	0	1	2	3
$p(x)$	6	2	0	0	2

Based on the table, what are the zeroes of the polynomial?

A. -1 and 0

B. 1 and 2

C. 0 and 2

D. 1 and 3

Q39. If the sum of the zeroes of a quadratic polynomial $x^2 + bx + c$ is 5 and the product is 6, then the polynomial is:

A. $x^2 - 5x + 6$

B. $x^2 + 5x + 6$

C. $x^2 - 6x + 5$

D. $x^2 + 6x + 5$

Q40. Observe the graph of $y = x^2 - 4$. At which points does the graph intersect the x-axis?

A. (2, 0) and (0, -2)

B. (2, 0) and (-2, 0)


C. (0, 2) and (0, -2)

D. (-2, 0) and (0, 0)

Question Bank

Q41. If one zero of the polynomial $p(x) = (k-1)x^2 + kx + 1$ is -3 , then the value of k is:

- A. $\frac{4}{3}$
- B. $-\frac{4}{3}$
- C. $\frac{2}{3}$
- D. $-\frac{2}{3}$



Q42. Which of the following is a quadratic polynomial?

- A. $x + \frac{1}{x}$
- B. $x^2 + \frac{1}{x^2}$
- C. $\sqrt{3}x^2 + 5x + 7$
- D. $x^3 + 1$

Q43. The zeroes of the polynomial $x^2 - 2x - 8$ are:

- A. 4, -2

Question Bank

- B. -4, 2
- C. 4, 2
- D. -4, -2

Q44. Consider the following table for a linear polynomial $p(x) = ax + b$:

x	-2	0	2
$p(x)$	-1	3	7

What is the zero of this polynomial?

- A. -1.5
- B. 1.5
- C. -3
- D. 3

Q45. If α and β are the zeroes of the polynomial $p(x) = ax^2 + bx + c$, then $\alpha * \beta$ is:

- A. $-b/a$
- B. b/a
- C. c/a
- D. $-c/a$

Q46. A cubic polynomial has at most:

- A. 1 zero
- B. 2 zeroes
- C. 3 zeroes
- D. 4 zeroes

Question Bank

Q47. If the graph of a polynomial does not intersect the x-axis at any point, then the number of real zeroes is:

- A. 0
- B. 1
- C. 2
- D. Infinitely many

Q48. The zeroes of the polynomial $p(x) = x^2 - 15$ are:

- A. 15, -15
- B. $\sqrt{15}$, $-\sqrt{15}$
- C. 0, 15
- D. 0, -15

Q49. If the sum of the zeroes of the polynomial $p(x) = kx^2 - 3x + 5$ is 1, then the value of k is:

- A. 2
- B. 3
- C. 4
- D. 5

Q50. Observe the graph of $y =$

Q51. If $p(x) = ax + b$, a is not equal to 0, is a linear polynomial, then its zero is given by:

Question Bank

- A. a/b
- B. $-a/b$
- C. $-b/a$
- D. b/a

Q52. The degree of a cubic polynomial is:

- A. 1
- B. 2
- C. 3
- D. 0

Q53. A quadratic polynomial can have at most how many zeroes?

- A. 0
- B. 1
- C. 2
- D. 3

Q54. If one of the zeroes of the quadratic polynomial $(k-1)x^2 + kx + 1$ is -3 , then the value of k is:

- A. $4/3$
- B. $-4/3$
- C. $2/3$
- D. $-2/3$

Q55. Consider the polynomial $p(x) = x^2 - 2x - 8$. The zeroes of the polynomial are:

Question Bank

- A. 2, -4
- B. 4, -2
- C. -2, -4
- D. 2, 4

Q56. For the polynomial $p(x) = 2x^2 - 8x + 6$, the sum of the zeroes is:

- A. 3
- B. 4
- C. -4
- D. 6

Q57. For the polynomial $p(x) = 2x^2 - 8x + 6$, the product of the zeroes is:

- A. 3
- B. 4
- C. 8
- D. 6

Q58. Observe the following table for $p(x) = x^2 - 3x - 4$. Identify the row that correctly represents a zero of the polynomial.

$x \mid p(x)$

- A. -1 | 0
- B. 0 | -4
- C. 1 | -6
- D. 2 | -6

Question Bank

Q59. If alpha and beta are the zeroes of the polynomial $ax^2 + bx + c$, then the value of $\alpha + \beta$ is:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$

Q60. If alpha and beta are the zeroes of the polynomial $ax^2 + bx + c$, then the value of $\alpha * \beta$ is:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$

Q61. A quadratic polynomial whose sum of zeroes is -3 and product of zeroes is 2 is:

- A. $x^2 - 3x + 2$
- B. $x^2 + 3x + 2$
- C. $x^2 + 3x - 2$
- D. $x^2 - 3x - 2$

Q62. If the zeroes of the quadratic polynomial $x^2 + (a+1)x + b$ are 2 and -3, then:

- A. $a = -7, b = -1$
- B. $a = 5, b = -1$
- C. $a = 2, b = -6$
- D. $a = 0, b = -6$


Question Bank

Q63. Observe the graph of $y = x^2 - 4$. The zeroes of the polynomial are the points where the graph intersects:

- A. The y-axis
- B. The x-axis
- C. The origin
- D. The line $y = x$

Q64. The number of zeroes of the polynomial $y = x^3 - x$ is:

- A. 1
- B. 2
- C. 3
- D. 0



Q65. If the graph of a polynomial intersects the x-axis at exactly two points, then it is:

- A. Always a linear polynomial
- B. Always a quadratic polynomial

Question Bank

- C. A polynomial of degree at least 2
- D. A polynomial of degree exactly 3

Q66. Consider the table below for a polynomial $p(x)$. Based on the values, which of the following is a zero of the polynomial?

$x \mid p(x)$

-2 $\mid$ 5

-1 $\mid$ 0

0 $\mid$ -3

1 $\mid$ -4

2 $\mid$ -3

- A. -2
- B. -1
- C. 0
- D. 1

Q67. The quadratic polynomial $p(x) = x^2 + 7x + 10$ has zeroes:

- A. 2 and 5
- B. -2 and 5
- C. -2 and -5
- D. 2 and -5

Q68. If the sum of the

Question Bank

Q69. Which of the following is a polynomial?

- A. $x^2 - 3x + 5$
- B. $x^2 + 1/x + 2$
- C. square root of $x + 3$
- D. $x^{-2} + 5x + 6$

Q70. The degree of the polynomial $4x^3 - 2x^2 + 7x - 5$ is:

- A. 1
- B. 2
- C. 3
- D. 4

Q71. If $p(x) = x^2 - 2x - 3$, then the value of $p(3)$ is:

- A. 0
- B. 3
- C. 6
- D. -3

Q72. A quadratic polynomial whose sum of zeroes is 0 and product of zeroes is -5 is:

- A. $x^2 - 5$
- B. $x^2 + 5$
- C. $x^2 + 5x$
- D. $x^2 - 5x$

Question Bank

Q73. If one zero of the polynomial $x^2 - 4x + 1$ is $2 + \sqrt{3}$, then the other zero is:

- A. $2 - \sqrt{3}$
- B. $2 + \sqrt{3}$
- C. $-2 + \sqrt{3}$
- D. $-2 - \sqrt{3}$

Q74. Observe the following table representing values of $p(x) = x^2 - 3x - 4$ for different values of x .

x	-1	0	1	2	3	4
$p(x)$	0	-4	-6	-6	-4	0

Based on the table, the zeroes of the polynomial are:

- A. -1 and 4
- B. 0 and -4
- C. 1 and 2
- D. 3 and 4

Q75. If the graph of $y = ax^2 + bx + c$ intersects the x -axis at exactly two distinct points, then the quadratic polynomial has:

- A. No zeroes
- B. One zero
- C. Two distinct zeroes
- D. More than two zeroes


Q76. Observe the graph of $y = x^2 - 4$. How many times does it intersect the x -axis?

Question Bank

- A. 0
- B. 1
- C. 2
- D. 3

Q77. The zeroes of the quadratic polynomial $x^2 + 7x + 10$ are:

- A. 2 and 5
- B. -2 and -5
- C. -2 and 5
- D. 2 and -5



Q78. For a polynomial $p(x) = ax^2 + bx + c$, the sum of zeroes is given by:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$

Question Bank

Q79. The product of zeroes of the polynomial $2x^2 - 7x + 3$ is:

- A. $\frac{3}{2}$
- B. $-\frac{3}{2}$
- C. $\frac{7}{2}$
- D. $-\frac{7}{2}$

Q80. If a and b are zeroes of $x^2 - 5x + k$ such that $a - b = 1$, the value of k is:

- A. 4
- B. 5
- C. 6
- D. 2

Q81. Consider the table of a linear polynomial $p(x) = ax + b$:

x	-2	0	2
$p(x)$	-7	-1	5

What are the values of a and b ?

- A. $a=3, b=-1$
- B. $a=2, b=-1$
- C. $a=3, b=1$
- D. $a=2, b=1$

Q82. A cubic polynomial can have at most:

- A. 1 zero

Question Bank

- B. 2 zeroes
- C. 3 zeroes
- D. 0 zeroes

Q83. The graph of $y = x^2 + 1$:

- A. Intersects the x-axis at two points
- B. Touches the x-axis at one point
- C. Does not intersect the x-axis
- D. Passes through the origin

Q84. If the sum of zeroes of the polynomial $kx^2 + 2x + 3k$ is equal to their product, then k is:

- A. $\frac{1}{3}$
- B. $-\frac{1}{3}$
- C. $\frac{2}{3}$
- D. $-\frac{2}{3}$

Q85. The quadratic polynomial whose zeroes are 0 and 5 is:

Q86. Which of the following is a polynomial?

- A. $x^2 - 3x + 5$
- B. $x^2 + \frac{1}{x} + 2$
- C. $\sqrt{x} + 3$
- D. $x^3 + x^{-1} + 5$

Question Bank

Q87. The degree of the polynomial $p(x) = 4x^3 - 2x^2 + 7x - 9$ is:

- A. 1
- B. 2
- C. 3
- D. 4

Q88. A polynomial of degree 1 is called:

- A. Quadratic
- B. Linear
- C. Cubic
- D. Bi-quadratic

Q89. If $p(x) = x^2 - 2x - 3$, then the value of $p(3)$ is:

- A. 0
- B. 3
- C. -3
- D. 6

Q90. The number of zeroes that a cubic polynomial can have is:

- A. At most 1
- B. At most 2
- C. At most 3
- D. Exactly 3

Question Bank

Q91. Consider the polynomial $p(x) = x^2 - 4$. The zeroes of the polynomial are:

- A. 2, -2
- B. 4, -4
- C. 2, 2
- D. -2, -2

Q92. If one zero of the polynomial $p(x) = x^2 - kx + 6$ is 3, then the value of k is:

- A. 2
- B. 5
- C. -5
- D. 6

Q93. Observe the following table of values for a polynomial $p(x) = ax + b$:

x	-1	0	1
$p(x)$	-2	1	4

The values of a and b are:

- A. $a=3$, $b=1$
- B. $a=1$, $b=3$
- C. $a=2$, $b=1$
- D. $a=3$, $b=0$

Q94. The sum of the zeroes of the quadratic polynomial $x^2 + 7x + 10$ is:

Question Bank

- A. 7
- B. -7
- C. 10
- D. -10

Q95. The product of the zeroes of the quadratic polynomial $x^2 - 5x + 6$ is:

- A. 5
- B. -5
- C. 6
- D. -6

Q96. If α and β are the zeroes of the polynomial $x^2 - 5x + k$ such that $\alpha - \beta = 1$, the value of k is:

- A. 4
- B. 5
- C. 6
- D. 2

Q97. Observe the following table representing a polynomial $p(x)$:

x	-2	-1	0	1	2
$p(x)$	0	-3	-4	-3	0

The zeroes of the polynomial based on the table are:

- A. -2 and 2

Question Bank

- B. -1 and 1
- C. -4 and 0
- D. -2 and 0

Q98. A quadratic polynomial whose sum of zeroes is 0 and product of zeroes is -5 is:

- A. $x^2 - 5$
- B. $x^2 + 5$
- C. $x^2 - 5x$
- D. $x^2 + 5x$

Q99. If the graph of $y = ax^2 + bx + c$ intersects the x-axis at two distinct points, the quadratic polynomial has:

- A. No zeroes
- B. Two distinct zeroes
- C. One zero
- D. Three zeroes


Q100. Consider the graph of $y = x^2 - 3x + 2$. The points where the graph intersects the x-axis are:

- A. (1, 0) and (2, 0)
- B. (-1, 0) and (-2, 0)
- C. (0, 1) and (0, 2)
- D. (1, 0) and (-2, 0)

Q101. If the graph of a polynomial $p(x)$ does not intersect the x-axis at any point, then the number of zeroes of $p(x)$ is:

Question Bank

- A. 0
- B. 1
- C. 2
- D. 3



Q102. The zeroes of the polynomial $p(x) = x^2 + 7x + 12$ are:

- A. -3, -4
- B. 3, 4
- C. -

Q103. If $p(x) = ax + b$, a is not equal to 0, is a linear polynomial, then its zero is given by:

- A. a/b
- B. $-b/a$
- C. b/a
- D. $-a/b$

Question Bank

Q104. A quadratic polynomial can have at most how many zeroes?

- A. 0
- B. 1
- C. 2
- D. 3

Q105. If the graph of a polynomial $y = p(x)$ intersects the x-axis at exactly three points, then the degree of the polynomial is:

- A. 1
- B. 2
- C. 3
- D. 4

Q106. Consider the following table representing the values of $p(x) = x^2 - 3x - 4$ for different values of x :

x	-2	-1	0	1	2	3	4	5
$p(x)$	6	0	-4	-6	-6	-4	0	6

Based on the table, the zeroes of the polynomial are:

- A. -2 and 5
- B. 0 and -4
- C. -1 and 4
- D. 1 and 2

Question Bank

Q107. If alpha and beta are the zeroes of the polynomial $p(x) = x^2 - 5x + 6$, then the value of $\alpha + \beta$ is:

- A. 5
- B. -5
- C. 6
- D. -6

Q108. If alpha and beta are the zeroes of the polynomial $p(x) = x^2 - 5x + 6$, then the value of $\alpha * \beta$ is:

- A. 5
- B. -5
- C. 6
- D. -6

Q109. The quadratic polynomial whose sum of zeroes is -3 and product of zeroes is 2 is:

- A. $x^2 - 3x + 2$
- B. $x^2 + 3x + 2$
- C. $x^2 + 2x - 3$
- D. $x^2 - 2x + 3$


Q110. Observe the graph of $y = x^2 - 4$. How many times does the graph intersect the x-axis?

- A. 0
- B. 1
- C. 2
- D. 3

Question Bank

Q111. For the polynomial $p(x) = ax^2 + bx + c$, the sum of zeroes is:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$



Q112. A cubic polynomial $x^3 - x$ has how many zeroes?

- A. 1
- B. 2
- C. 3
- D. 0

Q113. The value of the polynomial $p(x) = 5x - 4x^2 + 3$ at $x = -1$ is:

- A. -6
- B. -2
- C. 2

Question Bank

D. 12

Q114. Consider the table for $p(x) = x^2 - 2x - 3$:

x	-2	-1	0	1	2	3	4
$p(x)$	5	0	-3	-4	-3	0	5

Which of the following are the zeroes of the polynomial?

A. -2 and 4

B. -1 and 3

C. 0 and -3

D. 1 and 2

Q115. If one zero of the polynomial $p(x) = (k-1)x^2 + kx + 1$ is -3, then the value of k is:

A. $\frac{4}{3}$

B. $-\frac{4}{3}$

C. $\frac{2}{3}$

D. $-\frac{2}{3}$

Q116. The graph of a quadratic polynomial $y = ax^2 + bx + c$ is a parabola opening downwards if:

A. $a > 0$

B. $a < 0$

C. $a = 0$

D. $a = 1$

Question Bank

Q117. The number of zeroes of the polynomial $y = x^2 + 1$ is:

- A. 0
- B. 1
- C. 2
- D. 3

Q118. If the sum of the zeroes of the polynomial $p(x) = 2x^2 - 8x + 6$ is $\alpha + \beta$, then the value of $\alpha + \beta$ is:

- A. 8
- B. 4
- C. 3
- D. -4

Q1

Q119. Which of the following is a quadratic polynomial?

- A. $x + 5$
- B. $x^2 + 3x + 2$
- C. $x^3 + 2x^2 + x + 1$
- D. $\sqrt{x} + 2$

Q120. If $p(x) = ax^2 + bx + c$, then the condition for it to be a quadratic polynomial is:

- A. $a = 0$

Question Bank

B. a is not equal to 0

C. $b = 0$

D. $c = 0$

Q121. The maximum number of zeroes a cubic polynomial can have is:

A. 1

B. 2

C. 3

D. 4

Q122. If the graph of a polynomial $y = p(x)$ intersects the x -axis at two distinct points, then the number of zeroes of the polynomial is:

A. 0

B. 1

C. 2

D. 3

Q123. Consider the polynomial $p(x) = x^2 - 4x + 3$. Identify the zeroes from the following table:

x	-1	0	1	2	3	4
$p(x)$	8	3	0	-1	0	3

A. 0 and 3

B. 1 and 3

C. 2 and 4

D. -1 and 0

Question Bank

Q124. For a quadratic polynomial $ax^2 + bx + c$, the sum of zeroes is given by:

- A. c/a
- B. $-c/a$
- C. b/a
- D. $-b/a$

Q125. If α and β are the zeroes of $x^2 - 7x + 10$, then the product of zeroes ($\alpha * \beta$) is:

- A. 7
- B. -7
- C. 10
- D. -10

Q126. A linear polynomial has a degree of:

- A. 0
- B. 1
- C. 2
- D. 3

Q127. Observe the following table for a polynomial $p(x) = ax + b$:

x	-2	-1	0	1	2
$p(x)$	-7	-4	-1	2	5

What is the zero of this polynomial?

- A. $1/3$

Question Bank

- B. $\frac{1}{2}$
- C. $\frac{1}{4}$
- D. $\frac{1}{5}$

Q128. If α , β , and γ are the zeroes of a cubic polynomial $ax^3 + bx^2 + cx + d$, then $\alpha * \beta * \gamma$ is:

- A. $\frac{d}{a}$
- B. $-\frac{d}{a}$
- C. $\frac{c}{a}$
- D. $-\frac{c}{a}$

Q129. The graph of $y = x^2 - 4$ represents:

- A. A straight line
- B. A parabola opening upwards
- C. A parabola opening downwards
- D. A circle

Q130. If one zero of the polynomial $x^2 - 3x + k$ is 2, then the value of k is:

- A. 1
- B. 2
- C. 3
- D. 4

Q131. Which of the following is not a polynomial?

Question Bank

- A. $x^2 + 1$
- B. $x^3 - 5x + 7$
- C. $x^2 + 1/x$
- D. 5

Q132. The sum of zeroes of the polynomial $2x^2 - 8x + 6$ is:

- A. 4
- B. -4
- C. 3
- D. -3

Q133. Given the following observation table for $y = x^2 - 2x - 8$:

x	-2	-1	0	1	2	4
y	0	-5	-8	-9	-8	0

Identify the zeroes of the polynomial from the table:

- A. -2 and 4
- B. -1 and 2
- C. 0 and -8
- D. 1 and -9

Q134. If the sum of the zeroes of the polynomial $kx^2 + 2x + 3k$ is equal to their product, then the value of k is:

- A. $-2/3$
- B. $2/3$

Question Bank

C. $\frac{3}{2}$

D. $-\frac{3}{2}$

Q135. A polynomial of degree 3 is called:

A. Linear

B. Quadratic

C. Cubic

D. Biquadratic

Q18