

AGA KHAN UNIVERSITY EXAMINATION BOARD

HIGHER SECONDARY SCHOOL CERTIFICATE

CLASS XI




















MODEL EXAMINATION PAPER 2018

Chemistry Paper I

Time: 50 minutes Marks: 35

INSTRUCTIONS


1. Read each question carefully.
2. Answer the questions on the separate answer sheet provided. DO NOT write your answers on the question paper.
3. There are 100 answer numbers on the answer sheet. Use answer numbers 1 to 35 only.
4. In each question there are four choices A, B, C, D. Choose ONE. On the answer grid black out the circle for your choice with a pencil as shown below.

Correct Way				Incorrect Ways				
1				1				
				2				
				3				
				4				

Candidate's Signature


5. If you want to change your answer, ERASE the first answer completely with a rubber, before blacking out a new circle.
6. DO NOT write anything in the answer grid. The computer only records what is in the circles.
7. You may use a scientific calculator if you wish.

1. The number of moles of Ca(OH)_2 that completely reacts with 50 g of NH_4Cl to produce 15.87 g of NH_3 gas in the given reaction is




(Note: Atomic mass of C = 12 amu, H = 1 amu, N = 14 amu, Cl = 35.5 amu and Ca = 40 amu)

- A. 0.467
 B. 0.934
 C. 1.868
 D. 3.736
2. Methane burns exothermically in the presence of free oxygen at standard temperature and pressure (STP) as shown in the given chemical equation.



If 25 moles of CH_4 combust to give equal moles of CO_2 , then the volume of CO_2 formed will be

- A. 0.897 dm^3
 B. 22.414 dm^3
 C. 25.00 dm^3
 D. 560.35 dm^3
3. Sodium metal vigorously reacts with water to form sodium hydroxide and hydrogen gas as shown in the given equation.



An experiment under controlled environment in a laboratory gives 80% yield of sodium hydroxide on reacting 2 moles of sodium metal with 2 moles of water. How much actual yield of sodium hydroxide in grams is obtained in the given reaction?

- A. 32 g
 B. 50 g
 C. 64 g
 D. 100 g
4. In the discharge tube, the original glow disappears when the pressure inside the tube is reduced to
- A. 0.01 mm Hg
 B. 0.5 cm Hg
 C. 1 mm Hg
 D. 1 cm Hg

5. When rapidly moving electrons collide with heavy metal anode in the discharge tube, it produces
- X-rays.
 - gamma rays.
 - cathode rays.
 - positive rays.
6. Visible light is NOT used to determine the position of electrons because the wavelength of visible light is millions of times larger as compared to the
- energy of electrons.
 - velocity of electrons.
 - diameter of electrons.
 - momentum of electrons.
7. Mixing of acetic acid with sodium acetate will give buffer with pH
- < 7.0
 - > 7.0
 - $= 7.0$
- I only.
 - II only.
 - III only.
 - II and III.
8. If CH_3COONa is added in excess to CH_3COOH solution, then the
- pH of solution will increase.
 - pH of solution will decrease.
 - dissociation of CH_3COOH will increase.
 - dissociation of CH_3COONa will decrease.
9. In which of the following molecules, the highlighted carbon atom contains two unhybridised p-orbitals?
- CH_4
 - $\text{CH}=\text{CH}$
 - CH_3-CH_3
 - $\text{CH}_2=\text{CH}_2$
10. Which of the following bonds between two atoms has the LARGEST bond length?
- B-F
 - B-Cl
 - Si-H
 - Si-F

PLEASE TURN OVER THE PAGE

11. Ionic compounds are unable to show isomerism because in ionic compounds the bonds are
- rigid and directional.
 - non-rigid and directional.
 - rigid and non-directional.
 - non-rigid and non-directional.
12. If the electronegativity difference increases between hydrogen and halogen in a hydrogen halide molecule, then what will be the effect on its ionic character and bond strength?


	Ionic Character	Bond Strength
A	Increases	Increases
B	Increases	Decreases
C	Decreases	Increases
D	Decreases	Decreases

13. The density of $\text{CH}_{4(g)}$ at 17°C and at 1 atmospheric pressure is
- (Note: Atomic mass of C = 12 amu, H = 1 amu, and the value of $R = 0.0821 \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$)
- 0.67 g/dm^3
 - 11.46 g/dm^3
 - 22.32 g/dm^3
 - 380.94 g/dm^3
14. According to Charles's law, if a gas is warmed by 1°C at constant pressure, it will expand
- twice its original volume at 0°C .
 - twice its original volume at 100°C .
 - $1/273$ of its original volume at 0°C .
 - $1/273$ of its original volume at 100°C .
15. When water is poured into a glass test tube, the surface of the water is concaved. This meniscus forms because of
- weak adhesive forces.
 - strong adhesive forces.
 - strong cohesive forces.
 - equal adhesive and cohesive forces.
16. Liquid crystals have wide practical application due to their optical and electrical properties.
- Which of the following is INCORRECT about the use of liquid crystals?
- They can be used as temperature sensors.
 - They are used in the dial of analogue watches.
 - They are used in skin thermography to detect blockage of veins.
 - They can be used to find the point of potential failure in electrical circuits.

17. Graphite exists in the form of layers. It is a conductor parallel to the layers but insulator across the layers. This property of graphite is known as
- symmetry.
 - anisotropy.
 - isomorphism.
 - polymorphism.
18. In ionic solids, ions are held together by strong electrostatic force of attraction. However, ionic crystals are highly brittle because they
- have low density.
 - have loose packing of ions.
 - exist as neutral independent molecules.
 - contain opposite ions in parallel layers.
19. If, in a unit cell, all angles are right angles and all edges are of equal length, the crystal system must be
- cubic.
 - triclinic.
 - monoclinic.
 - orthorhombic.
20. The equilibrium constant (K_c) for the decomposition of HF at 2000°C is 10^{-13} which indicates that it has

	Stability	Rate of Decomposition
A	High	Low
B	Low	High
C	High	High
D	Low	Low

21. Consider the given reaction at equilibrium.




Which of the following would be the result if the concentration of oxygen is increased?

	$[\text{SO}_2]$	$[\text{SO}_3]$	Equilibrium shift
A	Increased	Decreased	Right
B	Decreased	Increased	Right
C	Increased	Decreased	Left
D	Decreased	Increased	Left

PLEASE TURN OVER THE PAGE

22. The ionisation of KClO_3 in the given reaction can be suppressed by adding



- A. KCl
 B. HCl
 C. NaCl
 D. NH_4Cl
23. An activated complex is a combination of all the atoms involved in the reaction. It
- A. is a short lived species.
 B. is a highly stable species.
 C. slowly decomposes into products.
 D. involves atoms having minimum energy.
24. The order of reaction can be deduced by simply observing the
- A. number of reactants involved in a reaction only.
 B. time taken to process the chemical reaction only.
 C. rate expression which is obtained experimentally.
 D. chemical equation showing the slow and fast steps.
25. The rate of a chemical reaction can be retarded by adding a/ an
- A. indicator.
 B. activator.
 C. inhibitor.
 D. co-factor.
26. A solution is prepared by mixing three different components, **P**, **Q** and **R**. If the number of moles of each component is $n_{\text{P}} = 4$ moles, $n_{\text{Q}} = 6$ moles, $n_{\text{R}} = 10$ moles, then the mole percent of each component will be

	Mole Percent		
	P	Q	R
A	20	30	50
B	0.2	0.3	0.5
C	0.4	0.6	0.10
D	40	60	100

27. When a non-volatile solute is added to a pure solvent, the
- A. boiling point of the solution decreases.
 B. freezing point of the solution increases.
 C. vapour pressure of the solvent increases.
 D. rate of evaporation of the solvent decreases.

28. Amongst the given set of liquids, the completely miscible set is
- A. aniline and water.
 - B. aniline and hexane.
 - C. cyclohexane and water.
 - D. benzene and cyclohexane.
29. What is the molality of a solution prepared by dissolving 148 g of butanol ($C_4H_{10}O$) in 0.1 kg of benzene?
- (Note: Atomic mass of C = 12 amu, O = 16 amu, H = 1 amu)
- A. 20 mol/kg
 - B. 2 mol/kg
 - C. 0.2 mol/kg
 - D. 0.05 mol/kg
30. In thermochemistry, a liquid whose boiling point is to be determined is considered as a/ an
- A. system.
 - B. final state.
 - C. initial state.
 - D. surrounding.
31. Which of the following is the CORRECT formula for calculating the amount of heat evolved during a neutralisation reaction in a glass calorimeter?
- A. Quantity of heat = mass \times heat capacity \times temperature
 - B. Quantity of heat = mass \times specific heat capacity \times temperature
 - C. Quantity of heat = mass \times specific heat capacity \times change in temperature
 - D. Quantity of heat = mass \times specific latent heat of fusion \times change in temperature
32. If the specific heat capacity of copper is $387 \text{ J/kg/}^\circ\text{C}$, then how much energy is needed to raise the temperature of 400 g of copper from 30°C to 55°C ?
- A. 3.870 kJ
 - B. 8.514 kJ
 - C. 3870 kJ
 - D. 8514 kJ
33. Iron is extracted from its ore, haematite, in a blast furnace which involves all of the following chemical reactions. In which of these reactions is carbon reduced?
- A. $C_{(s)} + O_{2(g)} \rightarrow CO_{2(g)}$
 - B. $CO_{2(g)} + C_{(s)} \rightarrow 2CO_{(g)}$
 - C. $CaCO_{3(s)} \rightarrow CaO_{(s)} + CO_{2(g)}$
 - D. $Fe_2O_{3(s)} + 3CO_{(g)} \rightarrow 2Fe_{(s)} + 3CO_{2(g)}$

PLEASE TURN OVER THE PAGE

34. Which of the following statements best defines standard electrode potential?

- A. The potential set up when an electrode undergoes reduction.
- B. The potential set up when an electrode undergoes oxidation.
- C. The potential of an electrode in 1M solution at 25°C and 1atm.
- D. The potential of an electrode in 1M solution at 100°C and 1atm.

35. The reaction between $\text{Cr}_2\text{O}_7^{-2}$ and H_2SO_3 gives the following half reactions.

Reduction half reaction: $\text{Cr}_2\text{O}_7^{-2} \rightarrow \text{Cr}^{+3}$

Oxidation half reaction: $\text{H}_2\text{SO}_3 \rightarrow \text{HSO}_4^{-}$

The equation which represents the balanced reduction half reaction is

- A. $\text{Cr}_2\text{O}_7^{-2} + 14\text{H}^+ + 3\text{e}^- \rightarrow \text{Cr}^{+3} + 7\text{H}_2\text{O}$
- B. $\text{Cr}_2\text{O}_7^{-2} + 14\text{H}^+ + 6\text{e}^- \rightarrow \text{Cr}^{+3} + 7\text{H}_2\text{O}$
- C. $\text{Cr}_2\text{O}_7^{-2} + 14\text{H}^+ + 3\text{e}^- \rightarrow 2\text{Cr}^{+3} + 7\text{H}_2\text{O}$
- D. $\text{Cr}_2\text{O}_7^{-2} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{+3} + 7\text{H}_2\text{O}$

Please use this page for rough work

AKU-EB
Model Paper 2018
for Teaching & Learning

Please use this page for rough work

AKU-EB
Model Paper 2018
for Teaching & Learning

Please use this page for rough work

AKU-EB
Model Paper 2018
for Teaching & Learning

Please use this page for rough work

AKU-EB
Model Paper 2018
for Teaching & Learning