

Module 4 :

→ Principles of counting - II

- Principle of Inclusion and Exclusion.
- Derangements.
- Rook polynomials.

→ Recurrence Relations

- first order linear Recurrence Relation.
- second order linear homogeneous Recurrence relⁿ with constant coefficient

Recurrence Relations

+ First order Recurrence relations:

A 1st order recurrence relation with constant coefficients is of the form

$$a_n = C a_{n-1} + f(n) \rightarrow \textcircled{1} \quad \text{for } n \geq 1.$$

where $C \rightarrow$ a known constant.

$f(n) \rightarrow$ a known function of n .

If $f(n) = 0$, then $\textcircled{1}$ is called a homogeneous recurrence relation; otherwise it is called a non-homogeneous recurrence relation.

Note:- 1) The General solⁿ of a homogeneous R.R is $a_n = C^n a_0$ for $n \geq 1$.

2) The General solⁿ of a non-homogeneous R.R of order 1 is given by $a_n = C^n a_0 + \sum_{k=1}^n C^{n-k} f(k)$, for $n \geq 1$.

Problems:-

1) Solve the recurrence relation $a_n = 7a_{n-1}$, where $n \geq 1$ given that $a_2 = 98$.

Solⁿ:- $a_n = 7a_{n-1} \rightarrow \textcircled{1}$ is a homogeneous 1st order recurrence relation.

General solⁿ is given by

$$a_n = C^n a_0.$$

$$a_n = 7^n a_0 \rightarrow \textcircled{2}$$

$$\text{for } n=2; a_2 = 7^2 a_0 \Rightarrow a_2 = 49 a_0$$

$$\Rightarrow 98 = 49 a_0 \Rightarrow \boxed{a_0 = 2}$$

Sub in $\textcircled{2}$, $\boxed{a_n = 2 \cdot 7^n}$ is the General solⁿ.

2) Solve the recurrence relation $a_n = n a_{n-1}$ for $n \geq 1$

given that $a_0 = 1$.

Soln:- $a_n = n a_{n-1}$

$$n=1; a_1 = 1 \times a_0$$

$$n=2; a_2 = 2 \times a_1 = (2 \times 1) \times a_0$$

$$n=3; a_3 = 3 \times a_2 = (3 \times 2 \times 1) \times a_0$$

$$n=4; a_4 = 4 \times a_3 = (4 \times 3 \times 2 \times 1) \times a_0 \text{ and so on.}$$

\therefore General soln is

$$a_n = n! a_0 \text{ for } n \geq 1.$$

using $a_0 = 1 \Rightarrow \boxed{a_n = n!}$ is the required soln.

3) Solve the recurrence relation $a_n - 3a_{n-1} = 5 \times 3^n$ for $n \geq 1$,

given that $a_0 = 2$.

Soln:- Given $a_n = 3a_{n-1} + (5 \times 3^n) \rightarrow \textcircled{1}$ is a non-homo.

relation with $c = 3$, $f(n) = 5 \times 3^n$.

General soln is given by

$$a_n = 3^n a_0 + \sum_{k=1}^n 3^{n-k} f(k)$$

$$a_n = 3^n a_0 + 3^{n-1} f(1) + 3^{n-2} f(2) + \dots + 3^0 f(n)$$

$$\Rightarrow a_n = 3^n \times 2 + 3^{n-1} (5 \times 3^1) + 3^{n-2} (5 \times 3^2) + \dots + 3^0 \times (5 \times 3^n)$$

$$= 2 \times 3^n + 5 \times [3^n + 3^n + \dots + 3^n] \text{ (n times)}$$

$$= 2 \times 3^n + 5 \times n \times 3^n$$

$$\boxed{a_n = 3^n (2 + 5n)} \text{ is the required soln.}$$

4) Solve the recurrence relation $a_n - 3a_{n-1} = 5 \times 7^n$, for $n \geq 1$,

given that $a_0 = 2$.

P.T.O.

Soln :: Given: $a_n = 3a_{n-1} + (5 \times 7^n) \rightarrow \textcircled{1}$ is a non-homo.
 recurrence relation with $c=3$, $f(n) = 5 \times 7^n$.

The general soln is given by

$$a_n = 3^n a_0 + \sum_{k=1}^n 3^{n-k} f(k).$$

$$= 3^n \times 2 + \sum_{k=1}^n 3^{n-k} \times (5 \times 7^k)$$

$$= 2 \times 3^n + (5 \times 3^n) \sum_{k=1}^n 3^{-k} \cdot 7^k.$$

$$= 2 \times 3^n + (5 \times 3^n) \sum_{k=1}^n \left(\frac{7}{3}\right)^k.$$

$$= 2 \times 3^n + (5 \times 3^n) \left[\frac{7}{3} + \left(\frac{7}{3}\right)^2 + \dots + \left(\frac{7}{3}\right)^n \right]$$

$$= 2 \times 3^n + (5 \times 3^n) \times \frac{7}{3} \left[1 + \left(\frac{7}{3}\right) + \left(\frac{7}{3}\right)^2 + \dots + \left(\frac{7}{3}\right)^{n-1} \right]$$

$$a + ar + ar^2 + \dots = \frac{a(r^n - 1)}{r - 1}; r > 1$$

here $a=1$, $r=7/3$

$$\therefore a_n = 2 \times 3^n + (35 \times 3^{n-1}) \left[\frac{\left(\frac{7}{3}\right)^n - 1}{\left(\frac{7}{3} - 1\right)} \right]$$

$$= (2 \times 3^n) + (35 \times 3^{n-1}) \times \frac{3}{4} \left[\frac{7^n - 3^n}{3^n} \right]$$

$$= (2 \times 3^n) + \left(\frac{35}{4}\right) (7^n - 3^n)$$

$$= 3^n \left[2 - \frac{35}{4} \right] + \frac{35}{4} \cdot 7^n$$

$$= -3^n \times \frac{27}{4} + \frac{5 \times 7 \times 7^n}{4}$$

$$= -\frac{1}{4} \cdot 3^{n+3} + \frac{5}{4} \cdot 7^{n+1}$$

$$\boxed{a_n = \frac{1}{4} [5 \times 7^{n+1} - 3^{n+3}]}$$

is the required soln.

5) Solve the recurrence relation

$$a_n = 2a_{n/2} + (n-1) \quad \text{for } n = 2^k, \quad k \geq 1, \quad \text{given } a_1 = 0.$$

Soln: $a_n = 2a_{n/2} + (n-1)$

$$\Rightarrow a_n - 2a_{n/2} = (n-1).$$

we obtain the following successive eqns

$$a_{n/2} - 2a_{n/4} = \left(\frac{n}{2} - 1\right)$$

$$a_{n/4} - 2a_{n/8} = \left(\frac{n}{4} - 1\right)$$

⋮

$$a_{n/2^{k-1}} - 2a_{n/2^k} = \left(\frac{n}{2^{k-1}} - 1\right)$$

These can be written as

$$a_n - 2a_{n/2} = (n-1).$$

$$2a_{n/2} - 2^2a_{n/4} = (n-2)$$

$$2^2a_{n/4} - 2^3a_{n/8} = (n-2^2)$$

⋮

$$2^{k-1}a_{n/2^{k-1}} - 2^k a_{n/2^k} = (n-2^{k-1}).$$

adding these, we get

$$a_n - 2^k a_{n/2^k} = (n-1) + (n-2) + (n-2^2) + \dots + (n-2^{k-1})$$

since $n = 2^k$, $a_{n/2^k} = a_1 = 0$ (given)

$$\therefore a_n = (n+n+\dots+n)_{k \text{ times}} - \frac{(1+2+2^2+\dots+2^{k-1})}{a+a^2+a^3+\dots} = \frac{a(r^k-1)}{r-1}, \quad r > 1$$

$$a=1, r=2$$

$$= kn - \frac{(1)(2^k-1)}{2-1}$$

$$= kn - (2^k-1) = kn - (n-1) \quad (\because n=2^k)$$

$$= 1 + (k-1)n$$

$$\boxed{a_n = 1 + [\log_2 n - 1]n}$$

(\because

$$2^k = n.$$

$$\log 2^k = \log n$$

$$k \cdot \log 2 = \log n.$$

$$k = \frac{\log_e n}{\log_e 2} = \log_2 n)$$

6) Find the recurrence relation and the initial condⁿ for the sequence 0, 2, 6, 12, 20, 30, 42, ... Hence find the general term of the sequence.

Soln:-

Given $a_0 = 0, a_1 = 2, a_2 = 6, a_3 = 12, a_4 = 20, \dots$

Consider $a_1 - a_0 = 2 - 0 = 2 = 2 \times 1$

$$a_2 - a_1 = 6 - 2 = 4 = 2 \times 2$$

$$a_3 - a_2 = 12 - 6 = 6 = 2 \times 3$$

$$a_4 - a_3 = 20 - 12 = 8 = 2 \times 4$$

\vdots

$$a_n - a_{n-1} = 2 \times n \text{ is the R.R with the initial cond}^n$$

$$a_0 = 0.$$

adding all these,

$$a_n - a_0 = (2 \times 1) + (2 \times 2) + (2 \times 3) + (2 \times 4) + \dots + (2 \times (n-1)) + (2 \times n)$$

$$a_n - 0 = 2 [1 + 2 + 3 + \dots + n]$$

$$a_n = 2 \frac{n(n+1)}{2}$$

$$\boxed{a_n = n(n+1)}$$

7) The number of virus affected files in a system is 1000 (to start with) and this increases 250% every 2 hours. Use a recurrence relation to determine the no. of virus affected files in the system after one day.

Soln:- Let $a_0 = 1000$

Let a_n denote the no. of virus affected files after $2n$ hours.

It is given that the no. increases by 250% every 2 hours.

$$\therefore a_1 = a_0 + 250\% \cdot a_0$$

$$a_2 = a_1 + 250\% \cdot a_1$$

\vdots

$$a_n = a_{n-1} + 250\% \cdot a_{n-1}$$

$$\begin{aligned}\therefore a_n &= a_{n-1} \left[1 + 250\% \right] \\ &= a_{n-1} \left[1 + \frac{250}{100} \right] \\ &= a_{n-1} (1 + 2.5)\end{aligned}$$

$$\boxed{a_n = 3.5 a_{n-1}} \quad \forall n \geq 1.$$

This is the recurrence relation for the no. of virus affected files.

\therefore General soln of the recurrence relation is given by

$$a_n = c^n a_0.$$

$$a_n = (3.5)^n \times 1000.$$

This gives the no. of virus affected files after 2n hours.

\therefore No. of virus affected files after 24 hrs (1 day)

(when $n=12$) is

$$a_{12} = (3.5)^{12} \times 1000 = 3379220508.$$

8) A person invests Rs. 10,000 at 10.5% interest (per year) compounded monthly. find and solve the recurrence relation for the value of the investment at the end of n months. what is the value of the investment at the end of the 1 year? How long will it take to double the investment?

Soln:- Let a_0 denote the initial investment.

Let a_1, a_2, \dots, a_n denote the investments after 1, 2, 3, ..., n months respectively.

Given: annual rate of interest = 10.5%.

$$\therefore \text{Monthly rate of interest} = \frac{10.5\%}{12} = 0.875\%.$$

Thus $a_0 = 10000$

$$a_1 = a_0 + (0.875\%) a_0.$$

$$a_2 = a_1 + (0.875\%) a_1$$

\vdots

$$a_n = a_{n-1} + (0.875\%) a_{n-1}$$

$$a_n = a_{n-1} [1 + 0.875\%]$$

$$\Rightarrow a_n = a_{n-1} \left[1 + \frac{0.875}{100} \right]$$

(4)

$$\boxed{a_n = 1.00875 a_{n-1}}, \quad \forall n \geq 1.$$

This is the Recurrence relation at the end of 'n' months.

\therefore General soln of the recurrence relation is given by

$$a_n = c^n a_0.$$

$$a_n = (1.00875)^n \times 10000$$

\therefore Investment at the end of first year is ($n=12$)

$$a_{12} = (1.00875)^{12} \times 10000$$

$$= 11102.03$$

$$\underline{a_{12} \approx 11,102}$$

Next, to find n given that

$$a_n = 2a_0.$$

$$\Rightarrow (1.00875)^n \times 10000 = 2 \times 10000$$

$$\Rightarrow (1.00875)^n = 2$$

$$\Rightarrow n \log_e(1.00875) = \log_e 2$$

$$\Rightarrow n = \frac{\log_e 2}{\log_e(1.00875)} = 79.56.$$

$$\boxed{n \approx 80}.$$

Thus the investment will be doubled in about 80 months
time i.e. 6 years and 8 months.

9) A bank pays a certain % of annual interest on deposits, compounding the interest once in 3 months. If a deposit doubles in 6 years and 6 months, what is the annual % of interest paid by the bank?

Soln: Let the annual rate of interest be $x\%$.

\therefore Quarterly rate of interest is $\left(\frac{x}{4}\right)\%$.

Let a_0 be the initial deposit and a_n be the deposit after at the end of n^{th} quarters.

$$a_1 = a_0 + \left(\frac{x}{4}\right) a_0$$

$$a_2 = a_1 + \left(\frac{x}{4}\right) a_1$$

$$a_n = a_{n-1} + \left(\frac{x}{4}\right) a_{n-1}$$

$$= a_{n-1} \left[1 + \frac{x}{4}\right]$$

$$\boxed{a_n = a_{n-1} \left(1 + \frac{x}{400}\right)} \rightarrow \text{is the recurrence relation.} \quad \text{--- (1)}$$

General solⁿ of (1) is

$$a_n = c^n a_0$$

$$\Rightarrow a_n = \left(1 + \frac{x}{400}\right)^n a_0$$

Given that the deposit doubles in 6 yrs, 6 months (i.e. 78 months) i.e. deposit doubles in 26 quarters ($\because \frac{78}{3} = 26$)

$$\therefore n = 26$$

$$\text{we have } a_n = 2 a_0$$

$$\text{i.e. } a_{26} = 2 a_0$$

$$\Rightarrow \left(1 + \frac{x}{400}\right)^{26} = 2$$

$$\Rightarrow 26 \log_e \left(1 + \frac{x}{400}\right) = \log_e 2$$

$$\Rightarrow \log_e \left(1 + \frac{x}{400}\right) = 0.0266595$$

$$\Rightarrow 1 + \frac{x}{400} = e^{0.0266595} = 1.027$$

$$\Rightarrow \frac{x}{400} = 0.027 \Rightarrow \boxed{x = 10.8}$$

Thus the annual rate of interest paid by the bank is 10.8% (compounding the interest once in 3 months).

10) A bank pays 6% interest compound quarterly. If Laura invests Rs. 100 then how many months must she wait for her money to double?

HINT:- 3 months - 6% interest
1 months - ?
 $1 \times \frac{6\%}{3} = 2\%$

$$\left. \begin{array}{l} a_0 = 100 \\ a_n = a_{n-1} (1 + 2\%) \\ a_n = 1.02 a_{n-1} \end{array} \right\} \begin{array}{l} a_n = 2 a_0 \\ \Rightarrow n = 35 \text{ months} \end{array}$$

Second Order homogenous Recurrence Relation

A second order homogenous recurrence relation is of the form $C_n a_n + C_{n-1} a_{n-1} + C_{n-2} a_{n-2} = 0 \quad \forall n \geq 2 \rightarrow \textcircled{1}$ where C_n, C_{n-1}, C_{n-2} are real constants.

The auxiliary equation of eqⁿ $\textcircled{1}$ is given by

$$C_n k^2 + C_{n-1} k + C_{n-2} = 0$$

Suppose k_1 and k_2 are the roots of A.E

Case (i): If k_1 and k_2 are real and distinct, then general soln of eqⁿ $\textcircled{1}$ is given by

$$a_n = A k_1^n + B k_2^n; \text{ where } A \text{ \& } B \text{ are arbitrary constants}$$

Case (ii): If $k_1 = k_2 = k$, then general soln of eqⁿ $\textcircled{1}$ is given by

$$a_n = (A + Bn) k^n.$$

Case (iii): If k_1 and k_2 are imaginary i.e. if $k_1 = p + iq$, and $k_2 = p - iq$, then the general soln of eqⁿ $\textcircled{1}$ is given by

$$a_n = r^n [A \cos n\theta + B \sin n\theta] \quad \text{where } r = \sqrt{p^2 + q^2} \\ \theta = \tan^{-1}\left(\frac{q}{p}\right).$$

Solve the following Recurrence relations:

$\Rightarrow a_n + a_{n-1} - 6a_{n-2} = 0 \quad \forall n \geq 2$ given $a_0 = -1, a_1 = 8$.
 $\hookrightarrow \textcircled{1}$

Soln:- comparing with $C_n a_n + C_{n-1} a_{n-1} + C_{n-2} a_{n-2} = 0$, we have

$$C_n = 1, C_{n-1} = 1, C_{n-2} = -6.$$

$$\text{A.E is } C_n k^2 + C_{n-1} k + C_{n-2} = 0.$$

$$\Rightarrow k^2 + k - 6 = 0$$

$$\Rightarrow (k+3)(k-2) = 0.$$

$$\Rightarrow \text{Roots are } \boxed{k_1 = -3, k_2 = 2} \quad \text{real \& distinct roots.}$$

\therefore General soln of $\textcircled{1}$ is given by

$$a_n = A \cdot (-3)^n + B \cdot 2^n \rightarrow \textcircled{2}.$$

Given $a_0 = -1, a_1 = 8$.

sub $n=0$ in (2),

$$a_0 = A(-3)^0 + B(2)^0$$

$$-1 = A + B \rightarrow (3)$$

sub $n=1$ in (2)

$$a_1 = A(-3)^1 + B(2)^1$$

$$8 = -3A + 2B \rightarrow (4)$$

Solving (3) & (4), $\boxed{A = -2, B = 1}$

sub in (2), $a_n = -2(-3)^n + 1 \cdot (2)^n$

$$\underline{\underline{a_n = 2^n - 2(-3)^n}}$$

2) $2a_n = 7a_{n-1} - 3a_{n-2}, n \geq 2, a_0 = 2, a_1 = 5$.

soln:- $2a_n - 7a_{n-1} + 3a_{n-2} = 0 \rightarrow (1)$

AE: $2k^2 - 7k + 3 = 0$.

Roots are $k_1 = 3, k_2 = \frac{1}{2}$. (real & distinct)

General soln is given by

$$a_n = A \cdot 3^n + B \cdot \left(\frac{1}{2}\right)^n \rightarrow (2)$$

given $a_0 = 2, a_1 = 5$

sub $n=0$ in (2), $a_0 = A + B$.

$$\Rightarrow 2 = A + B \rightarrow (3)$$

sub $n=1$ in (2), $a_1 = 3A + \frac{1}{2}B$

$$\Rightarrow 5 = \frac{6A + B}{2} \text{ (or) } 6A + B = 10 \rightarrow (4)$$

solving (3) & (4), $\boxed{A = \frac{8}{5}}, \boxed{B = \frac{2}{5}}$.

sub in (2), $a_n = \left(\frac{8}{5}\right)3^n + \left(\frac{2}{5}\right)\left(\frac{1}{2}\right)^n$

3) $a_n - 6a_{n-1} + 9a_{n-2} = 0, n \geq 2, a_0 = 5, a_1 = 12$.
 $\rightarrow (1)$

soln:-

AE: $k^2 - 6k + 9 = 0$.

$$\Rightarrow \boxed{K=3, 3}$$

(5)

Roots are real & repeated.

∴ General soln of ① is

$$a_n = (A+Bn)3^n \rightarrow (2)$$

Given $a_0 = 5, a_1 = 12$

sub $n=0$ in ② $\Rightarrow a_0 = A \cdot 3^0$

$$\boxed{5 = A}$$

sub $n=1$ in ② $\Rightarrow a_1 = (A+B)3^1$

$$12 = (5+B)3$$

$$12 = 15 + 3B$$

$$3B = -3 \Rightarrow \boxed{B=-1}$$

sub in ①, $\boxed{a_n = (5-n)3^n}$

4) $4a_n + 2a_{n+1} + a_{n+2} = 0$

Soln: AE: $4k^2 + 2k + 1 = 0$

$$k = \frac{-2 \pm \sqrt{4-16}}{8} = \frac{-2 \pm \sqrt{-12}}{8} = \frac{-2 \pm 2i\sqrt{3}}{8}$$

$$\boxed{k = \frac{-1 \pm \sqrt{3}i}{4}}$$

Roots are $k_1 = \frac{-1+\sqrt{3}i}{4}, k_2 = \frac{-1-\sqrt{3}i}{4}$ (imaginary roots)

comparing with $p \pm iq$, $p = -\frac{1}{4}, q = \frac{\sqrt{3}}{4}$

$$\therefore r = \sqrt{p^2 + q^2} = \sqrt{\frac{1}{16} + \frac{3}{16}} = \sqrt{\frac{4}{16}} = \sqrt{\frac{1}{4}} = \frac{1}{2}$$

$$\theta = \tan^{-1}\left(\frac{q}{p}\right) = \tan^{-1}\left(\frac{\frac{\sqrt{3}}{4}}{-\frac{1}{4}}\right) = \tan^{-1}(-\sqrt{3}) = -60^\circ = -\frac{\pi}{3}$$

∴ General soln of ① is

$$a_n = r^n [A \cos n\theta + B \sin n\theta]$$

$$a_n = \left(\frac{1}{2}\right)^n \left[A \cos\left(-\frac{n\pi}{3}\right) + B \sin\left(-\frac{n\pi}{3}\right) \right]$$

5) $a_n = 2(a_{n+1} - a_{n+2})$, for $n \geq 2$ given that $a_0 = 1$ & $a_1 = 2$.

Hw

Ans: $a_n = (\sqrt{2})^n \left[\cos \frac{n\pi}{4} + \sin \frac{n\pi}{4} \right]$

6) $D_n = bD_{n-1} - b^2 D_{n-2}$ for $n \geq 3$, given $D_1 = b > 0$, $D_2 = 0$

Soln:- $D_n - bD_{n-1} + b^2 D_{n-2} = 0 \rightarrow (1)$

AE: $k^2 - bk + b^2 = 0$

$$k = \frac{b \pm \sqrt{b^2 - 4b^2}}{2} = \frac{b \pm \sqrt{-3b^2}}{2} = \frac{b \pm i\sqrt{3}b}{2}$$

$\therefore k_1 = \frac{b}{2} + i\frac{\sqrt{3}b}{2}$ and $k_2 = \frac{b}{2} - i\frac{\sqrt{3}b}{2}$ (imaginary roots)

\therefore General soln for D_n is

$$D_n = r^n [A \cos n\theta + B \sin n\theta] \rightarrow (2)$$

where A and B are arbitrary constants.

$$r = \sqrt{p^2 + q^2} = \sqrt{\frac{b^2}{4} + \frac{3b^2}{4}} = \sqrt{\frac{4b^2}{4}} = b.$$

$$p = \frac{b}{2}, q = \frac{\sqrt{3}b}{2}$$

$$\theta = \tan^{-1}\left(\frac{q}{p}\right) = \tan^{-1}\left(\frac{\sqrt{3}b/2}{b/2}\right) = \tan^{-1} \sqrt{3} = \pi/3.$$

(2) $\Rightarrow D_n = b^n [A \cos \frac{n\pi}{3} + B \sin \frac{n\pi}{3}] \rightarrow (3)$ is the g. soln.

given $D_1 = b$, $D_2 = 0$.

Put $n=1$ in (3), $D_1 = b [A \cos \frac{\pi}{3} + B \sin \frac{\pi}{3}]$

$$b = b [A \cdot \frac{1}{2} + B \cdot \frac{\sqrt{3}}{2}]$$

$$\Rightarrow \boxed{1 = \frac{1}{2}A + \frac{\sqrt{3}}{2}B} \rightarrow (4)$$

Put $n=2$ in (3), $D_2 = b^2 [A \cos \frac{2\pi}{3} + B \sin \frac{2\pi}{3}]$

$$0 = b^2 [A \cos (180^\circ - 60^\circ) + B \sin (180^\circ - 60^\circ)]$$

$$\Rightarrow 0 = -A \cos 60^\circ + B \sin 60^\circ$$

$$\Rightarrow \boxed{0 = -\frac{1}{2}A + \frac{\sqrt{3}}{2}B} \rightarrow (5)$$

Solving (4) & (5), $\boxed{A=1, B=1/\sqrt{3}}$

sub in (3),

$$D_n = b^n \left[\cos \frac{n\pi}{3} + \frac{1}{\sqrt{3}} \sin \frac{n\pi}{3} \right].$$

7) $F_{n+2} = f_{n+1} + f_n$ for $n \geq 0$, given $F_0 = 0$, $F_1 = 1$.

Soln:- Rewriting as $f_n = f_{n+1} + f_{n-2}$

$$\Rightarrow f_n - f_{n+1} - f_{n-2} = 0 \quad \text{for } n \geq 2$$

\hookrightarrow (1)

AE: $k^2 - k - 1 = 0$.

$$k = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 \pm \sqrt{5}}{2} \quad (\text{real \& distinct roots})$$

$$k_1 = \frac{1+\sqrt{5}}{2}, \quad k_2 = \frac{1-\sqrt{5}}{2}$$

General solⁿ of (1) is

$$F_n = A \left(\frac{1+\sqrt{5}}{2} \right)^n + B \left(\frac{1-\sqrt{5}}{2} \right)^n, \text{ where } A \text{ \& } B \text{ are arbitrary constants.}$$

\hookrightarrow (2)

Given $F_0 = 0$, $F_1 = 1$.

sub $n=0$ in (2), $f_0 = A + B$

$$\boxed{0 = A + B} \rightarrow (3)$$

sub $n=1$ in (2), $F_1 = A \left(\frac{1+\sqrt{5}}{2} \right) + B \left(\frac{1-\sqrt{5}}{2} \right)$

$$\Rightarrow \boxed{1 = A \left(\frac{1+\sqrt{5}}{2} \right) + B \left(\frac{1-\sqrt{5}}{2} \right)} \rightarrow (4)$$

Eqⁿ (3) $\times \left(\frac{1+\sqrt{5}}{2} \right)$ gives $0 = \left(\frac{1+\sqrt{5}}{2} \right) A + \left(\frac{1+\sqrt{5}}{2} \right) B \rightarrow (5)$

$$(4) - (5) \Rightarrow 1 = B \left(\frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} \right)$$

$$1 = B \left(-\frac{2\sqrt{5}}{2} \right) \Rightarrow \boxed{B = -\frac{1}{\sqrt{5}}}$$

from (3), $A = -B \Rightarrow \boxed{A = \frac{1}{\sqrt{5}}}$

sub in (2),

$$F_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right]$$