

# Algebraic Geometry 2

## Tutorial session 5

Lecturer: Rami Aizenbud  
TA: Shai Shechter

May 23, 2020

## Proposition (2.3 in Hartshorne)

- ① *Let  $\varphi : A \rightarrow B$  be a homomorphism of rings. Then  $\varphi$  induces a natural morphism of locally ringed spaces*

$$(f, f^\#) : (\mathrm{Spec}(B), \mathcal{O}_{\mathrm{Spec}(B)}) \rightarrow (\mathrm{Spec}(A), \mathcal{O}_{\mathrm{Spec}(A)}).$$

- ② *Conversely, any morphism of locally ring spaces  $(f, f^\#)$  as above is induced from a ring homomorphism  $\varphi : A \rightarrow B$ .*

## Example

Let  $R = \overline{\mathbb{F}_p}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ , for  $q = p^\alpha$ .

## Example

Let  $R = \overline{\mathbb{F}_p}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ , for  $q = p^\alpha$ .  $T = \operatorname{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi$ .

## Example

Let  $R = \overline{\mathbb{F}_p}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ , for  $q = p^\alpha$ .  $T = \operatorname{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi$ .

- The associated map  $f = \varphi^{-1} : \operatorname{Spec}(R) \rightarrow \operatorname{Spec}(R)$  is the identity map.

## Example

Let  $R = \overline{\mathbb{F}_p}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ , for  $q = p^\alpha$ .  $T = \operatorname{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi$ .

- The associated map  $f = \varphi^{-1} : \operatorname{Spec}(R) \rightarrow \operatorname{Spec}(R)$  is the identity map.
- Given  $V \subseteq \operatorname{Spec}(R)$  and  $s \in \mathcal{O}_{\operatorname{Spec}(R)}(V)$ , we have

$$f^\#(s)(\cdot) = (s(\cdot))^q.$$

## Corollary (of the proposition)

*Let  $A, B$  be rings. Then  $A$  and  $B$  are isomorphic if and only if  $\operatorname{Spec}(A)$  and  $\operatorname{Spec}(B)$  are isomorphic as locally ringed spaces.*

## Corollary (of the proposition)

*Let  $A, B$  be rings. Then  $A$  and  $B$  are isomorphic if and only if  $\operatorname{Spec}(A)$  and  $\operatorname{Spec}(B)$  are isomorphic as locally ringed spaces.*

More precisely, the proposition says that

$$A \mapsto \operatorname{Spec}(A)$$

is a fully faithful functor from the category of commutative unital rings to the category of schemes; the image of this functor is the sub-category of *affine schemes*.



## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

A *scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  in which  $X$  admits an open cover  $X = \bigcup U_\alpha$  such that  $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$  is an affine scheme for all  $\alpha$ .

## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

A *scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  in which  $X$  admits an open cover  $X = \bigcup U_\alpha$  such that  $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$  is an affine scheme for all  $\alpha$ .

Here  $\mathcal{O}_X|_{U_\alpha}$  denotes the restricted sheaf  $\mathcal{O}_X|_{U_\alpha}(V) = \mathcal{O}_X(V)$  for  $V \subseteq U_\alpha$  open.

## Example

Let  $K$  be an infinite field and consider

$$X = ((\{0\} \times \mathbb{A}_K) \cup (\mathbb{A}_K \times \{1\})) / \sim,$$

where  $\sim$  is the smallest equivalence relation such that  $(0, x) \sim (x, 1)$  for all  $0 \neq x \in K$ .

## Example

Let  $K$  be an infinite field and consider

$$X = ((\{0\} \times \mathbb{A}_K) \cup (\mathbb{A}_K \times \{1\})) / \sim,$$

where  $\sim$  is the smallest equivalence relation such that  $(0, x) \sim (x, 1)$  for all  $0 \neq x \in K$ . Endow  $X$  with the quotient of the Zariski topology. Then  $X$  is homeomorphic to an affine line with two origins:

\_\_\_\_\_ : \_\_\_\_\_

## Example

Let  $K$  be an infinite field and consider

$$X = ((\{0\} \times \mathbb{A}_K) \cup (\mathbb{A}_K \times \{1\})) / \sim,$$

where  $\sim$  is the smallest equivalence relation such that  $(0, x) \sim (x, 1)$  for all  $0 \neq x \in K$ . Endow  $X$  with the quotient of the Zariski topology. Then  $X$  is homeomorphic to an affine line with two origins:

\_\_\_\_\_ : \_\_\_\_\_

$X$  is also endowed with a structure sheaf  $\mathcal{O}_X$ :

## Example

Let  $K$  be an infinite field and consider

$$X = ((\{0\} \times \mathbb{A}_K) \cup (\mathbb{A}_K \times \{1\})) / \sim,$$

where  $\sim$  is the smallest equivalence relation such that  $(0, x) \sim (x, 1)$  for all  $0 \neq x \in K$ . Endow  $X$  with the quotient of the Zariski topology. Then  $X$  is homeomorphic to an affine line with two origins:

$$\text{—————} : \text{—————}$$

$X$  is also endowed with a structure sheaf  $\mathcal{O}_X$ :

Let  $X_1, X_2$  be the images of  $\{0\} \times \mathbb{A}_K$  and  $\mathbb{A}_K \times \{1\}$  in  $X$ , respectively. Given  $U \subseteq X$  open, write  $U_i = U \cap X_i$  and define

$$\mathcal{O}_X(U) = \{(s_1, s_2) \in \mathcal{O}_{X_1}(U_1) \times \mathcal{O}_X(U_2) : s_1|_{U_1 \cap U_2} = s_2|_{U_1 \cap U_2}\}.$$

The locally ringed space  $(X, \mathcal{O}_X)$  from the previous example is a scheme, which is not affine.



The locally ringed space  $(X, \mathcal{O}_X)$  from the previous example is a scheme, which is not affine.

## Question

How do we know that it's not affine?

The locally ringed space  $(X, \mathcal{O}_X)$  from the previous example is a scheme, which is not affine.

## Question

How do we know that it's not affine?

## Answer

Compute it's global sections.

The locally ringed space  $(X, \mathcal{O}_X)$  from the previous example is a scheme, which is not affine.

## Question

How do we know that it's not affine?

## Answer

Compute it's global sections. For  $A = \Gamma(X, \mathcal{O}_X)$  we have that  $X$  is affine iff  $X$  is isomorphic, as a locally ringed spacem to  $\text{Spec}(A)$ .

The locally ringed space  $(X, \mathcal{O}_X)$  from the previous example is a scheme, which is not affine.

## Question

How do we know that it's not affine?

## Answer

Compute it's global sections. For  $A = \Gamma(X, \mathcal{O}_X)$  we have that  $X$  is affine iff  $X$  is isomorphic, as a locally ringed spacem to  $\text{Spec}(A)$ . But, by definition:

$$\mathcal{O}_X(X) := \{(s_1, s_2) \in \mathcal{O}_{X_1}(X_1) \times \mathcal{O}_{X_2}(X_2) : s_1|_{X_1} = s_2|_{X_2}\},$$

i.e. is the set of polynomial functions on  $X_i = \mathbb{A}_K$  which agree everywhere except zero. Since  $K$  is infinite, this can only occur if  $s_1 = s_2$  as polynomials.

# Examples of morphisms of affine schemes

Let  $A$  be a ring.

# Examples of morphisms of affine schemes

Let  $A$  be a ring.

## Example (Quotient by ideal)

Let  $I \triangleleft A$  be an ideal and  $\varphi : A \rightarrow A/I$  the quotient map.

# Examples of morphisms of affine schemes

Let  $A$  be a ring.

## Example (Quotient by ideal)

Let  $I \triangleleft A$  be an ideal and  $\varphi : A \rightarrow A/I$  the quotient map. The associated map  $f = \varphi^{-1} : \operatorname{Spec}(A/I) \rightarrow \operatorname{Spec}(A)$  is a closed injective map whose image is  $V(I)$ .

# Examples of morphisms of affine schemes

Let  $A$  be a ring.

## Example (Quotient by ideal)

Let  $I \triangleleft A$  be an ideal and  $\varphi : A \rightarrow A/I$  the quotient map. The associated map  $f = \varphi^{-1} : \operatorname{Spec}(A/I) \rightarrow \operatorname{Spec}(A)$  is a closed injective map whose image is  $V(I)$ . Indeed,  $f(V(J)) = V(\varphi^{-1}(J))$  for any  $J \triangleleft A/I$ .

For example, for  $A = \mathbb{C}[x, y]/(xy)$ ,  $\operatorname{Spec}(A)$  is naturally identified with a subset of  $\mathbb{A}_{\mathbb{C}}^2$  (the cross  $xy = 0$ ), with its minimal primes  $(x)$  and  $(y)$  corresponding the (generic points of the) irreducible components of  $\operatorname{Spec}(A)$ . All other points are closed and of the form  $(x - a, y)$  or  $(x, y - b)$ .



# Examples of morphisms of affine schemes

## Example (Localization)

Let  $S \subseteq A$  be a multiplicatively closed set, and  $\varphi : A \rightarrow S^{-1}A$  the localization map (i.e.  $\varphi(a) = a/1$ ).

# Examples of morphisms of affine schemes

## Example (Localization)

Let  $S \subseteq A$  be a multiplicatively closed set, and  $\varphi : A \rightarrow S^{-1}A$  the localization map (i.e.  $\varphi(a) = a/1$ ).

## Fact

The map  $\mathfrak{q} \mapsto \varphi^{-1}(\mathfrak{q})$  is an order-preserving (wrt inclusion) bijection of the primes of  $S^{-1}A$  onto the primes  $\mathfrak{q} \triangleleft A$  such that  $\mathfrak{q} \cap S = \emptyset$ .

# Examples of morphisms of affine schemes

- If  $S = A \setminus \mathfrak{p}$  for  $\mathfrak{p} \in \operatorname{Spec}(A)$ , then  $f : \operatorname{Spec}(A_{\mathfrak{p}}) \rightarrow \operatorname{Spec}(A)$  is the inclusion of  $\overline{\mathfrak{p}} = V(\mathfrak{p})$  in  $\operatorname{Spec}(A)$ .

# Examples of morphisms of affine schemes

- If  $S = A \setminus \mathfrak{p}$  for  $\mathfrak{p} \in \operatorname{Spec}(A)$ , then  $f : \operatorname{Spec}(A_{\mathfrak{p}}) \rightarrow \operatorname{Spec}(A)$  is the inclusion of  $\overline{\mathfrak{p}} = V(\mathfrak{p})$  in  $\operatorname{Spec}(A)$ .

For example, take  $A = \mathbb{C}[x, y]$  and  $\mathfrak{p} = (x)$ . Then

$$S^{-1}A = \mathbb{C}[x, y]_{(x)} = \{f(x, y)/g(x, y) \mid g(0, y) \neq 0\}$$

and  $f$  is given by  $f(\mathfrak{q}) = \mathfrak{q} \cap A$ , for  $\mathfrak{q} \in \operatorname{Spec}(S^{-1}A)$ . It maps  $\operatorname{Spec}(S^{-1}A)$  homeomorphically onto the affine line  $V((x)) \subseteq \mathbb{A}^2$ ,

# Examples of morphisms of affine schemes

- If  $S = \{1, a, a^2, a^3, \dots\}$  for  $a \in A$ , then  $f : \operatorname{Spec}(A_p) \rightarrow \operatorname{Spec}(A)$  is the inclusion of  $D(a)$  in  $\operatorname{Spec}(A)$ .

# Examples of morphisms of affine schemes

- If  $S = \{1, a, a^2, a^3, \dots\}$  for  $a \in A$ , then  $f : \operatorname{Spec}(A_p) \rightarrow \operatorname{Spec}(A)$  is the inclusion of  $D(a)$  in  $\operatorname{Spec}(A)$ .

For example, for  $A = \mathbb{C}[x]$  and  $a = (x)$ , we get  $S^{-1}A = A_x = \mathbb{C}[x, x^{-1}]$  and  $f$  is the open inclusion of  $D(x) = \mathbb{A}^1 \setminus V(x)$  (line without the origin) into  $\mathbb{A}^1$ .

From the previous two examples we now have a method for “separating” the points of an affine scheme  $X = \operatorname{Spec}(A)$ .

From the previous two examples we now have a method for “separating” the points of an affine scheme  $X = \operatorname{Spec}(A)$ . Specifically, given  $x \in X$ , let  $\mathfrak{m}_x \triangleleft \mathcal{O}_{X,x}$  be the maximal ideal within the stalk at  $x$  and put  $k(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$ .

We have an inclusion

$$\operatorname{Spec}(k(x)) \xrightarrow{\text{closed embedding}} \operatorname{Spec}(\mathcal{O}_{X,x}) \xrightarrow{\text{homeo onto } V(x)} \operatorname{Spec}(A)$$

of *locally ringed spaces* whose image is precisely the point  $x$ .



From the previous two examples we now have a method for “separating” the points of an affine scheme  $X = \operatorname{Spec}(A)$ . Specifically, given  $x \in X$ , let  $\mathfrak{m}_x \triangleleft \mathcal{O}_{X,x}$  be the maximal ideal within the stalk at  $x$  and put  $k(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$ .

We have an inclusion

$$\operatorname{Spec}(k(x)) \xrightarrow{\text{closed embedding}} \operatorname{Spec}(\mathcal{O}_{X,x}) \xrightarrow{\text{homeo onto } V(x)} \operatorname{Spec}(A)$$

of *locally ringed spaces* whose image is precisely the point  $x$ .

That is-

## Corollary

*For any scheme  $X$  and  $x \in X$ , there exists a field  $k = k(x)$  and a morphism  $\operatorname{Spec}(k) \rightarrow X$  whose image is precisely  $x$ .*

## Exercise (Hartshorne, Ex 2.7)

Let  $X$  be a scheme and  $K$  a field. Show that to give a morphism  $\mathrm{Spec}(K) \rightarrow X$  is equivalent to specifying a point  $x \in X$  and an inclusion of fields  $k(x) \rightarrow K$ .

## Exercise (Hartshorne, Ex 2.7)

Let  $X$  be a scheme and  $K$  a field. Show that to give a morphism  $\mathrm{Spec}(K) \rightarrow X$  is equivalent to specifying a point  $x \in X$  and an inclusion of fields  $k(x) \rightarrow K$ .

## Proof.

Assume given a morphism  $f : \mathrm{Spec}(K) \rightarrow X$  and let  $x$  be the image of the unique point  $* \in \mathrm{Spec}(K)$ .

## Exercise (Hartshorne, Ex 2.7)

Let  $X$  be a scheme and  $K$  a field. Show that to give a morphism  $\mathrm{Spec}(K) \rightarrow X$  is equivalent to specifying a point  $x \in X$  and an inclusion of fields  $k(x) \rightarrow K$ .

## Proof.

Assume given a morphism  $f : \mathrm{Spec}(K) \rightarrow X$  and let  $x$  be the image of the unique point  $* \in \mathrm{Spec}(K)$ . The associated sheaf morphism  $f^\#$  induces a *local* homomorphism

$$f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\mathrm{Spec}(K),*} \simeq K,$$

i.e.  $\mathfrak{m}_x = \mathrm{Ker}(f)$ .

## Exercise (Hartshorne, Ex 2.7)

Let  $X$  be a scheme and  $K$  a field. Show that to give a morphism  $\mathrm{Spec}(K) \rightarrow X$  is equivalent to specifying a point  $x \in X$  and an inclusion of fields  $k(x) \rightarrow K$ .

## Proof.

Assume given a morphism  $f : \mathrm{Spec}(K) \rightarrow X$  and let  $x$  be the image of the unique point  $* \in \mathrm{Spec}(K)$ . The associated sheaf morphism  $f^\#$  induces a *local* homomorphism

$$f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\mathrm{Spec}(K),*} \simeq K,$$

i.e.  $\mathfrak{m}_x = \mathrm{Ker}(f)$ . Thus  $f^\#$  reduces to an inclusion  $k(x) \hookrightarrow K$ .

## Exercise (Hartshorne, Ex 2.7)

Let  $X$  be a scheme and  $K$  a field. Show that to give a morphism  $\mathrm{Spec}(K) \rightarrow X$  is equivalent to specifying a point  $x \in X$  and an inclusion of fields  $k(x) \rightarrow K$ .

### Proof.

Assume given a morphism  $f : \mathrm{Spec}(K) \rightarrow X$  and let  $x$  be the image of the unique point  $* \in \mathrm{Spec}(K)$ . The associated sheaf morphism  $f^\#$  induces a *local* homomorphism

$$f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\mathrm{Spec}(K),*} \simeq K,$$

i.e.  $\mathfrak{m}_x = \mathrm{Ker}(f)$ . Thus  $f^\#$  reduces to an inclusion  $k(x) \hookrightarrow K$ . Conversely, given a map  $k(x) \rightarrow K$ , we get a morphism

$\mathrm{Spec}(K) \rightarrow \mathrm{Spec}(k(x))$ , which we can then compose with the map  $\mathrm{Spec}(k(x)) \rightarrow X$



## The functor of points approach

*Tell me who your friends are and I'll tell you who you are*

# The Yoneda Lemma

Let  $\underline{\mathbf{C}}$  be a category.



# The Yoneda Lemma

Let  $\underline{\mathbf{C}}$  be a category. Given  $A \in \underline{\mathbf{C}}$  we have a covariant functor  $h^A : \underline{\mathbf{C}} \rightarrow \mathbf{Set}$  defined by

$$h^A(B) = \text{Mor}_{\underline{\mathbf{C}}}(B, A),$$

acting on a morphism  $f : B \rightarrow C$  by

$$h^A(f) = (\psi \mapsto \psi \circ f) : \text{Mor}_{\underline{\mathbf{C}}}(C, A) \rightarrow \text{Mor}_{\underline{\mathbf{C}}}(B, A).$$

# The Yoneda Lemma

Let  $\underline{\mathbf{C}}$  be a category. Given  $A \in \underline{\mathbf{C}}$  we have a covariant functor  $h^A : \underline{\mathbf{C}} \rightarrow \underline{\mathbf{Set}}$  defined by

$$h^A(B) = \text{Mor}_{\underline{\mathbf{C}}}(B, A),$$

acting on a morphism  $f : B \rightarrow C$  by

$$h^A(f) = (\psi \mapsto \psi \circ f) : \text{Mor}_{\underline{\mathbf{C}}}(C, A) \rightarrow \text{Mor}_{\underline{\mathbf{C}}}(B, A).$$

## Definition

The **Yoneda functor**  $I : \underline{\mathbf{C}} \rightarrow \text{Mor}_{\underline{\mathbf{Cat}}}(\underline{\mathbf{C}}^{\text{op}}, \underline{\mathbf{Set}})$  is defined by

$$I(A) = h^A \quad \text{and} \quad I(\psi)(\varphi) = \psi \circ \varphi$$

for  $A, A', B \in \underline{\mathbf{C}}$ ,  $\psi : A \rightarrow A'$  and  $\varphi \in \text{Mor}_{\underline{\mathbf{C}}}(A, B)$ .

# Theorem (The Yoneda Lemma)

*The functor  $I$  is fully faithful.*

## Theorem (The Yoneda Lemma)

*The functor  $I$  is fully faithful. That is, given  $A, A' \in \underline{\mathbf{C}}$ , any natural transformation  $\varphi : h^A \rightarrow h^{A'}$  is of the form  $I(\psi)$  for a **unique** morphism  $\psi : A \rightarrow A'$ .*

## Theorem (The Yoneda Lemma)

*The functor  $I$  is fully faithful. That is, given  $A, A' \in \underline{\mathbf{C}}$ , any natural transformation  $\varphi : h^A \rightarrow h^{A'}$  is of the form  $I(\psi)$  for a **unique** morphism  $\psi : A \rightarrow A'$ .*

### Proof of fullness.

Let  $\varphi : h^A \rightarrow h^{A'}$ ,  $C \in \underline{\mathbf{C}}$  and  $\tau \in \text{Mor}_{\underline{\mathbf{C}}}(C, A)$  be given. We seek  $\psi \in \text{Mor}_{\underline{\mathbf{C}}}(A, A')$  such that  $\varphi_C(\tau) = I(\psi)(\tau) = \psi \circ \tau$ .

## Theorem (The Yoneda Lemma)

*The functor  $I$  is fully faithful. That is, given  $A, A' \in \underline{\mathbf{C}}$ , any natural transformation  $\varphi : h^A \rightarrow h^{A'}$  is of the form  $I(\psi)$  for a **unique** morphism  $\psi : A \rightarrow A'$ .*

### Proof of fullness.

Let  $\varphi : h^A \rightarrow h^{A'}$ ,  $C \in \underline{\mathbf{C}}$  and  $\tau \in \text{Mor}_{\underline{\mathbf{C}}}(C, A)$  be given. We seek  $\psi \in \text{Mor}_{\underline{\mathbf{C}}}(A, A')$  such that  $\varphi_C(\tau) = I(\psi)(\tau) = \psi \circ \tau$ . Follow  $\mathbf{1}_A$ :

$$\begin{array}{ccc} h^A(A) & \xrightarrow{h^A(\tau)} & h^A(C) \\ \varphi_A \downarrow & & \downarrow \varphi_C \\ h^{A'}(A) & \xrightarrow{h^{A'}(\tau)} & h^{A'}(C). \end{array}$$

## Theorem (The Yoneda Lemma)

*The functor  $I$  is fully faithful. That is, given  $A, A' \in \underline{\mathbf{C}}$ , any natural transformation  $\varphi : h^A \rightarrow h^{A'}$  is of the form  $I(\psi)$  for a **unique** morphism  $\psi : A \rightarrow A'$ .*

### Proof of fullness.

Let  $\varphi : h^A \rightarrow h^{A'}$ ,  $C \in \underline{\mathbf{C}}$  and  $\tau \in \text{Mor}_{\underline{\mathbf{C}}}(C, A)$  be given. We seek  $\psi \in \text{Mor}_{\underline{\mathbf{C}}}(A, A')$  such that  $\varphi_C(\tau) = I(\psi)(\tau) = \psi \circ \tau$ . Follow  $\mathbf{1}_A$ :

$$\begin{array}{ccc} h^A(A) & \xrightarrow{h^A(\tau)} & h^A(C) \\ \varphi_A \downarrow & & \downarrow \varphi_C \\ h^{A'}(A) & \xrightarrow{h^{A'}(\tau)} & h^{A'}(C). \end{array}$$

Specifically:

$$\varphi_C(\tau) = \varphi_C \circ h^A(\tau)(\mathbf{1}_A) = h^{A'}(\tau) \circ \varphi_A(\mathbf{1}_A) = \varphi(\mathbf{1}_A) \circ \tau.$$

Faithfulness will be proved in the home exercise.



Faithfulness will be proved in the home exercise.

In fact, the Yoneda lemma is even more general (see Wikipedia), but we will not require this generality at the moment.

## Definition

A functor  $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \underline{\mathbf{Sets}}$  is said to be *representable* if there exists  $A \in \underline{\mathbf{C}}$  such that  $F = h^A$ ; i.e.  $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$  for all  $B \in \underline{\mathbf{C}}$ .

## Definition

A functor  $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \underline{\mathbf{Sets}}$  is said to be *representable* if there exists  $A \in \underline{\mathbf{C}}$  such that  $F = h^A$ ; i.e.  $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$  for all  $B \in \underline{\mathbf{C}}$ .

There is also an analogous definition for covariant functors.

## Definition

A functor  $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \mathbf{Sets}$  is said to be *representable* if there exists  $A \in \underline{\mathbf{C}}$  such that  $F = h^A$ ; i.e.  $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$  for all  $B \in \underline{\mathbf{C}}$ .

There is also an analogous definition for covariant functors. There  $F : \underline{\mathbf{C}} \rightarrow \mathbf{Set}$  is said to be representable if

$$F = h_A = (B \mapsto \text{Mor}_{\underline{\mathbf{C}}}(A, B)) \text{ for some } A \in \underline{\mathbf{C}}.$$

# New definition of (affine) schemes

Given a scheme  $(X, \mathcal{O}_X)$ , we can define a functor  $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$  by

$$F_X(R) = \mathrm{Hom}_{\underline{\mathbf{Sch}}}(\mathrm{Spec}(R), X).$$

# New definition of (affine) schemes

Given a scheme  $(X, \mathcal{O}_X)$ , we can define a functor  $\mathbf{Ring} \rightarrow \mathbf{Set}$  by

$$F_X(R) = \mathrm{Hom}_{\mathbf{Sch}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of  $X$ . By an exercise, a scheme is affine iff  $F_X$  is representable; specifically,  $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$ .

# New definition of (affine) schemes

Given a scheme  $(X, \mathcal{O}_X)$ , we can define a functor  $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$  by

$$F_X(R) = \mathrm{Hom}_{\underline{\mathbf{Sch}}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of  $X$ . By an exercise, a scheme is affine iff  $F_X$  is representable; specifically,  $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$ .

## Definition (New definition of affine scheme)

An affine scheme is a representable functor  $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$ .

# New definition of (affine) schemes

Given a scheme  $(X, \mathcal{O}_X)$ , we can define a functor  $\mathbf{Ring} \rightarrow \mathbf{Set}$  by

$$F_X(R) = \mathrm{Hom}_{\mathbf{Sch}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of  $X$ . By an exercise, a scheme is affine iff  $F_X$  is representable; specifically,  $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$ .

## Definition (New definition of affine scheme)

An affine scheme is a representable functor  $\mathbf{Ring} \rightarrow \mathbf{Set}$ .

A general scheme, in this setting, would be a functor which is “locally representable”.



## Example

Consider the ring  $A = \mathbb{Z}[t, t^{-1}]$ .

## Example

Consider the ring  $A = \mathbb{Z}[t, t^{-1}]$ . Then  $\operatorname{Spec}(A) = \operatorname{Spec}(\mathbb{Z}[t] \setminus V(t))$ , which is, a-priori, not easy to describe.

## Example

Consider the ring  $A = \mathbb{Z}[t, t^{-1}]$ . Then  $\text{Spec}(A) = \text{Spec}(\mathbb{Z}[t] \setminus V(t))$ , which is, a-priori, not easy to describe.

The associated functor of points, however, is very easy. Namely-

$$F_{\text{Spec}(A)}(R) = \text{Hom}_{\underline{\text{Rings}}}(\mathbb{Z}[t, t^{-1}], R) = R^{\times},$$

since choosing a homomorphism  $\mathbb{Z}[t, t^{-1}] \rightarrow R$  amounts to choosing the image of  $t$ , which is necessarily in  $R^{\times}$ .

## Example

Consider the ring  $A = \mathbb{Z}[t, t^{-1}]$ . Then  $\mathrm{Spec}(A) = \mathrm{Spec}(\mathbb{Z}[t] \setminus V(t))$ , which is, a-priori, not easy to describe.

The associated functor of points, however, is very easy. Namely-


$$F_{\mathrm{Spec}(A)}(R) = \mathrm{Hom}_{\underline{\mathbf{Rings}}}(\mathbb{Z}[t, t^{-1}], R) = R^{\times},$$

since choosing a homomorphism  $\mathbb{Z}[t, t^{-1}] \rightarrow R$  amounts to choosing the image of  $t$ , which is necessarily in  $R^{\times}$ .

In fact, we can also encode the group laws of  $R^{\times}$  in terms natural transformations of  $F_{\mathrm{Spec}(A)}$ , giving rise to  $F_{\mathrm{Spec}(A)}$  as a *group scheme*. For example, the inversion in  $R^{\times}$  is “encoded” in the map

$$t \mapsto t^{-1} : A \rightarrow A.$$

Another very nice example is Mumford's doodle of  $\mathrm{Spec}(\mathbb{Z}[t])$ :



See <http://www.neverendingbooks.org/grothendiecks-functor-of-points> for more information.

In a sense, the functor of points approach and the spectrum manifest two possible interpretations of a phenomenon from classical algebraic topology. Given an affine variety  $V \subseteq \mathbb{C}^n$ , it is given as the solution set of a set of polynomials. However this set is not uniquely defined. What *is* uniquely defined is the (radical of) the ideal generated by these polynomials, or, equivalently, the associated coordinate algebra. If we start with the coordinate algebra  $A = \mathbb{C}[V]$ , instead of the variety, there are ways to reconstruct the variety:

- 1 Consider all maximal ideals of  $A$ ; or
- 2 Consider all homomorphisms  $A \rightarrow \mathbb{C}$ .

Over  $\mathbb{C}$ , Nullstellensatz tells us that these two give the “same” answer and are sufficient to describe  $V$ .

If we want to consider non-ac fields, or rings, these two methods diverge. The first generalizes to the spectrum, and the second to the functor of points.

# Questions?