

Approximating the Representation Zeta Function of Finite Groups of Lie-Type

Advanced School on Representations of Pro- p Groups
ICMAT, Madrid

Shai Shechter

Weizmann Institute of Science

July 9, 2019

Local set-up

- Let $\mathfrak{o} \supseteq \mathbb{Z}_p$ be a complete dvr with maximal ideal \mathfrak{p} and fraction field F .
- Let $\underline{\mathbf{G}} \subseteq \mathrm{GL}_N(F)$ be a semisimple \mathfrak{o} -defined algebraic group with Lie algebra $\underline{\mathfrak{g}}$ (e.g. $\underline{\mathbf{G}} = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\underline{\mathfrak{g}} = \mathfrak{sl}_n(\mathbb{Q}_p)$).
- Put $G = \underline{\mathbf{G}}(\mathfrak{o}) = \underline{\mathbf{G}} \cap \mathrm{GL}_N(\mathfrak{o})$ and $G^k = \mathrm{Ker}(G \rightarrow \underline{\mathbf{G}}(\mathfrak{o}/\mathfrak{p}^k)) = G \cap 1 + \mathfrak{p}^k M_N(\mathfrak{o})$.
Analogously, write $\mathfrak{g} = \underline{\mathfrak{g}}(\mathfrak{o}) = \underline{\mathfrak{g}} \cap M_N(\mathfrak{o})$ and $\mathfrak{g}^k = \mathfrak{p}^k \mathfrak{g}$.

Local set-up

- Let $\mathfrak{o} \supseteq \mathbb{Z}_p$ be a complete dvr with maximal ideal \mathfrak{p} and fraction field F .
- Let $\underline{\mathbf{G}} \subseteq \mathrm{GL}_N(F)$ be a semisimple \mathfrak{o} -defined algebraic group with Lie algebra $\underline{\mathfrak{g}}$ (e.g. $\underline{\mathbf{G}} = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\underline{\mathfrak{g}} = \mathfrak{sl}_n(\mathbb{Q}_p)$).
- Put $G = \underline{\mathbf{G}}(\mathfrak{o}) = \underline{\mathbf{G}} \cap \mathrm{GL}_N(\mathfrak{o})$ and $G^k = \mathrm{Ker}(G \rightarrow \underline{\mathbf{G}}(\mathfrak{o}/\mathfrak{p}^k)) = G \cap 1 + \mathfrak{p}^k M_N(\mathfrak{o})$.
Analogously, write $\mathfrak{g} = \underline{\mathfrak{g}}(\mathfrak{o}) = \underline{\mathfrak{g}} \cap M_N(\mathfrak{o})$ and $\mathfrak{g}^k = \mathfrak{p}^k \mathfrak{g}$.

Local set-up

- Let $\mathfrak{o} \supseteq \mathbb{Z}_p$ be a complete dvr with maximal ideal \mathfrak{p} and fraction field F .
- Let $\underline{\mathbf{G}} \subseteq \mathrm{GL}_N(F)$ be a semisimple \mathfrak{o} -defined algebraic group with Lie algebra $\underline{\mathfrak{g}}$ (e.g. $\underline{\mathbf{G}} = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\underline{\mathfrak{g}} = \mathfrak{sl}_n(\mathbb{Q}_p)$).
- Put $G = \underline{\mathbf{G}}(\mathfrak{o}) = \underline{\mathbf{G}} \cap \mathrm{GL}_N(\mathfrak{o})$ and $G^k = \mathrm{Ker}(G \rightarrow \underline{\mathbf{G}}(\mathfrak{o}/\mathfrak{p}^k)) = G \cap 1 + \mathfrak{p}^k M_n(\mathfrak{o})$.

Analogously, write $\mathfrak{g} = \underline{\mathfrak{g}}(\mathfrak{o}) = \underline{\mathfrak{g}} \cap M_N(\mathfrak{o})$ and $\mathfrak{g}^k = \mathfrak{p}^k \mathfrak{g}$.

Assumption

We assume throughout that $\mathrm{char}(\mathfrak{o}/\mathfrak{p})$ is large enough so G^1 is amenable to the Kirillov Orbit Method. In particular, the series $\exp(x) = 1 + x + \frac{x^2}{2} + \cdots$ defines a bijection $\mathfrak{g}^1 \xrightarrow{1-1} G^1$.

Local set-up

- Let $\mathfrak{o} \supseteq \mathbb{Z}_p$ be a complete dvr with maximal ideal \mathfrak{p} and fraction field F .
- Let $\underline{\mathbf{G}} \subseteq \mathrm{GL}_N(F)$ be a semisimple \mathfrak{o} -defined algebraic group with Lie algebra $\underline{\mathfrak{g}}$ (e.g. $\underline{\mathbf{G}} = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\underline{\mathfrak{g}} = \mathfrak{sl}_n(\mathbb{Q}_p)$).
- Put $G = \underline{\mathbf{G}}(\mathfrak{o}) = \underline{\mathbf{G}} \cap \mathrm{GL}_N(\mathfrak{o})$ and $G^k = \mathrm{Ker}(G \rightarrow \underline{\mathbf{G}}(\mathfrak{o}/\mathfrak{p}^k)) = G \cap 1 + \mathfrak{p}^k M_N(\mathfrak{o})$. Analogously, write $\mathfrak{g} = \underline{\mathfrak{g}}(\mathfrak{o}) = \underline{\mathfrak{g}} \cap M_N(\mathfrak{o})$ and $\mathfrak{g}^k = \mathfrak{p}^k \mathfrak{g}$.

Assumption

We assume throughout that $\mathrm{char}(\mathfrak{o}/\mathfrak{p})$ is large enough so G^1 is amenable to the Kirillov Orbit Method. In particular, the series $\exp(x) = 1 + x + \frac{x^2}{2} + \cdots$ defines a bijection $\mathfrak{g}^1 \xrightarrow{1-1} G^1$.

Local set-up

- Let $\mathfrak{o} \supseteq \mathbb{Z}_p$ be a complete dvr with maximal ideal \mathfrak{p} and fraction field F .
- Let $\underline{\mathbf{G}} \subseteq \mathrm{GL}_N(F)$ be a semisimple \mathfrak{o} -defined algebraic group with Lie algebra $\underline{\mathfrak{g}}$ (e.g. $\underline{\mathbf{G}} = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\underline{\mathfrak{g}} = \mathfrak{sl}_n(\mathbb{Q}_p)$).
- Put $G = \underline{\mathbf{G}}(\mathfrak{o}) = \underline{\mathbf{G}} \cap \mathrm{GL}_N(\mathfrak{o})$ and $G^k = \mathrm{Ker}(G \rightarrow \underline{\mathbf{G}}(\mathfrak{o}/\mathfrak{p}^k)) = G \cap 1 + \mathfrak{p}^k M_N(\mathfrak{o})$.
Analogously, write $\mathfrak{g} = \underline{\mathfrak{g}}(\mathfrak{o}) = \underline{\mathfrak{g}} \cap M_N(\mathfrak{o})$ and $\mathfrak{g}^k = \mathfrak{p}^k \mathfrak{g}$.

Assumption

We assume throughout that $\mathrm{char}(\mathfrak{o}/\mathfrak{p})$ is large enough so G^1 is amenable to the Kirillov Orbit Method. In particular, the series $\exp(x) = 1 + x + \frac{x^2}{2} + \cdots$ defines a bijection $\mathfrak{g}^1 \xrightarrow{1-1} G^1$.

Motivation

“The anomaly of the zeroth principal congruence subgroup”

Observation (cf [AKOV 13, Proposition 3.7])

Write $q = |\mathfrak{o}/\mathfrak{p}|$ and $d = \dim \underline{\mathbf{G}}$. The sequence $(q^{-k \cdot d} \zeta_{G^k})_{k \geq 1}$ is a constant sequence of functions, i.e.

$$q^{-kd} \zeta_{G^k} = q^{-md} \zeta_{G^m} \quad \text{for all } k, m \geq 1.$$

Motivation

“The anomaly of the zeroth principal congruence subgroup”

Observation (cf [AKOV 13, Proposition 3.7])

Write $q = |\mathfrak{o}/\mathfrak{p}|$ and $d = \dim \underline{\mathbf{G}}$. The sequence $(q^{-k \cdot d} \zeta_{G^k})_{k \geq 1}$ is a constant sequence of functions, i.e.

$$q^{-kd} \zeta_{G^k} = q^{-md} \zeta_{G^m} \quad \text{for all } k, m \geq 1.$$

Proof

By the Kirillov orbit method and Lie-correspondence:

$$\zeta_{G^k}(s) = \sum_{\Omega \in \text{Ad}(G^k) \backslash \widehat{\mathfrak{g}^k}} |\Omega|^{-s/2} = \sum_{\omega \in \widehat{\mathfrak{g}^k}} |\mathfrak{g}^k : \text{Rad}_{\mathfrak{g}^k}(\omega)|^{-s/2-1} \quad (*)$$

where $\text{Rad}_{\mathfrak{g}^k}(\omega) = \{x \in \mathfrak{g}^k : \omega([x, y]) = 1 \text{ for all } y \in \mathfrak{g}^k\}$.

The bijection

$$\varphi = (x \mapsto \pi^{k-1}x) : \mathfrak{g}^1 \rightarrow \mathfrak{g}^k$$

where $\mathfrak{p} = \pi\mathfrak{o}$, induces a surjective $q^{(k-1)d}$ -to-one map

$$\varphi^* : \widehat{\mathfrak{g}^k} \rightarrow \widehat{\mathfrak{g}^1} \text{ given by } \varphi^*(\omega)(x) = \omega(\pi^{k-1}x)$$

with $\varphi(\text{Rad}_{\mathfrak{g}^1}(\varphi^*\omega)) = \text{Rad}_{\mathfrak{g}^k}(\omega)$.

$$(*) = q^{(k-1)d} \sum_{\omega' = \varphi^*(\omega) \in \widehat{\mathfrak{g}^1}} |\mathfrak{p}\mathfrak{g} : \text{Rad}_{\mathfrak{g}^1}(\omega')|^{-s/2-1} = q^{(k-1)d} \zeta_{G^1}(s).$$

Proof

By the Kirillov orbit method and Lie-correspondence:

$$\zeta_{G^k}(s) = \sum_{\Omega \in \text{Ad}(G^k) \backslash \widehat{\mathfrak{g}^k}} |\Omega|^{-s/2} = \sum_{\omega \in \widehat{\mathfrak{g}^k}} |\mathfrak{g}^k : \text{Rad}_{\mathfrak{g}^k}(\omega)|^{-s/2-1} \quad (*)$$

where $\text{Rad}_{\mathfrak{g}^k}(\omega) = \{x \in \mathfrak{g}^k : \omega([x, y]) = 1 \text{ for all } y \in \mathfrak{g}^k\}$.

The bijection

$$\varphi = (x \mapsto \pi^{k-1}x) : \mathfrak{g}^1 \rightarrow \mathfrak{g}^k$$

where $\mathfrak{p} = \pi\mathfrak{o}$, induces a surjective $q^{(k-1)d}$ -to-one map

$$\varphi^* : \widehat{\mathfrak{g}^k} \rightarrow \widehat{\mathfrak{g}^1} \text{ given by } \varphi^*(\omega)(x) = \omega(\pi^{k-1}x)$$

with $\varphi(\text{Rad}_{\mathfrak{g}^1}(\varphi^*\omega)) = \text{Rad}_{\mathfrak{g}^k}(\omega)$.

$$(*) = q^{(k-1)d} \sum_{\omega' = \varphi^*(\omega) \in \widehat{\mathfrak{g}^1}} |\mathfrak{p}\mathfrak{g} : \text{Rad}_{\mathfrak{g}^1}(\omega')|^{-s/2-1} = q^{(k-1)d} \zeta_{G^1}(s).$$

Proof

By the Kirillov orbit method and Lie-correspondence:

$$\zeta_{G^k}(s) = \sum_{\Omega \in \text{Ad}(G^k) \backslash \widehat{\mathfrak{g}^k}} |\Omega|^{-s/2} = \sum_{\omega \in \widehat{\mathfrak{g}^k}} |\mathfrak{g}^k : \text{Rad}_{\mathfrak{g}^k}(\omega)|^{-s/2-1} \quad (*)$$

where $\text{Rad}_{\mathfrak{g}^k}(\omega) = \{x \in \mathfrak{g}^k : \omega([x, y]) = 1 \text{ for all } y \in \mathfrak{g}^k\}$.

The bijection

$$\varphi = (x \mapsto \pi^{k-1}x) : \mathfrak{g}^1 \rightarrow \mathfrak{g}^k$$

where $\mathfrak{p} = \pi\mathfrak{o}$, induces a surjective $q^{(k-1)d}$ -to-one map

$$\varphi^* : \widehat{\mathfrak{g}^k} \rightarrow \widehat{\mathfrak{g}^1} \text{ given by } \varphi^*(\omega)(x) = \omega(\pi^{k-1}x)$$

with $\varphi(\text{Rad}_{\mathfrak{g}^1}(\varphi^*\omega)) = \text{Rad}_{\mathfrak{g}^k}(\omega)$.

$$(*) = q^{(k-1)d} \sum_{\omega' = \varphi^*(\omega) \in \widehat{\mathfrak{g}^1}} |\mathfrak{p}\mathfrak{g} : \text{Rad}_{\mathfrak{g}^1}(\omega')|^{-s/2-1} = q^{(k-1)d} \zeta_{G^1}(s).$$

Proof

By the Kirillov orbit method and Lie-correspondence:

$$\zeta_{G^k}(s) = \sum_{\Omega \in \text{Ad}(G^k) \backslash \widehat{\mathfrak{g}^k}} |\Omega|^{-s/2} = \sum_{\omega \in \widehat{\mathfrak{g}^k}} |\mathfrak{g}^k : \text{Rad}_{\mathfrak{g}^k}(\omega)|^{-s/2-1} \quad (*)$$

where $\text{Rad}_{\mathfrak{g}^k}(\omega) = \{x \in \mathfrak{g}^k : \omega([x, y]) = 1 \text{ for all } y \in \mathfrak{g}^k\}$.

The bijection

$$\varphi = (x \mapsto \pi^{k-1}x) : \mathfrak{g}^1 \rightarrow \mathfrak{g}^k$$

where $\mathfrak{p} = \pi\mathfrak{o}$, induces a surjective $q^{(k-1)d}$ -to-one map

$$\varphi^* : \widehat{\mathfrak{g}^k} \rightarrow \widehat{\mathfrak{g}^1} \text{ given by } \varphi^*(\omega)(x) = \omega(\pi^{k-1}x)$$

with $\varphi(\text{Rad}_{\mathfrak{g}^1}(\varphi^*\omega)) = \text{Rad}_{\mathfrak{g}^k}(\omega)$.

$$(*) = q^{(k-1)d} \sum_{\omega' = \varphi^*(\omega) \in \widehat{\mathfrak{g}^1}} |\mathfrak{p}\mathfrak{g} : \text{Rad}_{\mathfrak{g}^1}(\omega')|^{-s/2-1} = q^{(k-1)d} \zeta_{G^1}(s).$$

Motivation

“The anomaly of the zeroth principal congruence subgroup”

The observation **does not** extend to $k = 0$. E.g.

$$\zeta_{\mathrm{SL}_2(0)}(s) = \zeta_{\mathrm{SL}_2(\mathbb{F}_p)}(s) + \frac{4q \left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2} (q^2 - q)^{-s} + \frac{(q-1)^2}{2} (q^2 + q)^{-s}}{1 - q^{1-s}}$$

[Jaikin-Zapirain 06, Theorem 7.5] ,

$$q^{-3k} \zeta_{\mathrm{SL}_2^k(0)}(s) = \frac{1 - q^{-2-s}}{1 - q^{1-s}} \quad \text{for all } k \geq 1$$

[AKOV 12, Theorem 1.2].

Motivation

“The anomaly of the zeroth principal congruence subgroup”

The observation **does not** extend to $k = 0$. E.g.

$$\zeta_{\mathrm{SL}_2(o)}(s) = \zeta_{\mathrm{SL}_2(\mathbb{F}_p)}(s) + \frac{4q \left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2} (q^2 - q)^{-s} + \frac{(q-1)^2}{2} (q^2 + q)^{-s}}{1 - q^{1-s}}$$

[Jaikin-Zapirain 06, Theorem 7.5] ,

$$q^{-3k} \zeta_{\mathrm{SL}_2^k(o)}(s) = \frac{1 - q^{-2-s}}{1 - q^{1-s}} \quad \text{for all } k \geq 1$$

[AKOV 12, Theorem 1.2].

Motivation

Towards an approximate Kirillov formula

The ‘anomaly’ above is a consequence of the lack of a Kirillov formula for G . Namely,

$$\zeta_G(s) \neq \sum_{\omega \in \widehat{\mathfrak{g}}} |\mathfrak{g} : \text{Rad}(\omega)|^{-s/2-1}$$

However, when considering congruence quotients of G , empirical evidence suggest that certain aspects of such a formula emerge.

Motivation

Towards an approximate Kirillov formula

The ‘anomaly’ above is a consequence of the lack of a Kirillov formula for G . Namely,

$$\zeta_G(s) \neq \sum_{\omega \in \widehat{\mathfrak{g}}} |\mathfrak{g} : \text{Rad}(\omega)|^{-s/2-1}$$

However, when considering congruence quotients of G , empirical evidence suggest that certain aspects of such a formula emerge.

Conjecture/Goal [Onn-Sh, ongoing]

There exists a relation \approx , independent of the residual characteristic of \mathfrak{o} , such that

$$\zeta_{G_k}(s) \approx \sum_{\omega \in \widehat{\mathfrak{g}_k}} |\mathfrak{g}_k : \text{Rad}_{\mathfrak{g}_k}(\omega)|^{-s/2-1}.$$

Here $G_k = G/G^k$ and $\mathfrak{g}_k = \mathfrak{g}/\mathfrak{g}^k$.

Conjecture/Goal [Onn-Sh, ongoing]

There exists a relation \approx , independent of the residual characteristic of \mathfrak{o} , such that

$$\zeta_{G_k}(s) \approx \sum_{\omega \in \widehat{\mathfrak{g}}_k} |\mathfrak{g}_k : \text{Rad}_{\mathfrak{g}_k}(\omega)|^{-s/2-1}.$$

Here $G_k = G/G^k$ and $\mathfrak{g}_k = \mathfrak{g}/\mathfrak{g}^k$.

The main result

Some definitions

Let $0 \in C \subseteq \mathbb{Q}$ a finite set and $r \in \mathbb{N}$

- ① Given $q \in \mathbb{N}$, let $\mathcal{D}_{C,r}(q)$ denote the set of Dirichlet polynomials with non-negative coefficients of the form

$$f(s) = \sum_{i=1}^r u_i(q) v_i(q)^{-s} \quad (s \in \mathbb{C})$$

with $u_1, \dots, u_r, v_1, \dots, v_r \in \mathbb{Q}[t]$ with coefficients in C

The main result

Some definitions

2. Given $q \in \mathbb{N}$ and f_1, f_2 Dirichlet polynomials with non-negative coefficients, write

$$f_1 \sim_{(C,q,r)} f_2$$

if

- $f_j(s) = \sum_{i=1}^r u_{i,j}(q) v_{i,j}(q)^{-s} \in \mathcal{D}_{C,r}(q)$, for $j = 1, 2$
- $\deg(u_{i,1}) = \deg(u_{i,2})$ and $\deg(v_{i,1}) = \deg(v_{i,2})$ for all $i = 1, \dots, r$

The main result

Some definitions

2. Given $q \in \mathbb{N}$ and f_1, f_2 Dirichlet polynomials with non-negative coefficients, write

$$f_1 \sim_{(C,q,r)} f_2$$

if

- $f_j(s) = \sum_{i=1}^r u_{i,j}(q) v_{i,j}(q)^{-s} \in \mathcal{D}_{C,r}(q)$, for $j = 1, 2$
- $\deg(u_{i,1}) = \deg(u_{i,2})$ and $\deg(v_{i,1}) = \deg(v_{i,2})$ for all $i = 1, \dots, r$

The main result

Some definitions

2. Given $q \in \mathbb{N}$ and f_1, f_2 Dirichlet polynomials with non-negative coefficients, write

$$f_1 \sim_{(C,q,r)} f_2$$

if

- $f_j(s) = \sum_{i=1}^r u_{i,j}(q) v_{i,j}(q)^{-s} \in \mathcal{D}_{C,r}(q)$, for $j = 1, 2$
- $\deg(u_{i,1}) = \deg(u_{i,2})$ and $\deg(v_{i,1}) = \deg(v_{i,2})$ for all $i = 1, \dots, r$

The main result

An example

Recall

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)}(s) = (q-1) \left(1 + q^{-s} + \frac{q-2}{2}(q+1)^{-s} + \frac{q-1}{2}(q-1)^{-s} \right).$$

Let $\mathcal{S}_2(\mathbb{F}_q)$ denote the set of similarity classes of 2×2 matrices over \mathbb{F}_q and define $\epsilon_{\mathrm{gl}_2(\mathbb{F}_q)}(s) = \sum_{[x] \in \mathcal{S}_2(\mathbb{F}_q)} |\mathrm{gl}_2(\mathbb{F}_q) : \mathbb{C}_{\mathrm{gl}_2(\mathbb{F}_q)}(x)|^{-s/2}$. Then:

$$\begin{aligned} \epsilon_{\mathrm{gl}_2(\mathbb{F}_q)} &= q(1 + (q-1)q^{-s}) \\ &= q \left(1 + \frac{1}{2}q^{-s} + \frac{q-2}{2}q^{-s} + \frac{q-1}{2}q^{-s} \right) \end{aligned}$$

Thus

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)} \sim_{(C,q,4)} \epsilon_{\mathrm{gl}_2(\mathbb{F}_q)} \text{ with } C = \frac{1}{2}\mathbb{Z} \cap [-3, 3].$$

The main result

An example

Recall

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)}(s) = (q-1) \left(1 + q^{-s} + \frac{q-2}{2}(q+1)^{-s} + \frac{q-1}{2}(q-1)^{-s} \right).$$

Let $\mathcal{S}_2(\mathbb{F}_q)$ denote the set of similarity classes of 2×2 matrices over \mathbb{F}_q and define $\epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)}(s) = \sum_{[x] \in \mathcal{S}_2(\mathbb{F}_q)} |\mathfrak{gl}_2(\mathbb{F}_q) : \mathbf{C}_{\mathfrak{gl}_2(\mathbb{F}_q)}(x)|^{-s/2}$. Then:

$$\begin{aligned} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} &= q(1 + (q-1)q^{-s}) \\ &= q \left(1 + \frac{1}{2}q^{-s} + \frac{q-2}{2}q^{-s} + \frac{q-1}{2}q^{-s} \right) \end{aligned}$$

Thus

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)} \sim_{(C,q,4)} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} \text{ with } C = \frac{1}{2}\mathbb{Z} \cap [-3, 3].$$

The main result

An example

Recall

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)}(s) = (q-1) \left(1 + q^{-s} + \frac{q-2}{2}(q+1)^{-s} + \frac{q-1}{2}(q-1)^{-s} \right).$$

Let $\mathcal{S}_2(\mathbb{F}_q)$ denote the set of similarity classes of 2×2 matrices over \mathbb{F}_q and define $\epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)}(s) = \sum_{[x] \in \mathcal{S}_2(\mathbb{F}_q)} |\mathfrak{gl}_2(\mathbb{F}_q) : \mathbf{C}_{\mathfrak{gl}_2(\mathbb{F}_q)}(x)|^{-s/2}$. Then:

$$\begin{aligned} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} &= q(1 + (q-1)q^{-s}) \\ &= q \left(1 + \frac{1}{2}q^{-s} + \frac{q-2}{2}q^{-s} + \frac{q-1}{2}q^{-s} \right) \end{aligned}$$

Thus

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)} \sim_{(C,q,4)} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} \text{ with } C = \frac{1}{2}\mathbb{Z} \cap [-3, 3].$$

The main result

An example

Recall

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)}(s) = (q-1) \left(1 + q^{-s} + \frac{q-2}{2}(q+1)^{-s} + \frac{q-1}{2}(q-1)^{-s} \right).$$

Let $\mathcal{S}_2(\mathbb{F}_q)$ denote the set of similarity classes of 2×2 matrices over \mathbb{F}_q and define $\epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)}(s) = \sum_{[x] \in \mathcal{S}_2(\mathbb{F}_q)} |\mathfrak{gl}_2(\mathbb{F}_q) : \mathbf{C}_{\mathfrak{gl}_2(\mathbb{F}_q)}(x)|^{-s/2}$. Then:

$$\begin{aligned} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} &= q(1 + (q-1)q^{-s}) \\ &= q \left(1 + \frac{1}{2}q^{-s} + \frac{q-2}{2}q^{-s} + \frac{q-1}{2}q^{-s} \right) \end{aligned}$$

Thus

$$\zeta_{\mathrm{GL}_2(\mathbb{F}_q)} \sim_{(C,q,4)} \epsilon_{\mathfrak{gl}_2(\mathbb{F}_q)} \text{ with } C = \frac{1}{2}\mathbb{Z} \cap [-3, 3].$$

The main result

Given a connected reductive \mathbb{F}_q -defined group $\underline{\mathbf{G}}$ with Lie algebra $\underline{\mathfrak{g}}$, define the *adjoint class function* of $\underline{\mathfrak{g}}(\mathbb{F}_q)$

$$\epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s) = \sum_{[x] \in \text{Ad}(\underline{\mathbf{G}}(\mathbb{F}_q)) \backslash \backslash \underline{\mathfrak{g}}(\mathbb{F}_q)} |\underline{\mathfrak{g}}(\mathbb{F}_q) : \mathbf{C}_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(x)|^{-s/2}.$$

Theorem (Sh)

Let $\mathcal{R} = (\Phi, X, \Phi^\vee, Y)$ be a root datum. There exist a prime $p_{\mathcal{R}}$, a finite set $C_{\mathcal{R}}$, and a natural number $r_{\mathcal{R}}$, such that the following holds for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) > p_{\mathcal{R}}$. For any $\underline{\mathbf{G}}$ connected reductive \mathbb{F}_q -defined group with root datum \mathcal{R} and Lie algebra $\underline{\mathfrak{g}}$

$$\zeta_{\underline{\mathbf{G}}(\mathbb{F}_q)}(s) \sim_{(C_{\mathcal{R}}, q, r_{\mathcal{R}})} \epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s).$$

The main result

Given a connected reductive \mathbb{F}_q -defined group $\underline{\mathbf{G}}$ with Lie algebra $\underline{\mathfrak{g}}$, define the *adjoint class function* of $\underline{\mathfrak{g}}(\mathbb{F}_q)$

$$\epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s) = \sum_{[x] \in \text{Ad}(\underline{\mathbf{G}}(\mathbb{F}_q)) \backslash \backslash \underline{\mathfrak{g}}(\mathbb{F}_q)} |\underline{\mathfrak{g}}(\mathbb{F}_q) : \mathbf{C}_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(x)|^{-s/2}.$$

Theorem (Sh)

Let $\mathcal{R} = (\Phi, X, \Phi^\vee, Y)$ be a root datum. There exist a prime $p_{\mathcal{R}}$, a finite set $\mathcal{C}_{\mathcal{R}}$, and a natural number $r_{\mathcal{R}}$, such that the following holds for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) > p_{\mathcal{R}}$. For any $\underline{\mathbf{G}}$ connected reductive \mathbb{F}_q -defined group with root datum \mathcal{R} and Lie algebra $\underline{\mathfrak{g}}$

$$\zeta_{\underline{\mathbf{G}}(\mathbb{F}_q)}(s) \sim_{(\mathcal{C}_{\mathcal{R}}, q, r_{\mathcal{R}})} \epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s).$$

The main result

Given a connected reductive \mathbb{F}_q -defined group $\underline{\mathbf{G}}$ with Lie algebra $\underline{\mathfrak{g}}$, define the *adjoint class function* of $\underline{\mathfrak{g}}(\mathbb{F}_q)$

$$\epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s) = \sum_{[x] \in \text{Ad}(\underline{\mathbf{G}}(\mathbb{F}_q)) \backslash \backslash \underline{\mathfrak{g}}(\mathbb{F}_q)} |\underline{\mathfrak{g}}(\mathbb{F}_q) : \mathbf{C}_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(x)|^{-s/2}.$$

Theorem (Sh)

Let $\mathcal{R} = (\Phi, X, \Phi^\vee, Y)$ be a root datum. There exist a prime $p_{\mathcal{R}}$, a finite set $C_{\mathcal{R}}$, and a natural number $r_{\mathcal{R}}$, such that the following holds for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) > p_{\mathcal{R}}$. For any $\underline{\mathbf{G}}$ connected reductive \mathbb{F}_q -defined group with root datum \mathcal{R} and Lie algebra $\underline{\mathfrak{g}}$

$$\zeta_{\underline{\mathbf{G}}(\mathbb{F}_q)}(s) \sim_{(C_{\mathcal{R}}, q, r_{\mathcal{R}})} \epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s).$$

The main result

Given a connected reductive \mathbb{F}_q -defined group $\underline{\mathbf{G}}$ with Lie algebra $\underline{\mathfrak{g}}$, define the *adjoint class function* of $\underline{\mathfrak{g}}(\mathbb{F}_q)$

$$\epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s) = \sum_{[x] \in \text{Ad}(\underline{\mathbf{G}}(\mathbb{F}_q)) \backslash \backslash \underline{\mathfrak{g}}(\mathbb{F}_q)} |\underline{\mathfrak{g}}(\mathbb{F}_q) : \mathbf{C}_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(x)|^{-s/2}.$$

Theorem (Sh)

Let $\mathcal{R} = (\Phi, X, \Phi^\vee, Y)$ be a root datum. There exist a prime $p_{\mathcal{R}}$, a finite set $C_{\mathcal{R}}$, and a natural number $r_{\mathcal{R}}$, such that the following holds for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) > p_{\mathcal{R}}$. For any $\underline{\mathbf{G}}$ connected reductive \mathbb{F}_q -defined group with root datum \mathcal{R} and Lie algebra $\underline{\mathfrak{g}}$

$$\zeta_{\underline{\mathbf{G}}(\mathbb{F}_q)}(s) \sim_{(C_{\mathcal{R}}, q, r_{\mathcal{R}})} \epsilon_{\underline{\mathfrak{g}}(\mathbb{F}_q)}(s).$$

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.
- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_p^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$
 - $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and
 - $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.

- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_{p'}^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$

• $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and

• $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.

- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_{p'}^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$

• $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and

• $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.
- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_{p'}^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$
 - $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and
 - $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.
- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_{p'}^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$
 - $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and
 - $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Key component of proof

Jordan-type decompositions

Write $G = \underline{G}(\mathbb{F}_q)$ and $\mathfrak{g} = \underline{\mathfrak{g}}(\mathbb{F}_q)$.

- $\epsilon_{\mathfrak{g}}(s) = \sum_{\substack{[x_{ss}] \in \text{Ad}(G) \backslash \mathfrak{g} \\ \text{semisimple}}} |\mathfrak{g} : \mathbf{C}_{\mathfrak{g}}(x_{ss})|^{-s/2} \cdot \epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$
 - $\epsilon_{\mathbf{C}_{\mathfrak{g}}(x_{ss})}^{\text{nil}}(s)$ enumerates *nilpotent* $\text{Ad}(\mathbf{C}_G(x_{ss}))$ -classes.
- $\zeta_G(s) = \sum_{\substack{[g_{ss}] \in \text{Ad}(G^*) \backslash G^* \\ \text{semisimple}}} |G^* : \mathbf{C}_{G^*}(g_{ss})|_{p'}^{-s} \cdot \zeta_{\mathbf{C}_{G^*}(g_{ss})}^{\text{unip}}(s)$
 - $G^* = \underline{G}^*(\mathbb{F}_q)$ is the dual algebraic group, and
 - $\zeta_H^{\text{unip}}(s) = \sum_{\chi \in \mathcal{E}(H, (1))} \chi(1)^{-s}$ enumerates dimensions of *unipotent* characters of a reductive \mathbb{F}_q -group $H = \underline{H}(\mathbb{F}_q)$ [Lusztig 84, Theorem 4.23].

Example- unipotent zeta function of $\mathrm{GL}_n(\mathbb{F}_q)$

By definition, an irreducible character $\chi \in \mathrm{Irr}(G)$ is *unipotent* if there exists a \mathbb{F}_q -defined maximal torus $\mathbf{T} \subseteq \mathbf{G}$ such that χ occurs in $R_{\mathbf{T}(\mathbb{F}_q)}^G(\mathbf{1})$.

In $\mathrm{GL}_n(\mathbb{F}_q)$ all unipotent characters are principal series, i.e. are constituents of $R_{\mathbf{D}_n(\mathbb{F}_q)}^{\mathrm{GL}_n(\mathbb{F}_q)}(\mathbf{1})$, with \mathbf{D}_n the diagonal torus. Furthermore:

Theorem (Steinberg 49, Green 54)

There exists a bijection

$$\left\{ \begin{array}{c} \text{Unipote characters} \\ \text{of } \mathrm{GL}_n(\mathbb{F}_q) \end{array} \right\} \leftrightarrow \{ \text{partitions of } n \}$$

Given $\lambda \vdash n$, the corresponding character χ_λ is of degree $f_\lambda(q)$, where $f_\lambda(t) \in \mathbb{Z}[t]$ is of degree $\frac{1}{2}(n^2 - \|\lambda\|_2^2)$ and independent of \mathbb{F}_q .

Example- unipotent zeta function of $\mathrm{GL}_n(\mathbb{F}_q)$

By definition, an irreducible character $\chi \in \mathrm{Irr}(G)$ is *unipotent* if there exists a \mathbb{F}_q -defined maximal torus $\mathbf{T} \subseteq \mathbf{G}$ such that χ occurs in $R_{\mathbf{T}(\mathbb{F}_q)}^G(\mathbf{1})$.

In $\mathrm{GL}_n(\mathbb{F}_q)$ all unipotent characters are principal series, i.e. are constituents of $R_{\mathbf{D}_n(\mathbb{F}_q)}^{\mathrm{GL}_n(\mathbb{F}_q)}(\mathbf{1})$, with \mathbf{D}_n the diagonal torus. Furthermore:

Theorem (Steinberg 49, Green 54)

There exists a bijection

$$\left\{ \begin{array}{c} \text{Unipote characters} \\ \text{of } \mathrm{GL}_n(\mathbb{F}_q) \end{array} \right\} \leftrightarrow \{ \text{partitions of } n \}$$

Given $\lambda \vdash n$, the corresponding character χ_λ is of degree $f_\lambda(q)$, where $f_\lambda(t) \in \mathbb{Z}[t]$ is of degree $\frac{1}{2}(n^2 - \|\lambda\|_2^2)$ and independent of \mathbb{F}_q .

Example- unipotent zeta function of $GL_n(\mathbb{F}_q)$

By definition, an irreducible character $\chi \in \text{Irr}(G)$ is *unipotent* if there exists a \mathbb{F}_q -defined maximal torus $\mathbf{T} \subseteq \mathbf{G}$ such that χ occurs in $R_{\mathbf{T}(\mathbb{F}_q)}^G(\mathbf{1})$.

In $GL_n(\mathbb{F}_q)$ all unipotent characters are principal series, i.e. are constituents of $R_{\mathbf{D}_n(\mathbb{F}_q)}^{GL_n(\mathbb{F}_q)}(\mathbf{1})$, with \mathbf{D}_n the diagonal torus. Furthermore:

Theorem (Steinberg 49, Green 54)

There exists a bijection

$$\left\{ \begin{array}{l} \text{Unipote characters} \\ \text{of } GL_n(\mathbb{F}_q) \end{array} \right\} \leftrightarrow \{\text{partitions of } n\}$$

Given $\lambda \vdash n$, the corresponding character χ_λ is of degree $f_\lambda(q)$, where $f_\lambda(t) \in \mathbb{Z}[t]$ is of degree $\frac{1}{2}(n^2 - \|\lambda\|_2^2)$ and independent of \mathbb{F}_q .

Example- unipotent zeta function of $GL_n(\mathbb{F}_q)$

Nilpotent similarity classes are parameterized by partitions of n as well, where given $\lambda = (\lambda_1, \dots, \lambda_r) \vdash n$, a representative for the associated similarity class is $x_\lambda = \text{diag}(J_{\lambda_1}(0), \dots, J_{\lambda_r}(0))$. One has $\dim_{\mathbb{F}_q} \mathcal{C}_{gl_n(\mathbb{F}_q)}(x_\lambda) = \|\lambda^*\|_2^2$, where λ^* denotes the dual partition. Hence

$$\epsilon_{gl_n(\mathbb{F}_q)}^{\text{nil}}(s) = \sum_{\lambda \vdash n} q^{-\frac{1}{2}(n^2 - \|\lambda^*\|_2^2)s}.$$

Example- unipotent zeta function of $GL_n(\mathbb{F}_q)$

Nilpotent similarity classes are parameterized by partitions of n as well, where given $\lambda = (\lambda_1, \dots, \lambda_r) \vdash n$, a representative for the associated similarity class is $x_\lambda = \text{diag}(J_{\lambda_1}(0), \dots, J_{\lambda_r}(0))$. One has $\dim_{\mathbb{F}_q} \mathbf{C}_{gl_n(\mathbb{F}_q)}(x_\lambda) = \|\lambda^*\|_2^2$, where λ^* denotes the dual partition.

Hence

$$\epsilon_{gl_n(\mathbb{F}_q)}^{\text{nil}}(s) = \sum_{\lambda \vdash n} q^{-\frac{1}{2}(n^2 - \|\lambda^*\|_2^2)s}.$$

Example- unipotent zeta function of $GL_n(\mathbb{F}_q)$

Nilpotent similarity classes are parameterized by partitions of n as well, where given $\lambda = (\lambda_1, \dots, \lambda_r) \vdash n$, a representative for the associated similarity class is $x_\lambda = \text{diag}(J_{\lambda_1}(0), \dots, J_{\lambda_r}(0))$. One has $\dim_{\mathbb{F}_q} \mathbf{C}_{\mathfrak{gl}_n(\mathbb{F}_q)}(x_\lambda) = \|\lambda^*\|_2^2$, where λ^* denotes the dual partition. Hence

$$\epsilon_{\mathfrak{gl}_n(\mathbb{F}_q)}^{\text{nil}}(s) = \sum_{\lambda \vdash n} q^{-\frac{1}{2}(n^2 - \|\lambda^*\|_2^2)s}.$$

Example Unipotent zeta functions of $\mathrm{GL}_n(\mathbb{F}_q)$

We deduce

Lemma

For any finite field \mathbb{F}_q ,

$$\zeta_{\mathrm{GL}_n(\mathbb{F}_q)}^{\mathrm{unip}}(s) = \sum_{\lambda \vdash n} f_{\lambda}(q)^{-s} \sim_{(C_n, q, r_n)} \sum_{\lambda \vdash n} q^{-\frac{1}{2}(n^2 - \|\lambda^*\|_2^2)s} = \epsilon_{\mathfrak{gl}_n(\mathbb{F}_q)}^{\mathrm{nil}}$$

where r_n is the number of partitions of n , and C_n is the set of all coefficients occurring in $\{f_{\lambda} : \lambda \vdash n\} \cup \{1, 0\}$.

Remark. The result above extends to the generality of $\underline{\mathbf{G}}(\mathbb{F}_q)$, with $\underline{\mathbf{G}}$ a connected reductive algebraic group.

Example Unipotent zeta functions of $\mathrm{GL}_n(\mathbb{F}_q)$

We deduce

Lemma

For any finite field \mathbb{F}_q ,

$$\zeta_{\mathrm{GL}_n(\mathbb{F}_q)}^{\mathrm{unip}}(s) = \sum_{\lambda \vdash n} f_{\lambda}(q)^{-s} \sim_{(C_n, q, r_n)} \sum_{\lambda \vdash n} q^{-\frac{1}{2}(n^2 - \|\lambda^*\|_2^2)s} = \epsilon_{\mathfrak{gl}_n(\mathbb{F}_q)}^{\mathrm{nil}}$$

where r_n is the number of partitions of n , and C_n is the set of all coefficients occurring in $\{f_{\lambda} : \lambda \vdash n\} \cup \{1, 0\}$.

Remark. The result above extends to the generality of $\underline{\mathbf{G}}(\mathbb{F}_q)$, with $\underline{\mathbf{G}}$ a connected reductive algebraic group.

Application of main result to arithmetic groups

Let K be a global number field with ring of integers \mathcal{O} and let $\underline{\mathbf{G}}$ be an affine algebraic group scheme over \mathcal{O} whose generic fiber $\underline{\mathbf{G}}_K$ is connected, simply connected and semisimple of absolute rank $r = \mathrm{rk}(\underline{\mathbf{G}})$ and absolute root system Φ .

The assumption of simply-connectedness implies that the constant coefficient of $\zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}$ is 1 for all but finitely many \mathfrak{p} 's. Therefore we may consider their infinite product.

Let $\mathrm{Irr}^{\mathrm{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))$ denote the set of equivalence classes of irreducible representations of $\underline{\mathbf{G}}(\mathcal{O})$, which factor through a quotient of the form $\underline{\mathbf{G}}(\mathcal{O}/I)$ with I a square-free ideal.

Application of main result to arithmetic groups

Let K be a global number field with ring of integers \mathcal{O} and let $\underline{\mathbf{G}}$ be an affine algebraic group scheme over \mathcal{O} whose generic fiber $\underline{\mathbf{G}}_K$ is connected, simply connected and semisimple of absolute rank $r = \mathrm{rk}(\underline{\mathbf{G}})$ and absolute root system Φ .

The assumption of simply-connectedness implies that the constant coefficient of $\zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}$ is 1 for all but finitely many \mathfrak{p} 's. Therefore we may consider their infinite product.

Let $\mathrm{Irr}^{\mathrm{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))$ denote the set of equivalence classes of irreducible representations of $\underline{\mathbf{G}}(\mathcal{O})$, which factor through a quotient of the form $\underline{\mathbf{G}}(\mathcal{O}/I)$ with I a square-free ideal.

Application of main result to arithmetic groups

Let K be a global number field with ring of integers \mathcal{O} and let $\underline{\mathbf{G}}$ be an affine algebraic group scheme over \mathcal{O} whose generic fiber $\underline{\mathbf{G}}_K$ is connected, simply connected and semisimple of absolute rank $r = \mathrm{rk}(\underline{\mathbf{G}})$ and absolute root system Φ .

The assumption of simply-connectedness implies that the constant coefficient of $\zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}$ is 1 for all but finitely many \mathfrak{p} 's. Therefore we may consider their infinite product.

Let $\mathrm{Irr}^{\mathrm{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))$ denote the set of equivalence classes of irreducible representations of $\underline{\mathbf{G}}(\mathcal{O})$, which factor through a quotient of the form $\underline{\mathbf{G}}(\mathcal{O}/I)$ with I a square-free ideal.

Application of main result to arithmetic groups

Theorem (Sh)

The Dirichlet series

$$\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}(s) = \sum_{\chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))} (\dim \chi)^{-s} = \prod_{\mathfrak{p} \nmid \mathcal{O}} \zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}(s)$$

converges on the right half-plane $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi+|} \right\}$.

Furthermore, there exists $\delta > 0$ such that $\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}$ extends to a meromorphic function on $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi+|} - \delta \right\}$ with a unique pole in this region. In particular, for suitable $\alpha, c > 0$ and $\beta \in \mathbb{N}$,

$$\# \{ \chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O})) : \dim \chi \leq N \} \sim c N^{\alpha} \log(N)^{\beta-1}.$$

Application of main result to arithmetic groups

Theorem (Sh)

The Dirichlet series

$$\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}(s) = \sum_{\chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))} (\dim \chi)^{-s} = \prod_{\mathfrak{p} \nmid \mathcal{O}} \zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}(s)$$

converges on the right half-plane $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi^+|} \right\}$.

Furthermore, there exists $\delta > 0$ such that $\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}$ extends to a meromorphic function on $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi^+|} - \delta \right\}$ with a unique pole in this region. In particular, for suitable $\alpha, c > 0$ and $\beta \in \mathbb{N}$,

$$\# \{ \chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O})) : \dim \chi \leq N \} \sim c N^\alpha \log(N)^{\beta-1}.$$

Application of main result to arithmetic groups

Theorem (Sh)

The Dirichlet series

$$\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}(s) = \sum_{\chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O}))} (\dim \chi)^{-s} = \prod_{\mathfrak{p} \nmid \mathcal{O}} \zeta_{\underline{\mathbf{G}}(\mathcal{O}/\mathfrak{p})}(s)$$

converges on the right half-plane $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi^+|} \right\}$.

Furthermore, there exists $\delta > 0$ such that $\zeta_{\underline{\mathbf{G}}(\mathcal{O})}^{\text{sqf}}$ extends to a meromorphic function on $\left\{ \text{Re}(s) > \frac{r+1}{|\Phi^+|} - \delta \right\}$ with a unique pole in this region. In particular, for suitable $\alpha, c > 0$ and $\beta \in \mathbb{N}$,

$$\# \{ \chi \in \text{Irr}^{\text{sqf}}(\underline{\mathbf{G}}(\mathcal{O})) : \dim \chi \leq N \} \sim c N^\alpha \log(N)^{\beta-1}.$$

Questions?

Thank you!

Unipotent representations and nilpotent classes

Given an irreducible root system Φ , let \underline{H}_Φ be the split simple group of adjoint type with root system Φ . We assume $\Psi \leq \Phi \Rightarrow \underline{H}_\Psi \subseteq \underline{H}_\Phi$.

Theorem (Sh)

Let $\Pi \subseteq \Phi$ be a basis and $\tilde{\Pi} = \Pi \cup \{-\alpha_0\}$, where α_0 is the longest root. There exists a prime p_Φ , a finite set $1 \in C_\Phi \subseteq \mathbb{Q}$ and $r_\Phi \in \mathbb{N}$ such that for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) = p$

$$\epsilon_{\text{Lie}(\underline{H}_{\Phi,\Psi})}^{\text{nil}}(\mathbb{F}_q)(s) \sim_{(C_\Phi, q, r_\Phi)} \sum_{\substack{\Psi \leq \Phi \\ |\Psi \cap \tilde{\Pi}| = |\Pi|}} |\underline{H}_\Phi(\mathbb{F}_q) : \underline{H}_\Psi(\mathbb{F}_q)|^{\frac{1}{p^s}} \cdot \zeta_{\underline{H}_\Psi(\mathbb{F}_q)}^{\text{unip}}(s).$$

Remark. Also true for non-split groups.

Unipotent representations and nilpotent classes

Given an irreducible root system Φ , let \underline{H}_Φ be the split simple group of adjoint type with root system Φ . We assume $\Psi \leq \Phi \Rightarrow \underline{H}_\Psi \subseteq \underline{H}_\Phi$.

Theorem (Sh)

Let $\Pi \subseteq \Phi$ be a basis and $\tilde{\Pi} = \Pi \cup \{-\alpha_0\}$, where α_0 is the longest root. There exists a prime p_Φ , a finite set $1 \in C_\Phi \subseteq \mathbb{Q}$ and $r_\Phi \in \mathbb{N}$ such that for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) = p$

$$\epsilon_{\text{Lie}(\underline{H}_{\Phi,\Psi})}^{\text{nil}}(\mathbb{F}_q)(s) \sim_{(C_\Phi, q, r_\Phi)} \sum_{\substack{\Psi \leq \Phi \\ |\Psi \cap \tilde{\Pi}| = |\Pi|}} |\underline{H}_\Phi(\mathbb{F}_q) : \underline{H}_\Psi(\mathbb{F}_q)|^{\frac{1}{r_\Phi}} \cdot \zeta_{\underline{H}_\Psi(\mathbb{F}_q)}^{\text{unip}}(s).$$

Remark. Also true for non-split groups.

Unipotent representations and nilpotent classes

Given an irreducible root system Φ , let $\underline{\mathbf{H}}_\Phi$ be the split simple group of adjoint type with root system Φ . We assume $\Psi \leq \Phi \Rightarrow \underline{\mathbf{H}}_\Psi \subseteq \underline{\mathbf{H}}_\Phi$.

Theorem (Sh)

Let $\Pi \subseteq \Phi$ be a basis and $\tilde{\Pi} = \Pi \cup \{-\alpha_0\}$, where α_0 is the longest root. There exists a prime p_Φ , a finite set $1 \in C_\Phi \subseteq \mathbb{Q}$ and $r_\Phi \in \mathbb{N}$ such that for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) = p$

$$\epsilon_{\text{Lie}(\underline{\mathbf{H}}_{\Phi^\vee})(\mathbb{F}_q)}^{\text{nil}}(s) \sim_{(C_\Phi, q, r_\Phi)} \sum_{\substack{\Psi \leq \Phi \\ |\Psi \cap \tilde{\Pi}| = |\Pi|}} |\underline{\mathbf{H}}_\Phi(\mathbb{F}_q) : \underline{\mathbf{H}}_\Psi(\mathbb{F}_q)|_{p'}^{-s} \cdot \zeta_{\underline{\mathbf{H}}_\Psi(\mathbb{F}_q)}^{\text{unip}}(s).$$

Remark. Also true for non-split groups.

Unipotent representations and nilpotent classes

Given an irreducible root system Φ , let $\underline{\mathbf{H}}_\Phi$ be the split simple group of adjoint type with root system Φ . We assume $\Psi \leq \Phi \Rightarrow \underline{\mathbf{H}}_\Psi \subseteq \underline{\mathbf{H}}_\Phi$.

Theorem (Sh)

Let $\Pi \subseteq \Phi$ be a basis and $\tilde{\Pi} = \Pi \cup \{-\alpha_0\}$, where α_0 is the longest root. There exists a prime p_Φ , a finite set $1 \in C_\Phi \subseteq \mathbb{Q}$ and $r_\Phi \in \mathbb{N}$ such that for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) = p$

$$\epsilon_{\text{Lie}(\underline{\mathbf{H}}_{\Phi^\vee})(\mathbb{F}_q)}^{\text{nil}}(s) \sim_{(C_\Phi, q, r_\Phi)} \sum_{\substack{\Psi \leq \Phi \\ |\Psi \cap \tilde{\Pi}| = |\Pi|}} |\underline{\mathbf{H}}_\Phi(\mathbb{F}_q) : \underline{\mathbf{H}}_\Psi(\mathbb{F}_q)|_{p'}^{-s} \cdot \zeta_{\underline{\mathbf{H}}_\Psi(\mathbb{F}_q)}^{\text{unip}}(s).$$

Remark. Also true for non-split groups.

Unipotent representations and nilpotent classes

Given an irreducible root system Φ , let $\underline{\mathbf{H}}_\Phi$ be the split simple group of adjoint type with root system Φ . We assume $\Psi \leq \Phi \Rightarrow \underline{\mathbf{H}}_\Psi \subseteq \underline{\mathbf{H}}_\Phi$.

Theorem (Sh)

Let $\Pi \subseteq \Phi$ be a basis and $\tilde{\Pi} = \Pi \cup \{-\alpha_0\}$, where α_0 is the longest root. There exists a prime p_Φ , a finite set $1 \in C_\Phi \subseteq \mathbb{Q}$ and $r_\Phi \in \mathbb{N}$ such that for any finite field \mathbb{F}_q with $\text{char}(\mathbb{F}_q) = p$

$$\epsilon_{\text{Lie}(\underline{\mathbf{H}}_{\Phi^\vee})(\mathbb{F}_q)}^{\text{nil}}(s) \sim_{(C_\Phi, q, r_\Phi)} \sum_{\substack{\Psi \leq \Phi \\ |\Psi \cap \tilde{\Pi}| = |\Pi|}} |\underline{\mathbf{H}}_\Phi(\mathbb{F}_q) : \underline{\mathbf{H}}_\Psi(\mathbb{F}_q)|_{p'}^{-s} \cdot \zeta_{\underline{\mathbf{H}}_\Psi(\mathbb{F}_q)}^{\text{unip}}(s).$$

Remark. Also true for non-split groups.

Remark

A non-example

The theorem *may fail* for non-reductive algebraic group.

Let \underline{U}_3 denote the (\mathbb{Z} -defined) algebraic group of upper unitriangular matrices, with Lie algebra \underline{u}_3 .

Then

$$\zeta_{\underline{U}_3(\mathbb{F}_q)}(s) = q^2 + (q-1)q^{-s}$$

and

$$\epsilon_{\underline{u}_3(\mathbb{F}_q)}(s) = q + q^2 \cdot q^{-s/2}.$$

Remark

A non-example

The theorem *may fail* for non-reductive algebraic group.

Let \underline{U}_3 denote the (\mathbb{Z} -defined) algebraic group of upper unitriangular matrices, with Lie algebra \underline{u}_3 .

Then

$$\zeta_{\underline{U}_3(\mathbb{F}_q)}(s) = q^2 + (q-1)q^{-s}$$

and

$$\epsilon_{\underline{u}_3(\mathbb{F}_q)}(s) = q + q^2 \cdot q^{-s/2}.$$

Remark

A non-example

The theorem *may fail* for non-reductive algebraic group.

Let \underline{U}_3 denote the (\mathbb{Z} -defined) algebraic group of upper unitriangular matrices, with Lie algebra \underline{u}_3 .

Then

$$\zeta_{\underline{u}_3(\mathbb{F}_q)}(s) = q^2 + (q - 1)q^{-s}$$

and

$$\epsilon_{\underline{u}_3(\mathbb{F}_q)}(s) = q + q^2 \cdot q^{-s/2}.$$

- [AKOV12] Nir Avni, Benjamin Klopsch, Uri Onn, and Christopher Voll. Representation zeta functions of some compact p -adic analytic groups. 566:295–330, 2012.
- [AKOV13] Nir Avni, Benjamin Klopsch, Uri Onn, and Christopher Voll. Representation zeta functions of compact p -adic analytic groups and arithmetic groups. *Duke Math. J.*, 162(1):111–197, 2013.
- [Gre55] J. A. Green. The characters of the finite general linear groups. *Trans. Amer. Math. Soc.*, 80:402–447, 1955.
- [JZ06] A. Jaikin-Zapirain. Zeta function of representations of compact p -adic analytic groups. *J. Amer. Math. Soc.*, 19(1):91–118, 2006.
- [Lus84] George Lusztig. *Characters of reductive groups over a finite field*, volume 107 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1984.
- [Ste51] R. Steinberg. A geometric approach to the representations of the full linear group over a Galois field. *Trans. Amer. Math. Soc.*, 71:274–282, 1951.