

Representation Zeta Function of Norm 1 Subgroups of Local Division Algebras

Shai Shechter

Ben Gurion University of the Negev, Beer Sheva

`shais@post.bgu.ac.il`

November 5, 2014

Acknowledgement

I wish to express my gratitude to Uri Onn, Benjamin Klopsch and Christopher Voll for their helpful guidance and advisory during the completion of the results presented here.

Definition of ζ_G

- Let G be a topological group. For any $n \in \mathbb{N}$, let $r_n(G) \in \mathbb{N} \cup \{0, \infty\}$ denote the number of isomorphism classes of complex continuous irreducible representations of G .

The group G is called **rigid** if $r_n(G)$ is finite for every n .

Definition of ζ_G

- Let G be a topological group. For any $n \in \mathbb{N}$, let $r_n(G) \in \mathbb{N} \cup \{0, \infty\}$ denote the number of isomorphism classes of complex continuous irreducible representations of G .
The group G is called **rigid** if $r_n(G)$ is finite for every n .
- The group G is said to have **polynomial representation growth** (PRG) if $r_n(G)$ is bounded above by a polynomial in n .

Definition of ζ_G

- Let G be a topological group. For any $n \in \mathbb{N}$, let $r_n(G) \in \mathbb{N} \cup \{0, \infty\}$ denote the number of isomorphism classes of complex continuous irreducible representations of G .
The group G is called **rigid** if $r_n(G)$ is finite for every n .
- The group G is said to have **polynomial representation growth** (PRG) if $r_n(G)$ is bounded above by a polynomial in n .
- In the case where G has PRG, one defines the **representation zeta function of G** to be the Dirichlet generating function

$$\zeta_G(s) = \sum_{n=1}^{\infty} r_n(G) n^{-s}, \quad s \in \mathbb{C}.$$

This series is absolutely convergent in a half-plane $\{z \in \mathbb{C} \mid \Re(z) > \alpha\}$, for some $\alpha \in \mathbb{R}$.

The infimum of such α 's is called the **abscissa of convergence** of ζ_G .

Arithmetic Groups

Let F/\mathbb{Q} be a number field with $\mathfrak{R} \subseteq F$ the ring of integers. Let G be a connected simply-connected algebraic group (e.g. $G = \mathrm{SL}_d$).

- The group $G(\mathfrak{R})$ is said to have the **congruence subgroup property** (CSP) if the map $\widehat{G(\mathfrak{R})} \rightarrow G(\widehat{\mathfrak{R}})$ is an isomorphism.
- It is said to have **weak CSP** (wCSP) if the kernel of the above map is finite.

Arithmetic Groups

Let F/\mathbb{Q} be a number field with $\mathfrak{R} \subseteq F$ the ring of integers. Let G be a connected simply-connected algebraic group (e.g. $G = \mathrm{SL}_d$).

- The group $G(\mathfrak{R})$ is said to have the **congruence subgroup property** (CSP) if the map $\widehat{G(\mathfrak{R})} \rightarrow G(\widehat{\mathfrak{R}})$ is an isomorphism.
- It is said to have **weak CSP** (wCSP) if the kernel of the above map is finite.

Theorem (Lubotzky, Martin)

The group $G(\mathfrak{R})$ has PRG if and only if $G(\mathfrak{R})$ has wCSP.

Arithmetic Groups

Let F/\mathbb{Q} be a number field with $\mathfrak{R} \subseteq F$ the ring of integers. Let G be a connected simply-connected algebraic group (e.g. $G = \mathrm{SL}_d$).

- The group $G(\mathfrak{R})$ is said to have the **congruence subgroup property** (CSP) if the map $\widehat{G(\mathfrak{R})} \rightarrow G(\widehat{\mathfrak{R}})$ is an isomorphism.
- It is said to have **weak CSP** (wCSP) if the kernel of the above map is finite.

Theorem (Lubotzky, Martin)

The group $G(\mathfrak{R})$ has PRG if and only if $G(\mathfrak{R})$ has wCSP.

Example

The group $\mathrm{SL}_d(\mathfrak{R})$ has PRG iff $d > 2$.

Euler Factorization

Theorem (Larsen, Lubotzky)

If $G(\mathfrak{R})$ has CSP then

$$\zeta_{G(\mathfrak{R})}(s) = \zeta_{G(\mathbb{C})}(s)^{|F:\mathbb{Q}|} \cdot \prod_{\substack{\mathfrak{p} \in \mathfrak{R} \\ \text{prime}}} \zeta_{G(\mathfrak{R}_{\mathfrak{p}})}(s).$$

- The archemedian factor $\zeta_{G(\mathbb{C})}(s)$ enumerates irreducible finite-dimensional rational representations of $G(\mathbb{C})$, which are reflected in the root system of G .

Euler Factorization

Theorem (Larsen, Lubotzky)

If $G(\mathfrak{R})$ has CSP then

$$\zeta_{G(\mathfrak{R})}(s) = \zeta_{G(\mathbb{C})}(s)^{|F:\mathbb{Q}|} \cdot \prod_{\substack{\mathfrak{p} \in \mathfrak{R} \\ \text{prime}}} \zeta_{G(\mathfrak{R}_{\mathfrak{p}})}(s).$$

- The archemedian factor $\zeta_{G(\mathbb{C})}(s)$ enumerates irreducible finite-dimensional rational representations of $G(\mathbb{C})$, which are reflected in the root system of G .
- The factors $\zeta_{G(\mathfrak{R}_{\mathfrak{p}})}(s)$ enumerate continuous representations of the compact p -adic analytic groups $G(\mathfrak{R}_{\mathfrak{p}})$, where $\mathfrak{R}_{\mathfrak{p}}$ is the completion of \mathfrak{R} at the prime \mathfrak{p} .

Compact p -adic Analytic Groups

Theorem (Jaikin-Zapirin)

Let G be a finitely generated compact p -adic analytic group. Assume $p > 2$ and that G is rigid. Then there exist rational functions $f_1(T), \dots, f_k(T) \in \mathbb{Q}(T)$ and numbers $n_1, \dots, n_k \in \mathbb{N}$ such that

$$\zeta_G(s) = \sum_{i=1}^k n_i^{-s} f_i(p^{-s}).$$

Compact p -adic Analytic Groups

Theorem (Jaikin-Zapirin)

Let G be a finitely generated compact p -adic analytic group. Assume $p > 2$ and that G is rigid. Then there exist rational functions $f_1(T), \dots, f_k(T) \in \mathbb{Q}(T)$ and numbers $n_1, \dots, n_k \in \mathbb{N}$ such that

$$\zeta_G(s) = \sum_{i=1}^k n_i^{-s} f_i(p^{-s}).$$

Remark

A similar result holds for compact 2-adic analytic groups, under some additional assumptions on G .

Some Notation

- From here on we fix K to be the completion of F at a fixed prime \mathfrak{p} , and let \mathcal{O} denote the completion of \mathfrak{K} and \mathfrak{o} be its maximal ideal, generated by an element $\pi \in \mathcal{O}$.
- We put p to denote the characteristic of the residue field \mathcal{O}/\mathfrak{o} and $q = p^\alpha$ to be its cardinality.

Forms of $\mathrm{SL}_\ell(\mathfrak{R})$

- Let ℓ be a prime number, distinct from p .
- Assume that the group $G(\mathfrak{R})$ is a form of $\mathrm{SL}_\ell(\mathfrak{R})$ for some $\ell \in \mathbb{N}$.
For the case where ℓ is prime, for all but finitely many primes \mathfrak{p} , the group $G(\mathcal{O})$ is isomorphic to one of the following groups:
 - $\mathrm{SL}_\ell(\mathcal{O})$,
 - $\mathrm{SU}_\ell(\mathcal{O})$

Forms of $\mathrm{SL}_\ell(\mathfrak{R})$

- Let ℓ be a prime number, distinct from p .
- Assume that the group $G(\mathfrak{R})$ is a form of $\mathrm{SL}_\ell(\mathfrak{R})$ for some $\ell \in \mathbb{N}$.
For the case where ℓ is prime, for all but finitely many primes \mathfrak{p} , the group $G(\mathcal{O})$ is isomorphic to one of the following groups:
 - $\mathrm{SL}_\ell(\mathcal{O})$,
 - $\mathrm{SU}_\ell(\mathcal{O})$

Forms of $\mathrm{SL}_\ell(\mathfrak{R})$

- Let ℓ be a prime number, distinct from p .
- Assume that the group $G(\mathfrak{R})$ is a form of $\mathrm{SL}_\ell(\mathfrak{R})$ for some $\ell \in \mathbb{N}$.
For the case where ℓ is prime, for all but finitely many primes \mathfrak{p} , the group $G(\mathcal{O})$ is isomorphic to one of the following groups:
 - $\mathrm{SL}_\ell(\mathcal{O})$,
 - $\mathrm{SU}_\ell(\mathcal{O})$
- For the finitely many exceptional primes, the group $G(\mathcal{O})$ is isomorphic to $\mathrm{SL}_1(D)$, the group of norm-1 elements of a central division algebra D of degree ℓ over K .

Forms of $\mathrm{SL}_\ell(\mathfrak{R})$

- Let ℓ be a prime number, distinct from p .
- Assume that the group $G(\mathfrak{R})$ is a form of $\mathrm{SL}_\ell(\mathfrak{R})$ for some $\ell \in \mathbb{N}$.
For the case where ℓ is prime, for all but finitely many primes \mathfrak{p} , the group $G(\mathcal{O})$ is isomorphic to one of the following groups:
 - $\mathrm{SL}_\ell(\mathcal{O})$,
 - $\mathrm{SU}_\ell(\mathcal{O})$
- For the finitely many exceptional primes, the group $G(\mathcal{O})$ is isomorphic to $\mathrm{SL}_1(D)$, the group of norm-1 elements of a central division algebra D of degree ℓ over K .
- In all 3 cases, the group $G(\mathcal{O})$ is a compact p -adic analytic group, and contains a maximal pro- p group $G^1(\mathcal{O})$, given explicitly as the kernel of the map $G(\mathcal{O}) \rightarrow G(\mathcal{O}/\mathfrak{p})$.

Example 1: The case $\ell = 2$

- Jaikin-Zapirin introduced a computation of $\zeta_{SL_2(\mathcal{O})}(s)$, based on a classification of the characters of $SL_2(\mathcal{O})$ and on Clifford theory.

Theorem (Jaikin-Zapirin)

Suppose $p \neq 2$. Then

$$\begin{aligned} \zeta_{SL_2(\mathcal{O})}(s) = & 1 + q^{-s} + \frac{q-3}{2}(q+1)^{-s} + 2\left(\frac{q+1}{2}\right)^{-s} + \frac{q-1}{2}(q-1)^{-s} + \\ & + \frac{4q\left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2}(q^2-q)^{-s} + \frac{(q-1)^2}{2}(q^2+q)^{-s}}{1 - q^{-s+1}}. \end{aligned}$$

Example 1: The case $\ell = 2$

- Jaikin-Zapirin introduced a computation of $\zeta_{\mathrm{SL}_2(\mathcal{O})}(s)$, based on a classification of the characters of $\mathrm{SL}_2(\mathcal{O})$ and on Clifford theory.

Theorem (Jaikin-Zapirin)

Suppose $p \neq 2$. Then

$$\begin{aligned} \zeta_{\mathrm{SL}_2(\mathcal{O})}(s) = & 1 + q^{-s} + \frac{q-3}{2}(q+1)^{-s} + 2\left(\frac{q+1}{2}\right)^{-s} + \frac{q-1}{2}(q-1)^{-s} + \\ & + \frac{4q\left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2}(q^2-q)^{-s} + \frac{(q-1)^2}{2}(q^2+q)^{-s}}{1 - q^{-s+1}}. \end{aligned}$$

- An important feature of this computation is the fact that it is actually independent of the characteristic of K , and holds for positive characteristic as well.

Example 1

- Avni, Klopsch, Onn and Voll introduced a p -adic formalism for the computation of the representation zeta function of a class of pro- p groups. This formalism allows for the calculation of the representation zeta function with via certain Igusa-type zeta integrals.

Example 1

- Avni, Klopsch, Onn and Voll introduced a p -adic formalism for the computation of the representation zeta function of a class of pro- p groups. This formalism allows for the calculation of the representation zeta function via certain Igusa-type zeta integrals.
- This development includes an application of the Kirillov orbit method and an analysis of the adjoint action in an associated Lie-ring defined over \mathcal{O} .
- As an outcome of this formalism, the authors managed to compute the representation zeta function of the maximal pro- p subgroup of $SL_2(\mathcal{O})$ and of $SL_1(D)$ for D a division algebra of degree 2 over K .

Example 1

Division Algebras of Degree 2

A division algebra D of degree 2 over K can be described by the following data:

- Let $a \in \mathcal{O}$ be an invertible element of \mathcal{O} which is not a square.

Example 1

Division Algebras of Degree 2

A division algebra D of degree 2 over K can be described by the following data:

- Let $a \in \mathcal{O}$ be an invertible element of \mathcal{O} which is not a square.
- As a vector space, the division algebra D is spanned by the set $\{1, \mathbf{i}, \mathbf{j}, \mathbf{ij}\}$, where \mathbf{i}, \mathbf{j} are formal symbols.

Example 1

Division Algebras of Degree 2

A division algebra D of degree 2 over K can be described by the following data:

- Let $a \in \mathcal{O}$ be an invertible element of \mathcal{O} which is not a square.
- As a vector space, the division algebra D is spanned by the set $\{1, \mathbf{i}, \mathbf{j}, \mathbf{ij}\}$, where \mathbf{i}, \mathbf{j} are formal symbols.
- The multiplication rules on D are given by the rules

$$\mathbf{i}^2 = a, \quad \mathbf{j}^2 = \pi \quad \text{and} \quad \mathbf{ij} = -\mathbf{ji}.$$

Example 1

Division Algebras of Degree 2

A division algebra D of degree 2 over K can be described by the following data:

- Let $a \in \mathcal{O}$ be an invertible element of \mathcal{O} which is not a square.
- As a vector space, the division algebra D is spanned by the set $\{1, \mathbf{i}, \mathbf{j}, \mathbf{ij}\}$, where \mathbf{i}, \mathbf{j} are formal symbols.
- The multiplication rules on D are given by the rules

$$\mathbf{i}^2 = a, \quad \mathbf{j}^2 = \pi \quad \text{and} \quad \mathbf{ij} = -\mathbf{ji}.$$

- The reduced norm of an element $\mathbf{x} = \alpha + \mathbf{i}\beta + \mathbf{j}\gamma + \mathbf{ij}\delta$ is given explicitly by the formula:

$$\mathrm{Nrd}_{D/K}(\mathbf{x}) = \alpha^2 - a\beta^2 + \pi(\gamma^2 - a\delta^2).$$

Remark

Compare to the case of \mathbb{H} , the Hamiltonian quaternions over \mathbb{R} .

Example 1

Under some mild assumptions on the ramification index of K/\mathbb{Q}_p , Avni, Klopsch, Onn and Voll computed the following.

Theorem (AKOV)

Suppose $p \neq 2$ and let D be a division algebra of degree 2 with $\mathbf{Z}(D) = K$. Then

$$\zeta_{SL_1^1(D)}(s) = q^3 \frac{q - q^{-1-s}}{1 - q^{1-s}},$$

where $SL_1^1(D)$ is the maximal pro- p subgroup of $SL_1(D)$.

Example 1

Applying tools from Clifford theory, the authors obtained the computation for $\mathrm{SL}_1(D)$ as well.

Theorem (AKOV)

Suppose $p \neq 2$ and let D be a division algebra of degree 2 with $\mathbf{Z}(D) = K$. Then

$$\zeta_{\mathrm{SL}_1(D)}(s) = \frac{(q+1)(1-q^{-s}) + 4(q-1)\left(\frac{q+1}{2}\right)^{-s}}{1-q^{1-s}}.$$

Example 2: $\ell = 3$

Theorem (AKOV)

Suppose $p \neq 3$, and let $G(\mathcal{O})$ be $\mathrm{SL}_3(\mathcal{O})$ or $\mathrm{SU}_3(\mathcal{O})$. Then

$$\zeta_{G(\mathcal{O})}(s) = q^8 \frac{1 + u(q)q^{-3-2s} + u(q^{-1})q^{-2-3s} + q^{-5-5s}}{(1 - q^{1-2s})(1 - q^{2-3s})}$$

where

$$u(T) = \begin{cases} T^3 + T^2 - T - 1 - T^{-1} & \text{if } G(\mathcal{O}) = \mathrm{SL}_3(\mathcal{O}) \\ -T^3 + T^2 - T + 1 - T^{-1} & \text{if } G(\mathcal{O}) = \mathrm{SU}_3(\mathcal{O}). \end{cases}$$

Example 2: $\ell = 3$

Theorem (AKOV)

Suppose $p \neq 3$, and let $G(\mathcal{O})$ be $\mathrm{SL}_3(\mathcal{O})$ or $\mathrm{SU}_3(\mathcal{O})$. Then

$$\zeta_{G(\mathcal{O})}(s) = q^8 \frac{1 + u(q)q^{-3-2s} + u(q^{-1})q^{-2-3s} + q^{-5-5s}}{(1 - q^{1-2s})(1 - q^{2-3s})}$$

where

$$u(T) = \begin{cases} T^3 + T^2 - T - 1 - T^{-1} & \text{if } G(\mathcal{O}) = \mathrm{SL}_3(\mathcal{O}) \\ -T^3 + T^2 - T + 1 - T^{-1} & \text{if } G(\mathcal{O}) = \mathrm{SU}_3(\mathcal{O}). \end{cases}$$

Remark

The computation for $\mathrm{SL}_3(\mathcal{O})$ and $\mathrm{SU}_3(\mathcal{O})$ is completed as well.

Example 2

Applying the p -adic formalism described by AKOV, and under some mild assumptions on the ramification index $e(K/\mathbb{Q}_p)$, we have the following

Theorem

Suppose $p \neq 3$ and assume that K contains a primitive cube root of unity. Let D be a division algebra of degree 3 with $\mathbf{Z}(D) = 3$. Then

$$\zeta_{SL_1^1(D)}(s) = q^3 \frac{1 + q^{-s+1}(1 - q^{-3} - q^{-3s-3})}{1 - q^{-3s+2}},$$

and

$$\zeta_{SL_1(D)}(s) = \frac{(1 + q + q^2)(1 - q^{-3s}) + 9 \left(\frac{1+q+q^2}{3} \right)^{-s} (q^{-s+1} + 1)(q - 1)}{1 - q^{-3s+2}}.$$

What About $\ell > 3$?

- A key ingredient to the computation of the zeta function for the groups under consideration is an understanding of the conjugation action of $G(\mathcal{O})$ on an associated lattice in its Lie-algebra.

What About $\ell > 3$?

- A key ingredient to the computation of the zeta function for the groups under consideration is an understanding of the conjugation action of $G(\mathcal{O})$ on an associated lattice in its Lie-algebra.
- The case of $\mathrm{SL}_2(\mathcal{O})$ is relatively easy.

What About $\ell > 3$?

- A key ingredient to the computation of the zeta function for the groups under consideration is an understanding of the conjugation action of $G(\mathcal{O})$ on an associated lattice in its Lie-algebra.
- The case of $\mathrm{SL}_2(\mathcal{O})$ is relatively easy.
- The case of $\mathrm{SL}_3(\mathcal{O})$ and $\mathrm{SU}_3(\mathcal{O})$ is significantly more involved and requires distinguishing between 8-10 types of conjugacy classes. The level of difficulty for $\ell > 3$ is still unclear.

What About $\ell > 3$?

- A key ingredient to the computation of the zeta function for the groups under consideration is an understanding of the conjugation action of $G(\mathcal{O})$ on an associated lattice in its Lie-algebra.
- The case of $\mathrm{SL}_2(\mathcal{O})$ is relatively easy.
- The case of $\mathrm{SL}_3(\mathcal{O})$ and $\mathrm{SU}_3(\mathcal{O})$ is significantly more involved and requires distinguishing between 8-10 types of conjugacy classes. The level of difficulty for $\ell > 3$ is still unclear.
- The case of $\mathrm{SL}_1(D)$ for division algebras of prime degree is, however, accessible.

Let ℓ be an arbitrary prime.

Assuming $p \neq \ell$ and some additional assumptions on $e(K/\mathbb{Q}_p)$, we have the following

Theorem

Suppose K contains a primitive ℓ -th root of unity, and let D be a division algebra of degree ℓ with $\mathbf{Z}(D) = K$. Then

$$\zeta_{\mathrm{SL}_1^1(D)}(s) = \frac{1 - q^{\binom{\ell}{2}s} + (q^\ell - 1) \cdot S_\ell(q^{1 - \frac{\ell-1}{2}s})}{1 - q^{-\binom{\ell}{2}s + \ell - 1}},$$

where $S_\ell(T) := T + \dots + T^{\ell-1}$.

By an application of tools from Clifford theory, we also obtain the following.

Theorem

Let K be as above, with a primitive ℓ -th root of unity, and let D be a division algebra of degree ℓ with $\mathbf{Z}(D) = K$. Then

$$\zeta_{\mathrm{SL}_1(D)}(s) = \frac{(1 - q^{-\binom{\ell}{2}s}) \cdot \frac{q^\ell - 1}{q - 1} + \left(\frac{q^\ell - 1}{q - 1}\right)^{-s-1} \ell^{s+2} \cdot (q^\ell - 1) \cdot S_\ell(q^{1 - \frac{\ell-1}{2}s})}{1 - q^{-\binom{\ell}{2}s + \ell - 1}},$$

where $S_\ell(T) = T + \dots + T^{\ell-1}$.

By an application of tools from Clifford theory, we also obtain the following.

Theorem

Let K be as above, with a primitive ℓ -th root of unity, and let D be a division algebra of degree ℓ with $\mathbf{Z}(D) = K$. Then

$$\zeta_{\mathrm{SL}_1(D)}(s) = \frac{(1 - q^{-\binom{\ell}{2}s}) \cdot \frac{q^\ell - 1}{q - 1} + \left(\frac{q^\ell - 1}{q - 1}\right)^{-s-1} \ell^{s+2} \cdot (q^\ell - 1) \cdot S_\ell(q^{1 - \frac{\ell-1}{2}s})}{1 - q^{-\binom{\ell}{2}s + \ell - 1}},$$

where $S_\ell(T) = T + \dots + T^{\ell-1}$.

Remark

It is possible that the assumption regarding the existence of primitive roots of unity can be omitted.

Questions ?

Kirillov Orbit Method

Prototype Case

- Consider the group $G = 1 + p\mathrm{Mat}_d(\mathbb{Z}_p)$ and the \mathbb{Z}_p -Lie algebra $\mathfrak{g} = \mathrm{Mat}_d(p\mathbb{Z}_p)$.

Kirillov Orbit Method

Prototype Case

- Consider the group $G = 1 + p\text{Mat}_d(\mathbb{Z}_p)$ and the \mathbb{Z}_p -Lie algebra $\mathfrak{g} = \text{Mat}_d(p\mathbb{Z}_p)$.
- The exponential series $X \mapsto \sum_{j=0}^{\infty} \frac{1}{j!} X^j$ is a bijection of \mathfrak{g} onto G . The map also induces a correspondence between sub-Lie algebras (resp. ideals) of \mathfrak{g} and subgroups (resp. normal subgroups) of G .

Kirillov Orbit Method

Prototype Case

- Consider the group $G = 1 + p\text{Mat}_d(\mathbb{Z}_p)$ and the \mathbb{Z}_p -Lie algebra $\mathfrak{g} = \text{Mat}_d(p\mathbb{Z}_p)$.
- The exponential series $X \mapsto \sum_{j=0}^{\infty} \frac{1}{j!} X^j$ is a bijection of \mathfrak{g} onto G . The map also induces a correspondence between sub-Lie algebras (resp. ideals) of \mathfrak{g} and subgroups (resp. normal subgroups) of G .
- The conjugation action of G on \mathfrak{g} induces an action on the Pontryagin dual $\hat{\mathfrak{g}}$ of \mathfrak{g} .
- To any G -orbit $\Omega \subseteq \hat{\mathfrak{g}}$ we associate the class function

$$\chi_{\Omega}(\exp(X)) = \frac{1}{\sqrt{|\Omega|}} \sum_{f \in \Omega} f(X).$$

- $\Omega \leftrightarrow \chi_{\Omega}$ is a bijection between the orbit space $\hat{\mathfrak{g}}/G$ and the set of finite-dimensional continuous irreducible characters of G .

Kirillov Orbit Method

- An application of Kirillov orbit method allows us to "linearize" the question of analysing the representation theory of a group.

Kirillov Orbit Method

- An application of Kirillov orbit method allows us to "linearize" the question of analysing the representation theory of a group.
- In a more general setting, suppose G is a pro- p group and \mathfrak{g} is a \mathbb{Z}_p -Lie algebra with a homeomorphism $\exp : \mathfrak{g} \rightarrow G$.
- Under assumptions on G guaranteeing the existence of a Lie-correspondence, as well as some other technical assumption, we have the following:

Kirillov Orbit Method

- An application of Kirillov orbit method allows us to "linearize" the question of analysing the representation theory of a group.
- In a more general setting, suppose G is a pro- p group and \mathfrak{g} is a \mathbb{Z}_p -Lie algebra with a homeomorphism $\exp : \mathfrak{g} \rightarrow G$.
- Under assumptions on G guaranteeing the existence of a Lie-correspondence, as well as some other technical assumption, we have the following:

Theorem (Howe)

The map $\Omega \mapsto \chi_\Omega$, where

$$\chi_\Omega((\exp(X))) = \frac{1}{\sqrt{|\Omega|}} \sum_{f \in \Omega} f(X),$$

is a bijection of $\hat{\mathfrak{g}}/G$ onto the set of irreducible continuous finite-dimensional characters of g .

Division Algebras

Let D be a division algebra of degree ℓ , with $K = \mathbf{Z}(D)$. Let L denote the unramified extension of K of degree L .

- It is known that L embeds into D as a maximal subfield of D . In particular L splits D , that is-

$$D \otimes_K L \cong \text{Mat}_\ell(L).$$

Division Algebras

Let D be a division algebra of degree ℓ , with $K = \mathbf{Z}(D)$. Let L denote the unramified extension of K of degree L .

- It is known that L embeds into D as a maximal subfield of D . In particular L splits D , that is-

$$D \otimes_K L \cong \text{Mat}_\ell(L).$$

- The map $x \mapsto x \otimes 1$ embeds D as a sub- K -algebra of $\text{Mat}_\ell(L)$ endowing D with a reduced norm $\text{Nrd}_{D/K}(\cdot)$ and trace $\text{Trd}_{D/K}(\cdot)$.

Division Algebras

Let D be a division algebra of degree ℓ , with $K = \mathbf{Z}(D)$. Let L denote the unramified extension of K of degree L .

- It is known that L embeds into D as a maximal subfield of D . In particular L splits D , that is-

$$D \otimes_K L \cong \text{Mat}_\ell(L).$$

- The map $x \mapsto x \otimes 1$ embeds D as a sub- K -algebra of $\text{Mat}_\ell(L)$ endowing D with a reduced norm $\text{Nrd}_{D/K}(\cdot)$ and trace $\text{Trd}_{D/K}(\cdot)$.
- The absolute value given on K extends uniquely to D and satisfies

$$|\mathbf{x}|_\wp = |\text{Nrd}_{D/K}(\mathbf{x})|_\wp^{1/\ell}, \quad \forall \mathbf{x} \in D.$$

- Put $\mathcal{O} = \{\mathbf{x} \mid |\mathbf{x}|_\wp \leq 1\}$ and $\mathcal{P} = \{\mathbf{x} \mid |\mathbf{x}|_\wp < 1\}$.

- The group $SL_1(D)$ consists of the element of norm 1 in D . It is a pro-finite compact p -adic analytic group.

- The group $SL_1(D)$ consists of the element of norm 1 in D . It is a pro-finite compact p -adic analytic group.
- The group $SL_1^1(D)$ consists of elements $x \in SL_1(D)$ such that $x - 1 \in \mathcal{P}$, and is a maximal pro- p sub group of $SL_1(D)$.
Under some mild assumptions on $e(K/\mathbb{Q}_p)$, the standard exponential map is convergent on the \mathcal{O} -Lie algebra $\mathfrak{sl}_1^1(D)$ of traceless elements in \mathcal{P} , mapping it bijectively onto $SL_1^1(D)$ and making $SL_1^1(D)$ amenable to the Kirrilov orbit method.

- The group $SL_1(D)$ consists of the element of norm 1 in D . It is a pro-finite compact p -adic analytic group.
- The group $SL_1^1(D)$ consists of elements $x \in SL_1(D)$ such that $x - 1 \in \mathcal{P}$, and is a maximal pro- p sub group of $SL_1(D)$.
Under some mild assumptions on $e(K/\mathbb{Q}_p)$, the standard exponential map is convergent on the \mathcal{O} -Lie algebra $\mathfrak{sl}_1^1(D)$ of traceless elements in \mathcal{P} , mapping it bijectively onto $SL_1^1(D)$ and making $SL_1^1(D)$ amenable to the Kirrilov orbit method.
- The quotient $SL_1(D)/SL_1^1(D)$ is embeddable into a subgroup of the cyclic group \mathbb{F}_{q^ℓ} .

Denote $H = \mathrm{SL}_1^1(D)$ and $\mathfrak{h} = \mathfrak{sl}_1^1(D)$.

$$\begin{aligned}
 \zeta_{\mathrm{SL}_1^1(D)}(s) &= \sum_{\Omega \in \widehat{\mathfrak{h}}/H} \chi_{\Omega}(1)^{-s} = \sum_{\Omega \in \widehat{\mathfrak{h}}/H} |\Omega|^{-s/2} \\
 &= \sum_{\omega \in \widehat{\mathfrak{h}}} |H \cdot \omega|^{-\frac{s}{2}-1} = \sum_{\omega \in \widehat{\mathfrak{h}}} |H : \mathbf{C}_H(\omega)|^{-\frac{s}{2}-1} \\
 &= 1 + \sum_{n=1}^{\infty} \sum_{\omega \in \widehat{\mathfrak{h}}/\pi^n \mathfrak{h}} |H : \mathbf{C}_H(\omega)|^{-\frac{s}{2}-1}
 \end{aligned}$$

- **Fact 1:** For any $n \in \mathbb{N}$, there exists a surjection of the set $\pi^{-1}\mathfrak{g} \setminus \mathfrak{g}$ onto $\widehat{\mathfrak{h}/\pi^n\mathfrak{h}}$, whose fibres are cosets of $\pi^{n-1}\mathfrak{g}$ (where \mathfrak{g} is the Lie-algebra of traceless elements in \mathcal{O}).

- **Fact 1:** For any $n \in \mathbb{N}$, there exists a surjection of the set $\pi^{-1}\mathfrak{g} \setminus \mathfrak{g}$ onto $\widehat{\mathfrak{h}/\pi^n\mathfrak{h}}$, whose fibres are cosets of $\pi^{n-1}\mathfrak{g}$ (where \mathfrak{g} is the Lie-algebra of traceless elements in \mathcal{O}).
- **Fact 2** The stabilizer of an element $\omega \in \widehat{\mathfrak{h}/\pi^n\mathfrak{h}}$ equals the set

$$\mathbf{St}^n(\mathbf{x}_\omega) := \{h \in H \mid \mathbf{x}_\omega - h^{-1}\mathbf{x}_\omega h \in \pi^{n-1}\mathfrak{g}\}$$

where \mathbf{x}_ω is any element which is mapped onto ω by the above bijection.

- **Philosophically:** *To understand the conjugation action of subgroups of D^\times on D , one must have an understanding of the subfields of D .*

- **Philosophically:** *To understand the conjugation action of subgroups of D^\times on D , one must have an understanding of the subfields of D .*
- Define

$$\mu(\mathbf{x}) := \ell \cdot \max \{ \text{val}(\mathbf{x} - \lambda) \mid \lambda \in K \} \in \mathbb{Z} \cup \{\infty\}, \quad \forall \mathbf{x} \in D,$$

where $\text{val}(\mathbf{x}) = -\log_q |\mathbf{x}|_\wp \in \frac{1}{\ell}\mathbb{Z}$ is the exponential valuation on D .

- **Philosophically:** *To understand the conjugation action of subgroups of D^\times on D , one must have an understanding of the subfields of D .*
- Define

$$\mu(\mathbf{x}) := \ell \cdot \max \{ \text{val}(\mathbf{x} - \lambda) \mid \lambda \in K \} \in \mathbb{Z} \cup \{\infty\}, \quad \forall \mathbf{x} \in D,$$

where $\text{val}(\mathbf{x}) = -\log_q |\mathbf{x}|_\wp \in \frac{1}{\ell}\mathbb{Z}$ is the exponential valuation on D .

Theorem

Let $\mathbf{x} \in \mathfrak{h}$ and $n \in \mathbb{N}$. Then

$$|\text{SL}_1^1(D) : \mathbf{St}^n(\mathbf{x})| = \begin{cases} q^{(\ell-1)(\ell n - \mu(\mathbf{x}))} & \text{if } \mu(\mathbf{x}) \in \ell\mathbb{Z}, \\ q^{(\ell-1)(\ell n - \mu(\mathbf{x}) - 1)} & \text{if } \mu(\mathbf{x}) \notin \ell\mathbb{Z}, \end{cases}$$

- From this, the computation of the representation zeta function of $\text{SL}_1^1(D)$ follows.

References



[Helmut Koch \(1990\)](#)

Eisenstein Polynomial Sequences and Conjugacy Classes of Division Algebras over Local Fields

Math. Nachr. 147, 307-336



[Richard S. Pierce \(1982\)](#)

Associative Algebras

Springer-Verlag New York, Heidelberg, Berlin



[Andrei Jaikin-Zapirin \(2006\)](#)

Zeta Function of Representation of Compact p -adic Analytic Groups

J. Amer. Math. Soc 19 91-118

Springer-Verlag New York, Heidelberg, Berlin



[Nir Avni, Benjamin Klopsch, Uri Onn and Chrisopher Voll \(2012\)](#)

Representation Zeta Functions of Compact p -adic Analytic Groups and Arithmetic Groups

J. Amer. Math. Soc 19(1) 91-118



[Jon Gonzáles-Sánchez \(2007\)](#)

Kirillov's Orbit Method for p -Groups and Pro- p Groups

Com. in Algebra 37(12) 4776-4488

Thank You