

# Algebraic Geometry 2

## Tutorial session 7

Lecturer: Rami Aizenbud  
TA: Shai Shechter

June 5, 2020

# Recollection – gluing two schemes along open subsets

Let  $(X_1, \mathcal{O}_1)$  be schemes and  $U_i \subseteq X_i$  open subsets.

# Recollection – gluing two schemes along open subsets

Let  $(X_1, \mathcal{O}_1)$  be schemes and  $U_i \subseteq X_i$  open subsets. Assume given an isomorphism  $\varphi : (U_1, \mathcal{O}_1|_{U_1}) \rightarrow (U_2, \mathcal{O}_2|_{U_2})$  of locally ringed spaces.

# Recollection – gluing two schemes along open subsets

Let  $(X_1, \mathcal{O}_1)$  be schemes and  $U_i \subseteq X_i$  open subsets. Assume given an isomorphism  $\varphi : (U_1, \mathcal{O}_1|_{U_1}) \rightarrow (U_2, \mathcal{O}_2|_{U_2})$  of locally ringed spaces.

## Definition

The gluing of  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$  via  $\varphi$  is  $(X, \mathcal{O}_X)$ , where:

# Recollection – gluing two schemes along open subsets

Let  $(X_1, \mathcal{O}_1)$  be schemes and  $U_i \subseteq X_i$  open subsets. Assume given an isomorphism  $\varphi : (U_1, \mathcal{O}_1|_{U_1}) \rightarrow (U_2, \mathcal{O}_2|_{U_2})$  of locally ringed spaces.

## Definition

The gluing of  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$  via  $\varphi$  is  $(X, \mathcal{O}_X)$ , where:

- ①  $|X| = (|X_1| \sqcup |X_2|) / \sim$  where  $\sim$  is the equivalence relation generated by  $\{(x, \varphi(x)) : x \in U_1\}$ .  $V \subseteq |X|$  is open iff  $V \cap |X_1|$  and  $V \cap |X_2|$  are both open.

# Recollection – gluing two schemes along open subsets

Let  $(X_1, \mathcal{O}_1)$  be schemes and  $U_i \subseteq X_i$  open subsets. Assume given an isomorphism  $\varphi : (U_1, \mathcal{O}_1|_{U_1}) \rightarrow (U_2, \mathcal{O}_2|_{U_2})$  of locally ringed spaces.

## Definition

The gluing of  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$  via  $\varphi$  is  $(X, \mathcal{O}_X)$ , where:

- ①  $|X| = (|X_1| \sqcup |X_2|) / \sim$  where  $\sim$  is the equivalence relation generated by  $\{(x, \varphi(x)) : x \in U_1\}$ .  $V \subseteq |X|$  is open iff  $V \cap |X_1|$  and  $V \cap |X_2|$  are both open.
- ② Given  $V \subseteq X$  open write  $V_i = V \cap |X_i|$  ( $i = 1, 2$ ) and set

$$\mathcal{O}_X(V)$$

$$= \{(s_1, s_2) \in \mathcal{O}_{X_1}(V_1) \times \mathcal{O}_{X_2}(V_2) : s_1|_{V_1 \cap U_1} = \varphi^\sharp(s_2|_{V_2 \cap U_2})\}.$$

## Remark

We slightly abused notation in the previous slide, by identifying  $|X_i|$  with a subspace of  $|X|$ . This is possible, because both are naturally embedded in  $|X|$ .

In Hartshorne, Example 2.3.5, the author is more righteous, and uses explicit maps  $i_j : X_j \rightarrow X$ .

# An example – $\mathbb{A}_k^2 \setminus \{\text{origin}\}$

## Exercise

Let  $k$  be a field, and, as usual  $\mathbb{A}_k^2 = \text{Spec}(k[x, y])$ . Show that  $\mathbb{A}_k^2 \setminus \{(x, y)\}$  is not affine.

# An example – $\mathbb{A}_k^2 \setminus \{\text{origin}\}$

## Exercise

Let  $k$  be a field, and, as usual  $\mathbb{A}_k^2 = \text{Spec}(k[x, y])$ . Show that  $\mathbb{A}_k^2 \setminus \{(x, y)\}$  is not affine.

Here  $(x, y)$  is the maximal ideal corresponding to the closed point of the origin.

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ .

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism;

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ;

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.
- ② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.
- ② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:
  - $\Gamma(D(x), \mathcal{O}_{D(x)}) = k[x, y, \frac{1}{x}]$  and  $\Gamma(D(y), \mathcal{O}_{D(y)}) = k[x, y, \frac{1}{y}]$ .

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.

② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:

- $\Gamma(D(x), \mathcal{O}_{D(x)}) = k[x, y, \frac{1}{x}]$  and  $\Gamma(D(y), \mathcal{O}_{D(y)}) = k[x, y, \frac{1}{y}]$ .
- $(s_1, s_2)$  is a global section on  $D(x) \cup D(y)$  iff  $s_1$  and  $s_2$  coincide elements of  $\mathcal{O}_{D(x)}(D(xy)) = \mathcal{O}_{D(y)}(D(xy)) = k[x, y, \frac{1}{xy}]$ .

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.
- ② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:
  - $\Gamma(D(x), \mathcal{O}_{D(x)}) = k[x, y, \frac{1}{x}]$  and  $\Gamma(D(y), \mathcal{O}_{D(y)}) = k[x, y, \frac{1}{y}]$ .
  - $(s_1, s_2)$  is a global section on  $D(x) \cup D(y)$  iff  $s_1$  and  $s_2$  coincide elements of  $\mathcal{O}_{D(x)}(D(xy)) = \mathcal{O}_{D(y)}(D(xy)) = k[x, y, \frac{1}{xy}]$ .
  - But  $s_1|_{D(xy)} = s_2|_{D(xy)}$  implies  $s_1, s_2 \in k[x, y]$ .

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.
- ② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:
  - $\Gamma(D(x), \mathcal{O}_{D(x)}) = k[x, y, \frac{1}{x}]$  and  $\Gamma(D(y), \mathcal{O}_{D(y)}) = k[x, y, \frac{1}{y}]$ .
  - $(s_1, s_2)$  is a global section on  $D(x) \cup D(y)$  iff  $s_1$  and  $s_2$  coincide elements of  $\mathcal{O}_{D(x)}(D(xy)) = \mathcal{O}_{D(y)}(D(xy)) = k[x, y, \frac{1}{xy}]$ .
  - But  $s_1|_{D(xy)} = s_2|_{D(xy)}$  implies  $s_1, s_2 \in k[x, y]$ .
  - Thus  $\Gamma(Z, \mathcal{O}_Z) = k[x, y]$  and, if  $Z$  is affine, then  $Z = \mathbb{A}_k^2$ .

## Solution.

Put  $Z = \mathbb{A}_k^2 \setminus \{(x, y)\}$ .

- ①  $Z = D(x) \cup D(y)$ , a union of principal open sets, with  $D(x) \cap D(y) = D(xy)$ . By the sheaf axioms,  $Z$  is isomorphic to the scheme obtained by gluing  $D(x)$  and  $D(y)$  along  $D(xy)$  via the identity isomorphism; the map  $Z \rightarrow D(x) \cup_{D(xy)} D(y)$  sends an element  $s \in \mathcal{O}_Z(V)$  to  $(s|_{V \cap D(x)}, s|_{V \cap D(y)})$ ; identity implies injective, locality implies surjective.
- ② Now compute  $\Gamma(Z, \mathcal{O}_Z)$  via the gluing:
  - $\Gamma(D(x), \mathcal{O}_{D(x)}) = k[x, y, \frac{1}{x}]$  and  $\Gamma(D(y), \mathcal{O}_{D(y)}) = k[x, y, \frac{1}{y}]$ .
  - $(s_1, s_2)$  is a global section on  $D(x) \cup D(y)$  iff  $s_1$  and  $s_2$  coincide elements of  $\mathcal{O}_{D(x)}(D(xy)) = \mathcal{O}_{D(y)}(D(xy)) = k[x, y, \frac{1}{xy}]$ .
  - But  $s_1|_{D(xy)} = s_2|_{D(xy)}$  implies  $s_1, s_2 \in k[x, y]$ .
  - Thus  $\Gamma(Z, \mathcal{O}_Z) = k[x, y]$  and, if  $Z$  is affine, then  $Z = \mathbb{A}_k^2$ .
- ③ However,  $(x)$  and  $(y)$  intersect non-trivially as elements of  $\mathbb{A}_k^2$ , and not of  $Z$ . Thus  $Z$  is not affine.



# An aside

## Corollary

*Any regular function on  $\mathbb{A}_k^2 \setminus \{(x, y)\}$  can be extended uniquely to a regular function on  $\mathbb{A}_k^2$ .*

This is, in fact, true without the assumption that  $k$  is a field.

# An aside

## Corollary

*Any regular function on  $\mathbb{A}_k^2 \setminus \{(x, y)\}$  can be extended uniquely to a regular function on  $\mathbb{A}_k^2$ .*

This is, in fact, true without the assumption that  $k$  is a field.

## More examples

Let  $A$  be a ring (or a field, if more convenient).

## More examples

Let  $A$  be a ring (or a field, if more convenient). Put

- ①  $X_1 = X_2 = \text{Spec}(k[t]) = \mathbb{A}_k^1$ ; and
- ②  $U_1 = U_2 = \text{Spec}(k[t, t^{-1}]) = \mathbb{A}_k^1 \setminus \{(t)\}$ .

## More examples

Let  $A$  be a ring (or a field, if more convenient). Put

- ①  $X_1 = X_2 = \text{Spec}(k[t]) = \mathbb{A}_k^1$ ; and
- ②  $U_1 = U_2 = \text{Spec}(k[t, t^{-1}]) = \mathbb{A}_k^1 \setminus \{(t)\}$ .

Upon specifying an isomorphism  $\varphi : U_1 \rightarrow U_2$ , we can glue  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$ .

## More examples

Let  $A$  be a ring (or a field, if more convenient). Put

- ①  $X_1 = X_2 = \text{Spec}(k[t]) = \mathbb{A}_k^1$ ; and
- ②  $U_1 = U_2 = \text{Spec}(k[t, t^{-1}]) = \mathbb{A}_k^1 \setminus \{(t)\}$ .

Upon specifying an isomorphism  $\varphi : U_1 \rightarrow U_2$ , we can glue  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$ .

- ① If  $\varphi$  is the identity map, then the resulting scheme is the affine line with doubled origin, which we've seen before.

\_\_\_\_\_ : \_\_\_\_\_

# More examples

Let  $A$  be a ring (or a field, if more convenient). Put

- ①  $X_1 = X_2 = \text{Spec}(k[t]) = \mathbb{A}_k^1$ ; and
- ②  $U_1 = U_2 = \text{Spec}(k[t, t^{-1}]) = \mathbb{A}_k^1 \setminus \{(t)\}$ .

Upon specifying an isomorphism  $\varphi : U_1 \rightarrow U_2$ , we can glue  $X_1$  and  $X_2$  along  $U_1$  and  $U_2$ .

- ① If  $\varphi$  is the identity map, then the resulting scheme is the affine line with doubled origin, which we've seen before.

\_\_\_\_\_ : \_\_\_\_\_

- ② We can also take  $\varphi$  to be the isomorphism induced from  $t \mapsto t^{-1}$  on  $U_1$ . The resulting space is denoted  $\mathbb{P}_A^1$ .

# Regarding $\mathbb{P}^1_k$

Assume now that  $A = k$  is an algebraically closed field.

# Regarding $\mathbb{P}_k^1$

Assume now that  $A = k$  is an algebraically closed field.


Consider “non-zero” closed points on  $\mathbb{A}_k^1$  which correspond to ideals  $(t - a)$  where  $a \neq 0$ . The morphism  $\varphi$  maps an ideal of the form  $(bt - a)$  to  $(bt^{-1} - a) = (at - b)$ , and preserves the generic point  $(0)$ .

# Regarding $\mathbb{P}_k^1$

Assume now that  $A = k$  is an algebraically closed field.

Consider “non-zero” closed points on  $\mathbb{A}_k^1$  which correspond to ideals  $(t - a)$  where  $a \neq 0$ . The morphism  $\varphi$  maps an ideal of the form  $(bt - a)$  to  $(bt^{-1} - a) = (at - b)$ , and preserves the generic point  $(0)$ .

The space  $\mathbb{P}_k^1$  can be thus identified with the space of pairs  $[a : b]$  with  $a, b \in k$  not both zero, upto multiplication of both  $a$  and  $b$  by a constant  $c \in k^\times$ . The identification is such that, if  $a \neq 0$  then  $[a : b]$  is mapped to the ideal  $(bt - a)$  on one affine line, and if  $b \neq 0$ , it is mapped to the ideal  $(at - b)$  on the second affine line.



## Exercise

Show that  $\mathbb{P}_k^1$  is not affine, for  $k$  a field.

## Exercise

Show that  $\mathbb{P}_k^1$  is not affine, for  $k$  a field.

## Solution.

Usual trick- compute global sections.

## Exercise

Show that  $\mathbb{P}_k^1$  is not affine, for  $k$  a field.

## Solution.

Usual trick- compute global sections. Let  $f \in \Gamma(\mathbb{P}_k^1, \mathcal{O}_{\mathbb{P}_k^1})$  be a global section, and considering  $f_1 = f|_{X_1}$  and  $f_2 = f|_{X_2}$ . Then  $f_1$  and  $f_2$  are global sections on subschemes which are isomorphic to  $\mathbb{A}_k^1$ , and therefore are given by polynomials. Furthermore, by considering their restrictions to  $U_1$  and  $U_2$  and taking into account the action of  $\varphi$  on them, we see that  $f_1(t) = f_2(1/t)$ .

## Exercise

Show that  $\mathbb{P}_k^1$  is not affine, for  $k$  a field.

## Solution.

Usual trick- compute global sections. Let  $f \in \Gamma(\mathbb{P}_k^1, \mathcal{O}_{\mathbb{P}_k^1})$  be a global section, and considering  $f_1 = f|_{X_1}$  and  $f_2 = f|_{X_2}$ . Then  $f_1$  and  $f_2$  are global sections on subschemes which are isomorphic to  $\mathbb{A}_k^1$ , and therefore are given by polynomials. Furthermore, by considering their restrictions to  $U_1$  and  $U_2$  and taking into account the action of  $\varphi$  on them, we see that  $f_1(t) = f_2(1/t)$ .

Since both  $f_1$  and  $f_2$  are polynomials, this is only possible if both are constant, thus  $\Gamma(\mathbb{P}_k^1, \mathcal{O}_{\mathbb{P}_k^1}) \simeq k$ . If  $\mathbb{P}_k^1$  were affine, it would have had only one point, which is false. □

# Gluing contd

Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and

## Gluing contd

### Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .

Then,  $\exists$  a scheme  $X$  with morphisms  $\psi_i : X_i \rightarrow X$  such that:

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .

Then,  $\exists$  a scheme  $X$  with morphisms  $\psi_i : X_i \rightarrow X$  such that:

- ①  $\psi_i$  is an isomorphism of  $X_i$  onto an open subscheme;

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .

Then,  $\exists$  a scheme  $X$  with morphisms  $\psi_i : X_i \rightarrow X$  such that:

- ①  $\psi_i$  is an isomorphism of  $X_i$  onto an open subscheme;
- ②  $\bigcup |\psi_i(X_i)| = |X|$ ;

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)

Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .

Then,  $\exists$  a scheme  $X$  with morphisms  $\psi_i : X_i \rightarrow X$  such that:

- ①  $\psi_i$  is an isomorphism of  $X_i$  onto an open subscheme;
- ②  $\bigcup |\psi_i(X_i)| = |X|$ ;
- ③  $\psi_i(U_{i,j}) = \psi_i(X_i) \cap \psi_j(X_j)$ ; and

# Gluing contd

## Lemma (The gluing lemma, HS Ex 2.12)


Let  $\{X_i\}$  be a, possibly infinite, family of schemes. For each  $i \neq j$ , assume given

- ① an open subset  $U_{i,j} \subseteq X_i$  with the induced scheme structure  $\mathcal{O}_{U_{i,j}} = \mathcal{O}_{X_i}|_{U_{i,j}}$ ; and
- ② an isomorphism  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$  such that
  - $\varphi_{j,i} = \varphi_{i,j}^{-1}$ ; and
  - for any  $i, j, k$ ,  $\varphi_{i,j}(U_{i,j} \cap U_{i,k}) = U_{j,i} \cap U_{j,k}$  and  $\varphi_{i,k} = \varphi_{j,k} \circ \varphi_{i,j}$  on  $U_{i,j} \cap U_{i,k}$ .


Then,  $\exists$  a scheme  $X$  with morphisms  $\psi_i : X_i \rightarrow X$  such that:

- ①  $\psi_i$  is an isomorphism of  $X_i$  onto an open subscheme;
- ②  $\bigcup |\psi_i(X_i)| = |X|$ ;
- ③  $\psi_i(U_{i,j}) = \psi_i(X_i) \cap \psi_j(X_j)$ ; and
- ④  $\psi_i = \psi_j \circ \varphi_{i,j}$  on  $U_{i,j}$ .

# Gluing input



# Gluing output



## Remark

The data  $(X_i, U_{i,j}, \varphi_{i,j})$  is often referred to as the **descent data** of the gluing.

# Proof of the gluing lemma

- ① **Topological space.** Let  $\tilde{X} = \bigsqcup |X_i|$ , and let  $\sim$  be the relation defined on  $\tilde{X}$  by

$$x \sim y \iff x = y \text{ or } \exists i, j \text{ st } x \in U_{i,j} \subseteq X_i \text{ and } y = \varphi_{i,j}(x).$$

The assumptions imply that  $\sim$  is an equivalence relation. Let  $|X| = \tilde{X}/\sim$  with the quotient topology. Let  $\psi_i : |X_i| \rightarrow |X|$  be the natural inclusions.

- ② **Structure sheaf** Given  $V \subseteq |X|$  open, put  $V_i = V \cap |X_i|$  and define

$$\mathcal{O}_X(V)$$

$$= \left\{ (s_i)_i \in \prod_i \mathcal{O}_{X_i}(V_i) : s_i|_{(V_i \cap V_j) \cap U_{i,j}} = \varphi_{i,j}^\sharp(s_j|_{(V_i \cap V_j) \cap U_{j,i}}) \right\}.$$

The restriction maps are defined coordinate-wise, i.e.

$$\text{res}_{V,W} = (\text{res}_{V_i,W_i})_i \text{ where } W_i = W \cap |X_i|.$$

- ② **Structure sheaf** Given  $V \subseteq |X|$  open, put  $V_i = V \cap |X_i|$  and define

$$\mathcal{O}_X(V)$$

$$= \left\{ (s_i)_i \in \prod_i \mathcal{O}_{X_i}(V_i) : s_i|_{(V_i \cap V_j) \cap U_{i,j}} = \varphi_{i,j}^\sharp(s_j|_{(V_i \cap V_j) \cap U_{j,i}}) \right\}.$$

The restriction maps are defined coordinate-wise, i.e.  
 $\text{res}_{V,W} = (\text{res}_{V_i,W_i})_i$  where  $W_i = W \cap |X_i|$ .

We need to verify

- ② **Structure sheaf** Given  $V \subseteq |X|$  open, put  $V_i = V \cap |X_i|$  and define

$$\mathcal{O}_X(V)$$

$$= \left\{ (s_i)_i \in \prod_i \mathcal{O}_{X_i}(V_i) : s_i|_{(V_i \cap V_j) \cap U_{i,j}} = \varphi_{i,j}^\sharp(s_j|_{(V_i \cap V_j) \cap U_{j,i}}) \right\}.$$

The restriction maps are defined coordinate-wise, i.e.  
 $\text{res}_{V,W} = (\text{res}_{V_i,W_i})_i$  where  $W_i = W \cap |X_i|$ .

We need to verify

- ①  $\mathcal{O}_X$  is a sheaf;

- ② **Structure sheaf** Given  $V \subseteq |X|$  open, put  $V_i = V \cap |X_i|$  and define

$$\mathcal{O}_X(V)$$

$$= \left\{ (s_i)_i \in \prod_i \mathcal{O}_{X_i}(V_i) : s_i|_{(V_i \cap V_j) \cap U_{i,j}} = \varphi_{i,j}^\sharp(s_j|_{(V_i \cap V_j) \cap U_{j,i}}) \right\}.$$

The restriction maps are defined coordinate-wise, i.e.  
 $\text{res}_{V,W} = (\text{res}_{V_i,W_i})_i$  where  $W_i = W \cap |X_i|$ .

We need to verify

- ①  $\mathcal{O}_X$  is a sheaf;
- ②  $(X, \mathcal{O}_X)$  is a locally ringed space; and

- ② **Structure sheaf** Given  $V \subseteq |X|$  open, put  $V_i = V \cap |X_i|$  and define

$$\mathcal{O}_X(V)$$

$$= \left\{ (s_i)_i \in \prod_i \mathcal{O}_{X_i}(V_i) : s_i|_{(V_i \cap V_j) \cap U_{i,j}} = \varphi_{i,j}^\sharp(s_j|_{(V_i \cap V_j) \cap U_{j,i}}) \right\}.$$

The restriction maps are defined coordinate-wise, i.e.  
 $\text{res}_{V,W} = (\text{res}_{V_i,W_i})_i$  where  $W_i = W \cap |X_i|$ .

We need to verify

- ①  $\mathcal{O}_X$  is a sheaf;
- ②  $(X, \mathcal{O}_X)$  is a locally ringed space; and
- ③  $(X, \mathcal{O}_X)$  is covered by affine schemes.

- ① The fact that  $\mathcal{O}_X$  is a presheaf and that it satisfies the sheaf axioms are verified for each coordinate separately, using the fact that any open cover of  $X$  gives rise to an open cover of the  $X_i$ 's.

- ① The fact that  $\mathcal{O}_X$  is a presheaf and that it satisfies the sheaf axioms are verified for each coordinate separately, using the fact that any open cover of  $X$  gives rise to an open cover of the  $X_i$ 's.
- ② It is clear that  $\mathcal{O}_X$  is a sheaf of rings. To verify that all stalks are local rings, we first consider the stalk of  $\prod_i \mathcal{O}_{X_i}$  at a point  $x \in X$ . This stalk is given by

$$\prod_i A_i \quad \text{where} \quad A_i = \begin{cases} \mathcal{O}_{X_i, x} & \text{if } x \in X_i \\ 0 & \text{otherwise.} \end{cases}$$

- ① The fact that  $\mathcal{O}_X$  is a presheaf and that it satisfies the sheaf axioms are verified for each coordinate separately, using the fact that any open cover of  $X$  gives rise to an open cover of the  $X_i$ 's.
- ② It is clear that  $\mathcal{O}_X$  is a sheaf of rings. To verify that all stalks are local rings, we first consider the stalk of  $\prod_i \mathcal{O}_{X_i}$  at a point  $x \in X$ . This stalk is given by

$$\prod_i A_i \quad \text{where} \quad A_i = \begin{cases} \mathcal{O}_{X_i, x} & \text{if } x \in X_i \\ 0 & \text{otherwise.} \end{cases}$$

Furthermore, if  $x$  is contained in  $X_i \cap X_j = U_{i,j}$  then the morphism  $\varphi_{i,j}$  sets up an isomorphism between the stalks.

- The fact that  $\mathcal{O}_X$  is a presheaf and that it satisfies the sheaf axioms are verified for each coordinate separately, using the fact that any open cover of  $X$  gives rise to an open cover of the  $X_i$ 's.
- It is clear that  $\mathcal{O}_X$  is a sheaf of rings. To verify that all stalks are local rings, we first consider the stalk of  $\prod_i \mathcal{O}_{X_i}$  at a point  $x \in X$ . This stalk is given by

$$\prod_i A_i \quad \text{where} \quad A_i = \begin{cases} \mathcal{O}_{X_i, x} & \text{if } x \in X_i \\ 0 & \text{otherwise.} \end{cases}$$

Furthermore, if  $x$  is contained in  $X_i \cap X_j = U_{i,j}$  then the morphism  $\varphi_{i,j}$  sets up an isomorphism between the stalks. Finally, the definition of  $\mathcal{O}_X$  gives that  $\mathcal{O}_{X,x}$  is embedded *diagonally* in the stalk above, and isomorphic to any of its components.

- The fact that  $\mathcal{O}_X$  is a presheaf and that it satisfies the sheaf axioms are verified for each coordinate separately, using the fact that any open cover of  $X$  gives rise to an open cover of the  $X_i$ 's.
- It is clear that  $\mathcal{O}_X$  is a sheaf of rings. To verify that all stalks are local rings, we first consider the stalk of  $\prod_i \mathcal{O}_{X_i}$  at a point  $x \in X$ . This stalk is given by

$$\prod_i A_i \quad \text{where} \quad A_i = \begin{cases} \mathcal{O}_{X_i, x} & \text{if } x \in X_i \\ 0 & \text{otherwise.} \end{cases}$$

Furthermore, if  $x$  is contained in  $X_i \cap X_j = U_{i,j}$  then the morphism  $\varphi_{i,j}$  sets up an isomorphism between the stalks. Finally, the definition of  $\mathcal{O}_X$  gives that  $\mathcal{O}_{X,x}$  is embedded *diagonally* in the stalk above, and isomorphic to any of its components.

- $X$  is covered by the  $X_i$ , and these are covered by affine schemes.

# The projective $n$ -space $\mathbb{P}^n$

We now construct the projective  $n$ -space as the gluing of  $n + 1$  affine  $n$ -spaces.

# The projective $n$ -space $\mathbb{P}^n$

We now construct the projective  $n$ -space as the gluing of  $n + 1$  affine  $n$ -spaces. We'll work over a field  $k$ , even though the construction holds for general rings.

For each  $i = 0, \dots, n$ , we let  $X_i$  be given by

$$\text{Spec}(k[x_0^i, \dots, x_n^i]/(x_i^i - 1)) \simeq \mathbb{A}_k^n.$$

Note that the variable  $x_i^i$  is a “dummy”, and we can think of  $X_i$  has having coordinates  $x_0^i, \dots, x_{i-1}^i, x_{i+1}^i, \dots, x_n^i$ .

# The projective $n$ -space $\mathbb{P}^n$

We now construct the projective  $n$ -space as the gluing of  $n + 1$  affine  $n$ -spaces. We'll work over a field  $k$ , even though the construction holds for general rings.

For each  $i = 0, \dots, n$ , we let  $X_i$  be given by

$$\text{Spec}(k[x_0^i, \dots, x_n^i]/(x_i^i - 1)) \simeq \mathbb{A}_k^n.$$

Note that the variable  $x_i^i$  is a “dummy”, and we can think of  $X_i$  has having coordinates  $x_0^i, \dots, x_{i-1}^i, x_{i+1}^i, \dots, x_n^i$ .

Given  $j = 0, \dots, n$ , we put  $U_{i,j} = D_{X_i}(x_j^i)$ . We wish to glue the  $X_i$ 's together along the  $U_{i,j}$ 's. To do so, we need to specify the isomorphisms  $\varphi_{i,j} : U_{i,j} \rightarrow U_{j,i}$ .

## Construction of $\varphi_{i,j}$

For  $j \neq i$ , we have  $\mathcal{O}_{X_i}(U_{i,j}) = \text{Spec}(k[x_0^i, \dots, x_n^i, \frac{1}{x_j^i}] / (x_i^i - 1))$ .

## Construction of $\varphi_{i,j}$

For  $j \neq i$ , we have  $\mathcal{O}_{X_i}(U_{i,j}) = \text{Spec}(k[x_0^i, \dots, x_n^i, \frac{1}{x_j^i}] / (x_i^i - 1))$ . The morphism  $\varphi_{i,j}$  is the one associated with the ring homomorphism

$$\begin{aligned} & \frac{k[x_0^j, \dots, x_n^j, \frac{1}{x_i^j}]}{(x_j^j - 1)} \rightarrow \frac{k[x_0^i, \dots, x_n^i, \frac{1}{x_j^i}]}{(x_i^i - 1)} \\ & x_k^j \mapsto \frac{x_k^i}{x_j^i} \quad \text{for } k = 0, \dots, n \end{aligned}$$

## Construction of $\varphi_{i,j}$

For  $j \neq i$ , we have  $\mathcal{O}_{X_i}(U_{i,j}) = \text{Spec}(k[x_0^i, \dots, x_n^i, \frac{1}{x_j^i}] / (x_i^i - 1))$ . The morphism  $\varphi_{i,j}$  is the one associated with the ring homomorphism

$$\begin{array}{ccc} k[x_0^j, \dots, x_n^j, \frac{1}{x_i^j}] & \longrightarrow & k[x_0^i, \dots, x_n^i, \frac{1}{x_j^i}] \\ (x_j^j - 1) & \rightarrow & (x_i^i - 1) \\ x_k^j & \mapsto & \frac{x_k^i}{x_j^i} \quad \text{for } k = 0, \dots, n \end{array}$$

## Lemma

The data  $(X_i, U_{i,j}, \varphi_{i,j})$  comprises a descent data, and therefore may be glued to a scheme  $X = \mathbb{P}_k^n$ .

## Proof of lemma

- ① The schemes  $U_i, U_j$  are clearly isomorphic (by change of variables), and that the map  $\varphi_{i,j}$  defined above is invertible, with  $\varphi_{i,j}^{-1} = \varphi_{j,i}$ .

## Proof of lemma

- ① The schemes  $U_i, U_j$  are clearly isomorphic (by change of variables), and that the map  $\varphi_{i,j}$  defined above is invertible, with  $\varphi_{i,j}^{-1} = \varphi_{j,i}$ . Indeed, the composition  $\varphi_{j,i} \circ \varphi_{i,j}$  acts on the ring polynomials by

$$x_k^j \xrightarrow{\varphi_{i,j}^\#} \frac{x_k^i}{x_j^i} \xrightarrow{\varphi_{j,i}^\#} \frac{x_k^j/x_i^j}{x_j^j/x_i^j} = x_k^j/x_j^j = x_k^j$$

as  $x_j^j = 1$  in  $X_j$ .

# Proof of lemma

- ① The schemes  $U_i, U_j$  are clearly isomorphic (by change of variables), and that the map  $\varphi_{i,j}$  defined above is invertible, with  $\varphi_{i,j}^{-1} = \varphi_{j,i}$ . Indeed, the composition  $\varphi_{j,i} \circ \varphi_{i,j}$  acts on the ring polynomials by

$$x_k^j \xrightarrow{\varphi_{i,j}^\#} \frac{x_k^i}{x_j^i} \xrightarrow{\varphi_{j,i}^\#} \frac{x_k^j/x_i^j}{x_j^j/x_i^j} = x_k^j/x_j^j = x_k^j$$

as  $x_j^j = 1$  in  $X_j$ .

- ② To show agreement on triple intersection, we compute the action of  $\varphi_{k,j} \circ \varphi_{i,j}$  on  $x_r^j$ .

$$\begin{array}{ccccc} x_r^k & \xrightarrow{\varphi_{j,k}^\#} & \frac{x_r^j}{x_k^j} & \xrightarrow{\varphi_{i,j}^\#} & \frac{x_r^i/x_j^i}{x_k^i/x_j^i} \\ & \swarrow \varphi_{i,k}^\# & & & \parallel \\ & & \frac{x_r^i}{x_k^i} & & \end{array}$$

Another construction which generalizes the notion of a projective variety appears in Hartshorne, p 76, using the Proj construction.

Another construction which generalizes the notion of a projective variety appears in Hartshorne, p 76, using the Proj construction. Using this construction, we have that

$$\mathbb{P}_k^n = \text{Proj}(k[x_0, \dots, x_n]).$$

The corresponding cover by affine spaces is given by  $X_i = D_+(x_i)$ , with  $U_{i,j} = D_+(x_i x_j)$ . We will recall the definitions and prove this in the next tutorial.

# Questions?