

# Regular Characters of Classical Groups

Forth International Workshop on  
Zeta Functions in Algebra and Geometry

Shai Shechter

Ben Gurion University of the Negev, Beer Sheva

Bielefeld, June 1, 2017

This work is part of the authors Ph.D thesis,  
under the supervision of Uri Onn.

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.
- ▶  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ .

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.
- ▶  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ .
- ▶  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$  the residue field.

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.
- ▶  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ .
- ▶  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$  the residue field.
- ▶  $G$  is a reductive algebraic group, defined over  $\mathfrak{o}$ .

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.
- ▶  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ .
- ▶  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$  the residue field.
- ▶  $\mathbf{G}$  is a reductive algebraic group, defined over  $\mathfrak{o}$ .
- ▶  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ .

# Setting

- ▶  $\mathfrak{o}$  is a complete discrete valuation ring.
- ▶  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ .
- ▶  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$  the residue field.
- ▶  $\mathbf{G}$  is a reductive algebraic group, defined over  $\mathfrak{o}$ .
- ▶  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ .
  
- ▶  $\text{Irr}(\mathbf{G}(\mathfrak{o}))$  is the set of (continuous) irreducible complex valued characters of  $\mathbf{G}(\mathfrak{o})$ .

- Put  $G = \mathbf{G}(\mathfrak{o})$ .



- Put  $G = \mathbf{G}(\mathfrak{o})$ . For any  $m \in \mathbb{N}$ , write  $G^m$  for the principal congruence subgroup  $G^m = \text{Ker}(G \rightarrow \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m))$ .

- Put  $G = \mathbf{G}(\mathfrak{o})$ . For any  $m \in \mathbb{N}$ , write  $G^m$  for the principal congruence subgroup  $G^m = \text{Ker}(G \rightarrow \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m))$ .

## Definition

The **level** of a continuous irreducible representation  $\rho$  of  $G$  is the minimal integer  $\ell$  such that  $\rho(G^{\ell+1}) = \{1\}$ . The level of a character  $\chi$  is the level of the associated representation.

# Quick recap- Regular characters of $GL_N(\mathfrak{o})$

Consider  $G = GL_N(\mathfrak{o})$ .

# Quick recap- Regular characters of $GL_N(\mathfrak{o})$

Consider  $G = GL_N(\mathfrak{o})$ .

The congruence subgroups of  $G$  are  $G^m = 1 + \mathfrak{p}^m M_N(\mathfrak{o})$ .

# Quick recap- Regular characters of $GL_N(\mathfrak{o})$

Consider  $G = GL_N(\mathfrak{o})$ .

The congruence subgroups of  $G$  are  $G^m = 1 + \mathfrak{p}^m M_N(\mathfrak{o})$ .

Given a character  $\chi \in \text{Irr}(G)$ , of level  $\ell > 0$ , identify  $\chi$  with a character of  $G/G^{\ell+1}$  and consider the its restriction to the group  $G^\ell/G^{\ell+1} \simeq M_n(\mathbb{F}_q)$ .

- By Clifford theory, the constituents of  $\chi|_{G^\ell/G^{\ell+1}}$  comprise a complete orbit  $\Omega$  in the Pontryagin dual of  $M_N(\mathbb{F}_q)$  for the coadjoint action of  $GL_N(\mathbb{F}_q)$ , i.e.

$$\chi|_{G^\ell/G^{\ell+1}} = \left( \sum_{\varphi \in \Omega} \varphi \right)^{\oplus e} \quad (e \in \mathbb{N}).$$

- By Clifford theory, the constituents of  $\chi|_{G^\ell/G^{\ell+1}}$  comprise a complete orbit  $\Omega$  in the Pontryagin dual of  $M_N(\mathbb{F}_q)$  for the coadjoint action of  $GL_N(\mathbb{F}_q)$ , i.e.

$$\chi|_{G^\ell/G^{\ell+1}} = \left( \sum_{\varphi \in \Omega} \varphi \right)^{\oplus e} \quad (e \in \mathbb{N}).$$

- By non-degeneracy of

$$(X, Y) \mapsto \text{Tr}(XY) : M_N(\mathbb{F}_q) \times M_N(\mathbb{F}_q) \rightarrow \mathbb{F}_q,$$

the orbit  $\Theta$  is associated uniquely with a similarity class  $\Theta \subseteq M_N(\mathbb{F}_q)$ .

$$\begin{array}{c}
 \chi \in \text{Irr}(G) \text{ of level } \ell > 0 \\
 \downarrow \\
 \Omega \in \text{Ad}^*(\text{GL}_N(\mathbb{F}_q)) \backslash \widehat{\text{M}_N(\mathbb{F}_q)} \\
 \updownarrow \\
 \Theta \in \text{Ad}(\text{GL}_N(\mathbb{F}_q)) \backslash \text{M}_N(\mathbb{F}_q).
 \end{array}$$



$$\begin{array}{c}
 \chi \in \text{Irr}(G) \text{ of level } \ell > 0 \\
 \downarrow \\
 \Omega \in \text{Ad}^*(\text{GL}_N(\mathbb{F}_q)) \backslash \widehat{\text{M}_N(\mathbb{F}_q)} \\
 \updownarrow \\
 \Theta \in \text{Ad}(\text{GL}_N(\mathbb{F}_q)) \backslash \text{M}_N(\mathbb{F}_q).
 \end{array}$$

## Definition (Regular characters of $\text{GL}_N(\mathfrak{o})$ )

A character  $\chi \in \text{Irr}(\text{GL}_N(\mathfrak{o}))$  is called **regular** if the associated similarity class  $\Theta$  consists of regular (cyclic) matrices.

# Regular Elements and Regular Characters

Returning to the case of  $\mathbf{G}$  a general reductive group and  $\mathfrak{g} = \mathrm{Lie}(\mathbf{G})$ .

# Regular Elements and Regular Characters

Returning to the case of  $\mathbf{G}$  a general reductive group and  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Recall that an element  $x \in \mathfrak{g}(\mathbb{F}_q)$  is called **regular** if its centralizer in  $\mathbf{G}(\mathbb{F}_q)$  is of minimal dimension.

# Regular Elements and Regular Characters

Returning to the case of  $\mathbf{G}$  a general reductive group and  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Recall that an element  $x \in \mathfrak{g}(\mathbb{F}_q)$  is called **regular** if its centralizer in  $\mathbf{G}(\mathbb{F}_q)$  is of minimal dimension.

Analogously to the case of  $\text{GL}_N(\mathbb{F}_q)$ , we get an association

# Regular Elements and Regular Characters

Returning to the case of  $\mathbf{G}$  a general reductive group and  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Recall that an element  $x \in \mathfrak{g}(\mathbb{F}_q)$  is called **regular** if its centralizer in  $\mathbf{G}(\mathbb{F}_q)$  is of minimal dimension.

Analogously to the case of  $\text{GL}_N(\mathbb{F}_q)$ , we get an association

$$\begin{array}{c} \chi \in \text{Irr}(G) \text{ of level } \ell > 0 \\ \downarrow \\ \Omega \in \text{Ad}^*(\mathbf{G}(\mathbb{F}_q)) \backslash \widehat{\mathfrak{g}(\mathbb{F}_q)} \\ \updownarrow \\ \Theta \in \text{Ad}(\mathbf{G}(\mathbb{F}_q)) \backslash \mathfrak{g}(\mathbb{F}_q). \end{array}$$

# Regular Elements and Regular Characters

Returning to the case of  $\mathbf{G}$  a general reductive group and  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Recall that an element  $x \in \mathfrak{g}(\mathbb{F}_q)$  is called **regular** if its centralizer in  $\mathbf{G}(\mathbb{F}_q)$  is of minimal dimension.

Analogously to the case of  $\text{GL}_N(\mathbb{F}_q)$ , we get an association

$$\begin{array}{c} \chi \in \text{Irr}(G) \text{ of level } \ell > 0 \\ \downarrow \\ \Omega \in \text{Ad}^*(\mathbf{G}(\mathbb{F}_q)) \backslash \widehat{\mathfrak{g}(\mathbb{F}_q)} \\ \updownarrow \\ \Theta \in \text{Ad}(\mathbf{G}(\mathbb{F}_q)) \backslash \mathfrak{g}(\mathbb{F}_q). \end{array}$$

A character  $\chi \in \text{Irr}(G)$  is said to be regular if the associated orbit  $\Theta$  consists of regular elements of  $\mathfrak{g}(\mathbb{F}_q)$ .

# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].

# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].
  1. Basic properties of regular elements of congruence quotients of  $M_N(\mathfrak{o})$ , including description of the centralizer of a regular element.



# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].
  1. Basic properties of regular elements of congruence quotients of  $M_N(\mathfrak{o})$ , including description of the centralizer of a regular element.
  2. Construction of regular characters of even degree.

# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].
  1. Basic properties of regular elements of congruence quotients of  $M_N(\mathfrak{o})$ , including description of the centralizer of a regular element.
  2. Construction of regular characters of even degree.
- ▶ Hill [Hill, 1995] studied the regular characters of  $GL_N(\mathfrak{o})$  as well.

# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].
  1. Basic properties of regular elements of congruence quotients of  $M_N(\mathfrak{o})$ , including description of the centralizer of a regular element.
  2. Construction of regular characters of even degree.
- ▶ Hill [Hill, 1995] studied the regular characters of  $GL_N(\mathfrak{o})$  as well.
  1. Independent development. Proof of structural properties via implementation of algebraic geometry over local rings of finite length (viz. the *Greenberg functor*).

# Some History

- ▶ Regular characters of  $GL_N(\mathfrak{o})$  we introduced by Shintani in [Shintani, 1968].
  1. Basic properties of regular elements of congruence quotients of  $M_N(\mathfrak{o})$ , including description of the centralizer of a regular element.
  2. Construction of regular characters of even degree.
- ▶ Hill [Hill, 1995] studied the regular characters of  $GL_N(\mathfrak{o})$  as well.
  1. Independent development. Proof of structural properties via implementation of algebraic geometry over local rings of finite length (viz. the *Greenberg functor*).
  2. Construction of several classes of regular characters of odd degree (*cuspidal* and *split-regular* characters).

# Recent Results

- ▶ Krakowski, Onn and Singla [Krakowski et al., 2016]

# Recent Results

- ▶ Krakowski, Onn and Singla [Krakovski et al., 2016]
  1. Classification and enumeration of all regular characters of  $GL_N(\mathfrak{o})$  and  $U_N(\mathfrak{o})$ , for  $\mathfrak{o}$  of odd residual characteristic.

# Recent Results

- ▶ Krakowski, Onn and Singla [Krakovski et al., 2016]
  1. Classification and enumeration of all regular characters of  $GL_N(\mathfrak{o})$  and  $U_N(\mathfrak{o})$ , for  $\mathfrak{o}$  of odd residual characteristic.
  2. Computation of regular representation zeta functions.

# Recent Results

- ▶ Krakowski, Onn and Singla [Krakowski et al., 2016]
  1. Classification and enumeration of all regular characters of  $GL_N(\mathfrak{o})$  and  $U_N(\mathfrak{o})$ , for  $\mathfrak{o}$  of odd residual characteristic.
  2. Computation of regular representation zeta functions.
- ▶ Independently, Stasinski and Stevens [Stasinski and Stevens, 2016] completed the construction of all regular characters of  $GL_N(\mathfrak{o})$ , without restrictions on the residual characteristic of  $\mathfrak{o}$ .



# The Current Research

Consider the case where  $\mathbf{G}$  is a classical group, i.e. a symplectic or an orthogonal group defined over  $\mathfrak{o}^N$ .

# The Current Research

Consider the case where  $\mathbf{G}$  is a classical group, i.e. a symplectic or an orthogonal group defined over  $\mathfrak{o}^N$ .

► **Methods.**

1. An extension of Hill's methods to the study of regular orbits in a classical Lie-algebra over  $\mathfrak{o}$ .
2. Classification of regular conjugacy classes in classical Lie-algebras.

# The Current Research

Consider the case where  $\mathbf{G}$  is a classical group, i.e. a symplectic or an orthogonal group defined over  $\mathfrak{o}^N$ .

## ► **Methods.**

1. An extension of Hill's methods to the study of regular orbits in a classical Lie-algebra over  $\mathfrak{o}$ .
2. Classification of regular conjugacy classes in classical Lie-algebras.

## ► **Results.**

1. A uniform formula for the regular representation zeta function of  $\mathbf{G}(\mathfrak{o})$ .
2. Computation of the regular zeta function of  $\mathbf{G}(\mathfrak{o})$ .

# Regular Zeta Functions of Classical Groups

The regular representation zeta function of  $G = \mathbf{G}(\mathfrak{o})$  is defined by the Dirichlet generating function

$$\zeta_G^{\text{reg.}}(s) = \sum_{\chi \in \text{Irr}^{\text{reg.}}(G)} \chi(1)^{-s} \quad (s \in \mathbb{C}).$$

## Theorem

*Assume  $\text{char}(\mathbb{F}_q) \neq 2$ , and let  $\mathbf{G}$  be a classical group of type  $B_n$ ,  $C_n$  or  $D_n$  over  $\mathfrak{o}$ , with  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ .*

## Theorem

Assume  $\text{char}(\mathbb{F}_q) \neq 2$ , and let  $\mathbf{G}$  be a classical group of type  $B_n$ ,  $C_n$  or  $D_n$  over  $\mathfrak{o}$ , with  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Let  $X \subseteq \text{Ad}(\mathbf{G}(\mathbb{F}_q)) \backslash \backslash \mathfrak{g}(\mathbb{F}_q)$  denote the set of regular orbits, and put  $\mathcal{D}_{\mathfrak{g}(\mathfrak{o})}(s) = \sum_{\Omega \in X} \frac{|\mathbf{G}(\mathbb{F}_q)|}{|\Omega|} \cdot |\Omega|^{-s}$ .

## Theorem

Assume  $\text{char}(\mathbb{F}_q) \neq 2$ , and let  $\mathbf{G}$  be a classical group of type  $B_n$ ,  $C_n$  or  $D_n$  over  $\mathfrak{o}$ , with  $\mathfrak{g} = \text{Lie}(\mathbf{G})$ . Let  $X \subseteq \text{Ad}(\mathbf{G}(\mathbb{F}_q)) \backslash \backslash \mathfrak{g}(\mathbb{F}_q)$  denote the set of regular orbits, and put  $\mathcal{D}_{\mathfrak{g}(\mathfrak{o})}(s) = \sum_{\Omega \in X} \frac{|\mathbf{G}(\mathbb{F}_q)|}{|\Omega|} \cdot |\Omega|^{-s}$ .

$$\zeta_{\mathbf{G}(\mathfrak{o})}^{\text{reg.}}(s) = \frac{\mathcal{D}_{\mathfrak{g}(\mathfrak{o})}(s)}{1 - q^{n-\alpha s}}$$

where  $\alpha = \frac{\dim \mathbf{G} - n}{2}$  and  $n = \text{rk} \mathbf{G}$ .

## Theorem (Cont.)

*Furthermore, there exists polynomials  $f_1, \dots, f_r, g_1, \dots, g_r \in \mathbb{Z}[t]$ , independent of  $\mathfrak{o}$ , such that*

$$\mathcal{D}_{\mathfrak{g}(\mathfrak{o})}(s) = \sum f_i(q) g_i(q)^{-s}.$$



# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ .

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*
  - ▶ Abelian centralizers of regular elements.

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*
  - ▶ Abelian centralizers of regular elements.
  - ▶ Enumeration of regular characters of abelian quotients of  $G = \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*
  - ▶ Abelian centralizers of regular elements.
  - ▶ Enumeration of regular characters of abelian quotients of  $G = \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .
- ▶ Enumeration and dimension computation of regular characters of  $\mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*
  - ▶ Abelian centralizers of regular elements.
  - ▶ Enumeration of regular characters of abelian quotients of  $G = \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .
- ▶ Enumeration and dimension computation of regular characters of  $\mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .

**$\therefore$  Theorem**

# Ingredients

- ▶ Analysis of the locus of regular elements in  $\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^m)$ , for  $m \geq 1$ . Namely
  - ▶ *Regular mod  $\mathfrak{p}^m \iff$  regular mod  $\mathfrak{p}$ .*
  - ▶ Abelian centralizers of regular elements.
  - ▶ Enumeration of regular characters of abelian quotients of  $G = \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .
- ▶ Enumeration and dimension computation of regular characters of  $\mathbf{G}(\mathfrak{o}/\mathfrak{p}^m)$ .

**$\therefore$  Theorem**

- ▶ Classification of regular adjoint orbits in  $\mathfrak{g}(\mathbb{F}_q)$ , and computation of the regular zeta function.

# Why not general reductive groups?

A pivotal point in the analysis of regular elements in [Hill, 1995] is the following.



# Why not general reductive groups?

A pivotal point in the analysis of regular elements in [Hill, 1995] is the following.

## Lemma (Hill's Lemma 3.5)

*Let  $x \in M_N(\mathfrak{o})$ . For any  $r \in \mathbb{N}$  let  $x_r$  denote the canonical image of  $x$  in  $M_N(\mathfrak{o}/\mathfrak{p}^r)$ .*

# Why not general reductive groups?

A pivotal point in the analysis of regular elements in [Hill, 1995] is the following.

## Lemma (Hill's Lemma 3.5)

*Let  $x \in M_N(\mathfrak{o})$ . For any  $r \in \mathbb{N}$  let  $x_r$  denote the canonical image of  $x$  in  $M_N(\mathfrak{o}/\mathfrak{p}^r)$ . Let  $\eta_{r,1} : M_N(\mathfrak{o}/\mathfrak{p}^r) \rightarrow M_N(\mathbb{F}_q)$  be the canonical projection.*

# Why not general reductive groups?

A pivotal point in the analysis of regular elements in [Hill, 1995] is the following.

## Lemma (Hill's Lemma 3.5)

*Let  $x \in M_N(\mathfrak{o})$ . For any  $r \in \mathbb{N}$  let  $x_r$  denote the canonical image of  $x$  in  $M_N(\mathfrak{o}/\mathfrak{p}^r)$ . Let  $\eta_{r,1} : M_N(\mathfrak{o}/\mathfrak{p}^r) \rightarrow M_N(\mathbb{F}_q)$  be the canonical projection. If  $x_1$  is a regular matrix, then*

$$\eta_{r,1} \left( \mathbf{C}_{M_N(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{M_N(\mathbb{F}_q)}(x_1)$$

In different terminology,

*For  $x$  as above, the Lie algebra shadow of  $x$  coincides with the Lie-algebra of  $\mathbb{F}_q$ -points of the algebraic group obtained from  $\mathbf{C}_{M_N(\mathfrak{o})}(x)$  by reduction modulo  $\mathfrak{p}$ .*

## Lemma (Hill's Lemma 3.5)

*Let  $x \in M_N(\mathfrak{o})$ . For any  $r \in \mathbb{N}$  let  $x_r$  denote the canonical image of  $x$  in  $M_N(\mathfrak{o}/\mathfrak{p}^r)$ . Let  $\eta_{r,1} : M_N(\mathfrak{o}/\mathfrak{p}^r) \rightarrow M_N(\mathbb{F}_q)$  be the canonical projection. If  $x_1$  is a regular matrix, then*

$$\eta_{r,1} \left( \mathbf{C}_{M_N(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{M_N(\mathbb{F}_q)}(x_1)$$

Hill's Lemma 3.5 generalizes to general reductive Lie-algebras,  
i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

Hill's Lemma 3.5 generalizes to general reductive Lie-algebras, i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

In order to facilitate the analysis of regular elements, one must be able to deduce an analogous result in the group setting, namely

$$\eta_{r,1} \left( \mathbf{C}_{G(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{G(\mathbb{F}_q)}(x_1).$$

Hill's Lemma 3.5 generalizes to general reductive Lie-algebras, i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

In order to facilitate the analysis of regular elements, one must be able to deduce an analogous result in the group setting, namely

$$\eta_{r,1} \left( \mathbf{C}_{G(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{G(\mathbb{F}_q)}(x_1).$$

As a first step- is it possible to show that the two groups are of the same dimension? Is it possible to show the Lie-algebra shadow and the group shadow of an element are of equal dimension?

Hill's Lemma 3.5 generalizes to general reductive Lie-algebras, i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

In order to facilitate the analysis of regular elements, one must be able to deduce an analogous result in the group setting, namely

$$\eta_{r,1} \left( \mathbf{C}_{G(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{G(\mathbb{F}_q)}(x_1).$$

As a first step- is it possible to show that the two groups are of the same dimension? Is it possible to show the Lie-algebra shadow and the group shadow of an element are of equal dimension?

- For  $\mathbf{G} = \mathrm{GL}_N$  this is done by adhering to the denseness of  $\mathrm{GL}_N$  in  $M_N$ .



Hill's Lemma 3.5 generalizes to general reductive Lie-algebras, i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

In order to facilitate the analysis of regular elements, one must be able to deduce an analogous result in the group setting, namely

$$\eta_{r,1} \left( \mathbf{C}_{G(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{G(\mathbb{F}_q)}(x_1).$$

As a first step- is it possible to show that the two groups are of the same dimension? Is it possible to show the Lie-algebra shadow and the group shadow of an element are of equal dimension?

- ▶ For  $\mathbf{G} = \mathrm{GL}_N$  this is done by adhering to the denseness of  $\mathrm{GL}_N$  in  $\mathbf{M}_N$ .
- ▶ For the classical groups, one may prove this statement by invoking the Cayley map  $y \mapsto (1 - y)^{-1}(1 + y)$ .

Hill's Lemma 3.5 generalizes to general reductive Lie-algebras, i.e.

$$\eta_{r,1} \left( \mathbf{C}_{\mathfrak{g}(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{\mathfrak{g}(\mathbb{F}_q)}(x_1).$$

In order to facilitate the analysis of regular elements, one must be able to deduce an analogous result in the group setting, namely

$$\eta_{r,1} \left( \mathbf{C}_{G(\mathfrak{o}/\mathfrak{p}^r)}(x_r) \right) = \mathbf{C}_{G(\mathbb{F}_q)}(x_1).$$

As a first step- is it possible to show that the two groups are of the same dimension? Is it possible to show the Lie-algebra shadow and the group shadow of an element are of equal dimension?

- ▶ For  $\mathbf{G} = \mathrm{GL}_N$  this is done by adhering to the denseness of  $\mathrm{GL}_N$  in  $\mathbf{M}_N$ .
- ▶ For the classical groups, one may prove this statement by invoking the Cayley map  $y \mapsto (1 - y)^{-1}(1 + y)$ .
- ▶ For a general reductive group - ???

# Examples

$$\zeta_{\mathrm{SL}_2(\mathfrak{o})}^{\mathrm{reg.}}(s) = \frac{4q\left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2}(q^2-q)^{-s} + \left(\frac{q-1}{2}\right)^2(q^2+q)^{-s}}{1-q^{1-s}}, \quad (\text{Jaikin}).$$

$$\zeta_{\mathrm{SL}_1(D)}^{\mathrm{reg.}}(s) = \sum_{\lambda=0}^{\ell-2} \iota^2 (q-1) \frac{q^\lambda \cdot \left( q^{-\lambda \frac{(\ell-1)}{2}} \cdot \frac{q^\ell-1}{\iota(q-1)} \right)^{-s}}{1-q^{(\ell-1)-\binom{\ell}{2}s}}, \quad (\text{Sh})$$

( $\deg D$  is prime  $\neq p$  and  $\iota = \#\mu_\ell(\mathbb{F}_q)$ ).

## Remark

In both cases above  $\zeta_{\mathbf{G}(\mathfrak{o})}(s) = \zeta_{\mathbf{G}(\mathbb{F}_q)}(s) + \zeta_{\mathbf{G}(\mathfrak{o})}^{\mathrm{reg.}}(s)$ .

For  $\mathbf{G}_+ = \mathrm{GL}_n$  and  $\mathbf{G}_- = \mathrm{U}_n$

$$\zeta_{\mathbf{G}_\epsilon(\mathfrak{o})}^{\mathrm{reg.}}(s) = \frac{1}{1 - q^{n - \binom{n}{2}s}} \cdot \sum_{\tau \in \mathcal{A}_n} u_\epsilon^\tau(q) \prod_{i=1}^n \binom{\sum_j \tau_{i,j}}{\tau_{i,1}, \tau_{i,2}, \dots, \tau_{i,n}} \binom{w_d(q)}{\sum_j \tau_{i,j}} \left( \frac{|\mathbf{G}_\epsilon(\mathbb{F}_q)|}{u_\epsilon^\tau(q)} \right)^{-s},$$

(KOS),

where

- ▶  $\mathcal{A}_n$  is the set of  $n$ -typical matrices,
- ▶  $w_d(q)$  is the number of irreducible polynomials of degree  $d$  over  $\mathbb{F}_q$ ,
- ▶  $u_\epsilon^\tau(q)$  is the cardinality of the centralizer in  $\mathbf{G}_\epsilon(\mathbb{F}_q)$  of a regular element associated to an  $n$ -typical matrix  $\tau \in \mathcal{A}_n$ .

# Regular Zeta Functions

Let  $\mathcal{X}_n$  consist of triplets  $(r, A, B) \in \mathbb{Z}_{\geq 0} \times M_n(\mathbb{Z}_{\geq 0}) \times M_n(\mathbb{Z}_{\geq 0})$  with  $r + \sum_{d,e} (A_{d,e} + B_{d,e}) = n$ , and  $\mathcal{X}_n^0$  the subset of elements with  $r = 0$ .

$$\begin{aligned} \mathcal{D}_{\mathfrak{sp}_{2n}(\mathfrak{o})} = & q^n \sum_{(0,A,B) \in \mathcal{X}_n^0} M_{(0,A,B)} \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}} \\ & \left( q^{2n^2} \frac{\prod_{i=1}^n (1 - q^{-2i})}{\prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}}} \right)^{-s} \\ + 4q^n & \sum_{(r,A,B) \in \mathcal{X}_n \setminus \mathcal{X}_n^0} M_{(r,A,B)} \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}} \\ & \left( q^{2n^2} \frac{\prod_{i=1}^n (1 - q^{-2i})}{2 \cdot \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}}} \right)^{-s}. \end{aligned}$$

$$\mathcal{D}_{\mathfrak{so}_{2n+1}(\mathfrak{o})} = q^n \sum_{(r,A,B) \in \mathcal{X}_n} M_{(r,A,B)} \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}} \left( q^{2n^2} \frac{\prod_{i=1}^n (1 - q^{-2i})}{\prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}}} \right)^{-s}.$$

Let  $\mathcal{X}_n^0$  denote the set of triplets  $\mathbf{t} = (r, A, B) \in \mathcal{X}_n$  with  $r = 0$ , and let  $\mathcal{X}_n^{0,+}$  denote the subset of  $\mathcal{X}_n^0$  consisting of element  $(0, A, B)$  such that  $\sum_{d,e} eA_{d,e}$  is even and  $\mathcal{X}_n^{0,-} = \mathcal{X}_n^0 \setminus \mathcal{X}_n^{0,+}$ .

$$\begin{aligned} \mathbb{D}_{\mathfrak{so}_{2n}^+(\mathfrak{o})} = & q^n \sum_{(0,A,B) \in \mathcal{X}_n^{0,+}} M_{(0,A,B)} \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}} \\ & \left( q^{2n^2-n} \frac{(1 - q^{-n}) \prod_{i=1}^{n-1} (1 - q^{-2i})}{\prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}}} \right)^{-s} \\ & + 2q^n \sum_{(r,A,B) \in \mathcal{X}_n \setminus \mathcal{X}_n^0} M_{(r,A,B)} \prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}} \\ & \left( q^{2n^2-n} \frac{(1 - q^{-n}) \prod_{i=1}^{n-1} (1 - q^{-2i})}{\prod_{d,e} (1 + q^{-d})^{A_{d,e}} (1 - q^{-d})^{B_{d,e}}} \right)^{-s}, \end{aligned}$$

$$\begin{aligned}
\mathbb{D}_{\mathfrak{so}_{2n}^-(\mathfrak{o})} = & q^n \sum_{(0,A,B) \in \mathcal{X}_n^{0,-}} M_{(0,A,B)} \prod_{d,e} (1+q^{-d})^{A_{d,e}} (1-q^{-d})^{B_{d,e}} \\
& \left( q^{2n^2-n} \frac{(1+q^{-n}) \prod_{i=1}^{n-1} (1-q^{-2i})}{\prod_{d,e} (1+q^{-d})^{A_{d,e}} (1-q^{-d})^{B_{d,e}}} \right)^{-s} \\
& + 2q^n \sum_{(r,A,B) \in \mathcal{X}_n \setminus \mathcal{X}_n^0} M_{(r,A,B)} \prod_{d,e} (1+q^{-d})^{A_{d,e}} (1-q^{-d})^{B_{d,e}} \\
& \left( q^{2n^2-n} \frac{(1+q^{-n}) \prod_{i=1}^{n-1} (1-q^{-2i})}{\prod_{d,e} (1+q^{-d})^{A_{d,e}} (1-q^{-d})^{B_{d,e}}} \right)^{-s}. \quad (0.a)
\end{aligned}$$



# Questions?

# Thank you!



Hill, G. (1995).

Regular elements and regular characters of  $GL_n(\mathcal{O})$ .  
*Journal of Algebra*, 174:610–635.



Krakovski, R., Onn, U., and Singla, P. (2016).

On regular representations of  $GL_n(\mathfrak{o})$  and  $GU_n(\mathfrak{o})$ .



Shintani, T. (1968).

On certain square integrable irreducible unitary representations of some  $p$ -adic linear groups.

*Proceedings of the Japan Academy, Series A, Mathematical Sciences*, 44(1):1–3.



Stasinski, A. and Stevens, S. (2016).

The regular representations of  $GL_N$  over finite local principal ideal rings.