

# Algebraic Geometry 2

## Tutorial session 4

Lecturer: Rami Aizenbud  
TA: Shai Shechter

May 17, 2020

# The pullback and direct image - recollections

Let  $f : X \rightarrow Y$  be a continuous map of topological space, and let  $\mathcal{F}$  be a sheaf on  $X$  and  $\mathcal{G}$  a sheaf on  $Y$ .

# The pullback and direct image - recollections

Let  $f : X \rightarrow Y$  be a continuous map of topological space, and let  $\mathcal{F}$  be a sheaf on  $X$  and  $\mathcal{G}$  a sheaf on  $Y$ . Recall that we defined the pullback of  $\mathcal{F}$  to be the sheaf  $f^*\mathcal{F}$  on  $Y$ , defined by

$$f^*\mathcal{F}(V) = \mathcal{F}(f^{-1}(V)) \quad \text{for } V \subseteq Y \text{ open.}$$

# The pullback and direct image - recollections

Let  $f : X \rightarrow Y$  be a continuous map of topological space, and let  $\mathcal{F}$  be a sheaf on  $X$  and  $\mathcal{G}$  a sheaf on  $Y$ . Recall that we defined the pullback of  $\mathcal{F}$  to be the sheaf  $f_*\mathcal{F}$  on  $Y$ , defined by

$$f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}(V)) \quad \text{for } V \subseteq Y \text{ open.}$$

We also defined the direct image sheaf  $f^{-1}\mathcal{G}$  on  $X$ , by sheafification of the presheaf defined by

$$f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U) \text{ open}} \mathcal{G}(V) \quad \text{for } U \subseteq X \text{ open.}$$

# Adjointness of pullback and direct image

## Exercise

Let  $f : X \rightarrow Y$  be a continuous map,  $\mathcal{F}$  a sheaf on  $X$  and  $\mathcal{G}$  a sheaf on  $Y$ . Then there exists a natural bijection

$$\mathrm{Hom}_{\mathbf{Sh}(X)}(f^{-1}\mathcal{G}, \mathcal{F}) \simeq \mathrm{Hom}_{\mathbf{Sh}(Y)}(\mathcal{G}, f_*\mathcal{F}).$$

# Adjontness of pullback and direct image

## Exercise

Let  $f : X \rightarrow Y$  be a continuous map,  $\mathcal{F}$  a sheaf on  $X$  and  $\mathcal{G}$  a sheaf on  $Y$ . Then there exists a natural bijection

$$\mathrm{Hom}_{\underline{\mathbf{Sh}}(X)}(f^{-1}\mathcal{G}, \mathcal{F}) \simeq \mathrm{Hom}_{\underline{\mathbf{Sh}}(Y)}(\mathcal{G}, f_*\mathcal{F}).$$

## Proof.

We construct natural maps going in both directions, and verify that their compositions are equivalent to identity (some details are left as exercises).

## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ :

## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ : Let  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ . We construct  $F(\varphi)_V$  for  $V$  in a basis for the topology on  $Y$ . Recall from exercise, that  $Y$  has a basis of subsets  $V$  such that for  $U \subseteq f^{-1}(V) \subseteq X$  open, we have  $f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U)} \mathcal{G}(V)$ .



## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ : Let  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ . We construct  $F(\varphi)_V$  for  $V$  in a basis for the topology on  $Y$ . Recall from exercise, that  $Y$  has a basis of subsets  $V$  such that for  $U \subseteq f^{-1}(V) \subseteq X$  open, we have  $f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U)} \mathcal{G}(V)$ . Let  $V \subseteq Y$  be such, and  $U = f^{-1}(V)$ .

## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ : Let  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ . We construct  $F(\varphi)_V$  for  $V$  in a basis for the topology on  $Y$ . Recall from exercise, that  $Y$  has a basis of subsets  $V$  such that for  $U \subseteq f^{-1}(V) \subseteq X$  open, we have  $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$ . Let  $V \subseteq Y$  be such, and  $U = f^{-1}(V)$ . By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ : Let  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ . We construct  $F(\varphi)_V$  for  $V$  in a basis for the topology on  $Y$ . Recall from exercise, that  $Y$  has a basis of subsets  $V$  such that for  $U \subseteq f^{-1}(V) \subseteq X$  open, we have  $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$ . Let  $V \subseteq Y$  be such, and  $U = f^{-1}(V)$ . By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

In particular,  $\text{Hom}(f^{-1}\mathcal{G}(U), \mathcal{F}(U)) = \text{Hom}(\mathcal{G}(V), f_*\mathcal{F}(V))$ , and we can define  $F(\varphi)_V = \varphi_{f^{-1}(V)}$ .

## Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ : Let  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ . We construct  $F(\varphi)_V$  for  $V$  in a basis for the topology on  $Y$ . Recall from exercise, that  $Y$  has a basis of subsets  $V$  such that for  $U \subseteq f^{-1}(V) \subseteq X$  open, we have  $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$ . Let  $V \subseteq Y$  be such, and  $U = f^{-1}(V)$ . By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

In particular,  $\text{Hom}(f^{-1}\mathcal{G}(U), \mathcal{F}(U)) = \text{Hom}(\mathcal{G}(V), f_*\mathcal{F}(V))$ , and we can define  $F(\varphi)_V = \varphi_{f^{-1}(V)}$ . This gives rise to a sheaf morphism  $F(\varphi) \in \text{Hom}(\mathcal{G}, f_*\mathcal{F})$  (Ex).



## Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$ : Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ ,  $U \subseteq X$  open and  $f(U) \subseteq V \subseteq Y$  open, we have a map  $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$ , given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

## Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$ : Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ ,  $U \subseteq X$  open and  $f(U) \subseteq V \subseteq Y$  open, we have a map  $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$ , given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative.

## Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$ : Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ ,  $U \subseteq X$  open and  $f(U) \subseteq V \subseteq Y$  open, we have a map  $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$ , given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative. The maps  $(g_U)_{U \subseteq X \text{ open}}$  comprise a morphism of presheaves from the presheaf  $(U \mapsto \lim_{V \supseteq f(U)} \mathcal{G}(V))$  to  $\mathcal{F}$  (Ex).

## Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$ : Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ ,  $U \subseteq X$  open and  $f(U) \subseteq V \subseteq Y$  open, we have a map  $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$ , given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative. The maps  $(g_U)_{U \subseteq X \text{ open}}$  comprise a morphism of presheaves from the presheaf  $(U \mapsto \lim_{V \supseteq f(U)} \mathcal{G}(V))$  to  $\mathcal{F}$  (Ex). Sheafifying, we set  $G(\psi) = \tilde{g} : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ .





## Proof.

We need to show  $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$  and  $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$ .

- Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$  and  $V \subseteq Y$  open, we have  $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$ , where the RHS is given by a direct limit over open sets containing  $f(f^{-1}(V)) = V$ .

## Proof.

We need to show  $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$  and  $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$ .

- Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$  and  $V \subseteq Y$  open, we have  $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$ , where the RHS is given by a direct limit over open sets containing  $f(f^{-1}(V)) = V$ . Hence it is a filtered limit with a terminal object, and equals to its value at  $V$ .

## Proof.

We need to show  $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$  and  $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$ .

- Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$  and  $V \subseteq Y$  open, we have  $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$ , where the RHS is given by a direct limit over open sets containing  $f(f^{-1}(V)) = V$ . Hence it is a filtered limit with a terminal object, and equals to its value at  $V$ . It follows that  $(FG\psi)_V = (G\psi)_{f^{-1}(V)} = \psi_V$ .



## Proof.

We need to show  $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$  and  $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$ .

- Given  $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$  and  $V \subseteq Y$  open, we have  $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$ , where the RHS is given by a direct limit over open sets containing  $f(f^{-1}(V)) = V$ . Hence it is a filtered limit with a terminal object, and equals to its value at  $V$ . It follows that  $(FG\psi)_V = (G\psi)_{f^{-1}(V)} = \psi_V$ .
- The equivalence  $(GF\varphi)_U = \varphi_U$  for all  $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$  follows similarly, by unfolding the definitions (Ex).



# Schemes

# The spectrum of a ring

Let  $A$  be a commutative unital ring.

## Definition

The spectrum of  $R$  is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

# The spectrum of a ring

Let  $A$  be a commutative unital ring.

## Definition

The spectrum of  $R$  is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

## Examples

- 1  $\operatorname{Spec}(k) = \{*\}$  for any field  $k$ .

# The spectrum of a ring

Let  $A$  be a commutative unital ring.

## Definition

The spectrum of  $R$  is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

## Examples

- 1  $\operatorname{Spec}(k) = \{*\}$  for any field  $k$ .
- 2  $\operatorname{Spec}(k[x]) =: \mathbb{A}_k^1 \approx \{\text{irreducible monic polynomials in } x\} \sqcup \{0\}$



# The spectrum of a ring

Let  $A$  be a commutative unital ring.

## Definition

The spectrum of  $R$  is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

## Examples

- 1  $\operatorname{Spec}(k) = \{*\}$  for any field  $k$ .
- 2  $\operatorname{Spec}(k[x]) =: \mathbb{A}_k^1 \approx \{\text{irreducible monic polynomials in } x\} \sqcup \{0\}$
- 3  $\operatorname{Spec}(\mathbb{Z}) = \{\langle 0 \rangle\} \sqcup \{\langle p \rangle : p \text{ prime}\}.$

# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

Exercise

# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

## Exercise

- 1  $V((0)) = A$  and  $V(A) = \emptyset$ .

# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

## Exercise

- 1  $V((0)) = A$  and  $V(A) = \emptyset$ .
- 2  $V(IJ) = V(I) \cup V(J)$ .

# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

## Exercise

- 1  $V((0)) = A$  and  $V(A) = \emptyset$ .
- 2  $V(IJ) = V(I) \cup V(J)$ .
- 3 Given a collection  $\{I_\alpha\}$  of ideals,  $V(\sum I_\alpha) = \bigcap V(I_\alpha)$ .

# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

## Exercise

- 1  $V((0)) = A$  and  $V(A) = \emptyset$ .
- 2  $V(IJ) = V(I) \cup V(J)$ .
- 3 Given a collection  $\{I_\alpha\}$  of ideals,  $V(\sum I_\alpha) = \bigcap V(I_\alpha)$ .

## Proof.

Home exercise.



# The spectrum of a ring - topology

Given  $I \triangleleft A$ , define  $V(I) := \{\mathfrak{p} \in \text{Spec}(A) : I \subseteq \mathfrak{p}\}$ .

## Exercise

- 1  $V((0)) = A$  and  $V(A) = \emptyset$ .
- 2  $V(IJ) = V(I) \cup V(J)$ .
- 3 Given a collection  $\{I_\alpha\}$  of ideals,  $V(\sum I_\alpha) = \bigcap V(I_\alpha)$ .

## Proof.

Home exercise.



The collection  $\{V(I) : I \triangleleft R\}$  is the set of closed sets for a topology on  $\text{Spec}(R)$ , which is known as the *Zariski Topology* of  $R$ .

## Exercise

Let  $A$  be a ring.

- 1 Show that  $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$ , for all  $\mathfrak{p} \in \operatorname{Spec}(A)$  and, in particular, that  $\{\mathfrak{p}\}$  is closed iff  $\mathfrak{p}$  is maximal.
- 2 Show that, if  $A$  is a domain, then  $\{(0)\}$  is a dense point.



## Exercise

Let  $A$  be a ring.

- 1 Show that  $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$ , for all  $\mathfrak{p} \in \operatorname{Spec}(A)$  and, in particular, that  $\{\mathfrak{p}\}$  is closed iff  $\mathfrak{p}$  is maximal.
- 2 Show that, if  $A$  is a domain, then  $\{(0)\}$  is a dense point.

## Solution.



## Exercise

Let  $A$  be a ring.

- 1 Show that  $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$ , for all  $\mathfrak{p} \in \operatorname{Spec}(A)$  and, in particular, that  $\{\mathfrak{p}\}$  is closed iff  $\mathfrak{p}$  is maximal.
- 2 Show that, if  $A$  is a domain, then  $\{(0)\}$  is a dense point.

## Solution.

- 1 By definition, and by the previous exercise:

$$\overline{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in F \text{ closed}} F = \bigcap_{\substack{I \triangleleft R \\ I \subseteq \mathfrak{p}}} V(I) = V\left(\sum_{I \subseteq \mathfrak{p}} I\right) = V(\mathfrak{p}).$$

In particular,  $\{\mathfrak{p}\}$  is closed iff  $\{\mathfrak{p}\} = V(\mathfrak{p})$  which occurs iff  $\mathfrak{p}$  is maximal (o/w, take  $\mathfrak{m} \supsetneq \mathfrak{p}$  maximal).



## Exercise

Let  $A$  be a ring.

- 1 Show that  $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$ , for all  $\mathfrak{p} \in \operatorname{Spec}(A)$  and, in particular, that  $\{\mathfrak{p}\}$  is closed iff  $\mathfrak{p}$  is maximal.
- 2 Show that, if  $A$  is a domain, then  $\{(0)\}$  is a dense point.

## Solution.

- 1 By definition, and by the previous exercise:

$$\overline{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in F \text{ closed}} F = \bigcap_{\substack{I \triangleleft R \\ I \subseteq \mathfrak{p}}} V(I) = V\left(\sum_{I \subseteq \mathfrak{p}} I\right) = V(\mathfrak{p}).$$

In particular,  $\{\mathfrak{p}\}$  is closed iff  $\{\mathfrak{p}\} = V(\mathfrak{p})$  which occurs iff  $\mathfrak{p}$  is maximal (o/w, take  $\mathfrak{m} \supsetneq \mathfrak{p}$  maximal).

- 2 Note:  $(0) \in \operatorname{Spec}(R)$  iff  $R$  is a domain, in which case  $V(0) = R$ .



## Exercise

Show that  $\text{Spec}(A)$  is irreducible iff  $A$  is a domain.

## Exercise

Show that  $\text{Spec}(A)$  is irreducible iff  $A$  is a domain.

## Solution.

Assume  $A$  is domain, and  $0 \neq I_1, I_2 \triangleleft A$  proper ideals such that  $\text{Spec}(A) = V(I_1) \cup V(I_2)$ . Let  $f_i \in I_i$  ( $i = 1, 2$ ). Then, since

$$V(I_1) \cup V(I_2) \subseteq V((f_1)) \cup V((f_2)) = V((f_1 f_2)),$$

we have that  $f_1 f_2$  is necessarily nilpotent, and since  $A$  is a domain, either  $f_1 = 0$  or  $f_2 = 0$ . By fixing  $0 \neq f_1 \in I_1$  (wlog, assuming such exists) and letting  $f_2 \in I_2$  vary we deduce that  $I_2 = (0)$  and hence  $V(I_2) = \text{Spec}(A)$ .

## Exercise

Show that  $\text{Spec}(A)$  is irreducible iff  $A$  is a domain.

## Solution.

Assume  $A$  is domain, and  $0 \neq I_1, I_2 \triangleleft A$  proper ideals such that  $\text{Spec}(A) = V(I_1) \cup V(I_2)$ . Let  $f_i \in I_i$  ( $i = 1, 2$ ). Then, since

$$V(I_1) \cup V(I_2) \subseteq V((f_1)) \cup V((f_2)) = V((f_1 f_2)),$$

we have that  $f_1 f_2$  is necessarily nilpotent, and since  $A$  is a domain, either  $f_1 = 0$  or  $f_2 = 0$ . By fixing  $0 \neq f_1 \in I_1$  (wlog, assuming such exists) and letting  $f_2 \in I_2$  vary we deduce that  $I_2 = (0)$  and hence  $V(I_2) = \text{Spec}(A)$ .

For the converse implication, assume  $\text{Spec}(A)$  is irreducible and let  $f_1, f_2 \in A$  non-nilpotents. By assumption

$V((f_1)) \cup V((f_2)) = V((f_1 f_2)) \subsetneq \text{Spec}(A)$ , and hence there exists  $\mathfrak{p} \in \text{Spec}(A)$  such that  $f_1 f_2 \notin \mathfrak{p}$ . In particular,  $f_1 f_2 \neq 0$ .



The last two exercises imply the following:

The last two exercises imply the following:

## Corollary

*If  $\text{Spec}(A)$  is irreducible then there exists  $\xi \in \text{Spec}(A)$  such that  $\overline{\xi} = \text{Spec}(A)$ .*



The last two exercises imply the following:

## Corollary

*If  $\text{Spec}(A)$  is irreducible then there exists  $\xi \in \text{Spec}(A)$  such that  $\overline{\xi} = \text{Spec}(A)$ . Such an element is called a generic point of  $\text{Spec}(A)$ .*

# Ringed spaces and locally ringed spaces

## Definition

# Ringed spaces and locally ringed spaces

## Definition

- 1 A *ringed space* is a pair  $(X, \mathcal{O}_X)$  consisting of a topological space and a sheaf of rings  $\mathcal{O}_X$  on  $X$ .

# Ringed spaces and locally ringed spaces

## Definition

- 1 A *ringed space* is a pair  $(X, \mathcal{O}_X)$  consisting of a topological space and a sheaf of rings  $\mathcal{O}_X$  on  $X$ .
- 2 A *morphism of ringed spaces* is a pair  $(f, f^\#)$ , in which  $f$  is a continuous map  $X \rightarrow Y$  and  $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$  is a morphism of sheaves of rings on  $Y$ .

# Ringed spaces and locally ringed spaces

## Definition

- 1 A *ringed space* is a pair  $(X, \mathcal{O}_X)$  consisting of a topological space and a sheaf of rings  $\mathcal{O}_X$  on  $X$ .
- 2 A *morphism of ringed spaces* is a pair  $(f, f^\#)$ , in which  $f$  is a continuous map  $X \rightarrow Y$  and  $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$  is a morphism of sheaves of rings on  $Y$ .
- 3  $(X, \mathcal{O}_X)$  is a *locally ringed space* if, in addition, the stalk  $\mathcal{O}_{X,x}$  is a local ring for all  $x \in X$ .

# Ringed spaces and locally ringed spaces

## Definition

- 1 A *ringed space* is a pair  $(X, \mathcal{O}_X)$  consisting of a topological space and a sheaf of rings  $\mathcal{O}_X$  on  $X$ .
- 2 A *morphism of ringed spaces* is a pair  $(f, f^\#)$ , in which  $f$  is a continuous map  $X \rightarrow Y$  and  $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$  is a morphism of sheaves of rings on  $Y$ .
- 3  $(X, \mathcal{O}_X)$  is a *locally ringed space* if, in addition, the stalk  $\mathcal{O}_{X,x}$  is a local ring for all  $x \in X$ .
- 4 A morphism of locally ringed spaces  $(f, f^\#)$  is a morphism of ringed spaces with the **added requirement** that  $f^\#_x : \mathcal{O}_{Y,f(x)} \rightarrow f_{*}\mathcal{O}_{X,x}$  is a local homomorphism (i.e. preimage of the maximal ideal is maximal) for any  $x \in X$ .

# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

- for any  $\mathfrak{p} \in \mathrm{Spec}(A)$ , the stalk  $\mathcal{O}_{\mathfrak{p}}$  of  $\mathcal{O}$  at  $\mathfrak{p}$  is isomorphic to the local ring  $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$ , and



# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

- for any  $\mathfrak{p} \in \mathrm{Spec}(A)$ , the stalk  $\mathcal{O}_{\mathfrak{p}}$  of  $\mathcal{O}$  at  $\mathfrak{p}$  is isomorphic to the local ring  $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$ , and
- for any  $f \in A$ , let  $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$ ; this is an open set of  $\mathrm{Spec}(A)$ . Then  $\mathcal{O}(D(f)) \simeq f^{-1}A$ .

# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

- for any  $\mathfrak{p} \in \mathrm{Spec}(A)$ , the stalk  $\mathcal{O}_{\mathfrak{p}}$  of  $\mathcal{O}$  at  $\mathfrak{p}$  is isomorphic to the local ring  $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$ , and
- for any  $f \in A$ , let  $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$ ; this is an open set of  $\mathrm{Spec}(A)$ . Then  $\mathcal{O}(D(f)) \simeq f^{-1}A$ .

Thus  $(\mathrm{Spec}(A), \mathcal{O})$  is a locally ring space, with  $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$ .

# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

- for any  $\mathfrak{p} \in \mathrm{Spec}(A)$ , the stalk  $\mathcal{O}_{\mathfrak{p}}$  of  $\mathcal{O}$  at  $\mathfrak{p}$  is isomorphic to the local ring  $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$ , and
- for any  $f \in A$ , let  $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$ ; this is an open set of  $\mathrm{Spec}(A)$ . Then  $\mathcal{O}(D(f)) \simeq f^{-1}A$ .

Thus  $(\mathrm{Spec}(A), \mathcal{O})$  is a locally ring space, with  $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$ .

## Explicit construction

The sheaf  $\mathcal{O}$  is defined by sheafifying the presheaf on  $\mathrm{Spec}(A)$  whose stalks are given by  $A_{\mathfrak{p}} = \lim_{\mathfrak{p} \in D(f), f \in A} A_f$ .

# The sheaf of rings on $\mathrm{Spec}(A)$

Let  $A$  be a ring. The spectrum of  $A$  has an associated sheaf of rings  $\mathcal{O}$ , which has the following properties:

- for any  $\mathfrak{p} \in \mathrm{Spec}(A)$ , the stalk  $\mathcal{O}_{\mathfrak{p}}$  of  $\mathcal{O}$  at  $\mathfrak{p}$  is isomorphic to the local ring  $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$ , and
- for any  $f \in A$ , let  $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$ ; this is an open set of  $\mathrm{Spec}(A)$ . Then  $\mathcal{O}(D(f)) \simeq f^{-1}A$ .

Thus  $(\mathrm{Spec}(A), \mathcal{O})$  is a locally ring space, with  $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$ .

## Explicit construction

The sheaf  $\mathcal{O}$  is defined by sheafifying the presheaf on  $\mathrm{Spec}(A)$  whose stalks are given by  $A_{\mathfrak{p}} = \lim_{\mathfrak{p} \in D(f), f \in A} A_f$ . Explicitly,  $\mathcal{O}(U)$  consists of functions  $s : U \rightarrow \bigsqcup_{\mathfrak{p} \in U} A_{\mathfrak{p}}$  for which there exists an open cover  $U = \bigcup_{\alpha} V_{\alpha}$  such that, given  $\alpha$ ,  $\exists a_{\alpha}, b_{\alpha} \in A$  with  $V_{\alpha} \subseteq D(b_{\alpha})$ , such that  $s(\mathfrak{q}) =$  “the image of  $a_{\alpha}/b_{\alpha}$  in  $A_{\mathfrak{q}}$ ”, for all  $\mathfrak{q} \in V_{\alpha}$ .

## Proposition (2.3 in Hartshorne)

- ① *Let  $\varphi : A \rightarrow B$  be a homomorphism of rings. Then  $\varphi$  induces a natural morphism of locally ringed spaces*

$$(f, f^\#) : (\mathrm{Spec}(B), \mathcal{O}_{\mathrm{Spec}(B)}) \rightarrow (\mathrm{Spec}(A), \mathcal{O}_{\mathrm{Spec}(A)}).$$

- ② *Conversely, any morphism of locally ring spaces  $(f, f^\#)$  as above is induced from a ring homomorphism  $\varphi : A \rightarrow B$ .*

## Example

Let  $R$  be a dvr and  $K = \text{Frac}(R)$ , its field of fractions.

## Example

Let  $R$  be a dvr and  $K = \text{Frac}(R)$ , its field of fractions. The space  $T = \text{Spec}(R)$  has two points. One of them, denoted  $t_0$ , is closed and corresponds to the maximal ideal  $\mathfrak{m}$  of  $R$ . The second, denoted  $t_0$  is open and dense, and corresponds to the zero ideal.

## Example

Let  $R$  be a dvr and  $K = \text{Frac}(R)$ , its field of fractions. The space  $T = \text{Spec}(R)$  has two points. One of them, denoted  $t_0$ , is closed and corresponds to the maximal ideal  $\mathfrak{m}$  of  $R$ . The second, denoted  $t_1$  is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$



## Example

Let  $R$  be a dvr and  $K = \text{Frac}(R)$ , its field of fractions. The space  $T = \text{Spec}(R)$  has two points. One of them, denoted  $t_0$ , is closed and corresponds to the maximal ideal  $\mathfrak{m}$  of  $R$ . The second, denoted  $t_1$  is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$

The map  $f : S = \text{Spec}(K) \rightarrow T$ , sending the only point in  $\text{Spec}(K)$  to  $t_1$ , is part of a morphism  $(f, f^\#)$  of **locally** ringed spaces, since the corresponding stalks are both  $K$ .

## Example

Let  $R$  be a dvr and  $K = \text{Frac}(R)$ , its field of fractions. The space  $T = \text{Spec}(R)$  has two points. One of them, denoted  $t_0$ , is closed and corresponds to the maximal ideal  $\mathfrak{m}$  of  $R$ . The second, denoted  $t_1$  is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$

The map  $f : S = \text{Spec}(K) \rightarrow T$ , sending the only point in  $\text{Spec}(K)$  to  $t_1$ , is part of a morphism  $(f, f^\#)$  of **locally** ringed spaces, since the corresponding stalks are both  $K$ . Therefore, it is induced from a ring homomorphism  $\varphi : R \rightarrow K$ .

## Example - contd

On the other hand, if  $\text{Im}(f) = \{t_0\}$ , and  $(f, f^\#)$  is a morphism of ringed spaces, then  $f^\#$  induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

## Example - contd

On the other hand, if  $\text{Im}(f) = \{t_0\}$ , and  $(f, f^\#)$  is a morphism of ringed spaces, then  $f^\#$  induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

Assuming  $R/\mathfrak{m} \not\cong K$ , this is **not** a local homomorphism.

## Example - contd

On the other hand, if  $\text{Im}(f) = \{t_0\}$ , and  $(f, f^\#)$  is a morphism of ringed spaces, then  $f^\#$  induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

Assuming  $R/\mathfrak{m} \not\cong K$ , this is **not** a local homomorphism. Therefore, in this case,  $(f, f^\#)$  is not induced from any morphism  $R \rightarrow K$ .

# Proof of proposition

Proof.

# Proof of proposition

## Proof.

- 1 Let  $\varphi : A \rightarrow B$  be a ring homomorphism. Define  $f : \operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$  by  $f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \in \operatorname{Spec}(A)$ .

# Proof of proposition

## Proof.

- ① Let  $\varphi : A \rightarrow B$  be a ring homomorphism. Define  $f : \operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$  by  $f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \in \operatorname{Spec}(A)$ .

$f$  is continuous. Given  $I \triangleleft A$ , we have

$$\mathfrak{p} \in f^{-1}(V(I)) \iff f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \supseteq I \iff \mathfrak{p} \supseteq \varphi(I).$$

Thus  $f^{-1}(V(I)) = V(\varphi(I))$ .





## Proof.

Construction of  $f^\sharp$ . For any  $\mathfrak{p} \in \operatorname{Spec} B$ , we have a localized map  $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$ , defined by  $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$ .

## Proof.

Construction of  $f^\#$ . For any  $\mathfrak{p} \in \operatorname{Spec} B$ , we have a localized map  $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$ , defined by  $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$ . Given  $V \subseteq \operatorname{Spec}(A)$  open, and  $s \in \mathcal{O}_{\operatorname{Spec}(A)}(V)$  define

$$f^\#(s)(\mathfrak{p}) = \varphi_{\mathfrak{p}} \circ s \circ f.$$

## Proof.

Construction of  $f^\#$ . For any  $\mathfrak{p} \in \operatorname{Spec} B$ , we have a localized map  $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$ , defined by  $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$ . Given  $V \subseteq \operatorname{Spec}(A)$  open, and  $s \in \mathcal{O}_{\operatorname{Spec}(A)}(V)$  define

$$f^\#(s)(\mathfrak{p}) = \varphi_{\mathfrak{p}} \circ s \circ f.$$

Then  $f^\#$  is a ring homomorphism. Furthermore, if  $U \subseteq V$  is such that there exist  $a, b \in A$  with  $U \subseteq D(b)$  such that  $s(\mathfrak{q}) = a/b \in A_{\mathfrak{q}}$  for all  $\mathfrak{q} \in U$ , then  $f^{-1}(U) \subseteq D(\varphi(b))$  and  $f^\#(s)(\mathfrak{q}') = \varphi(a)/\varphi(b) \in B_{\mathfrak{q}'}$  for all  $\mathfrak{q}' \in f^{-1}(U)$ . It follows that  $f^\#(s) \in \mathcal{O}_{\operatorname{Spec}(B)}(f^{-1}(V)) = f_* \mathcal{O}_{\operatorname{Spec}(A)}(V)$ .

## Proof.

- ② Given a morphism of locally ringed spaces  $(f, f^\#)$ , in particular, we get a map  $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$ , where we identify a ring  $R$  with  $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$ .

## Proof.

- ② Given a morphism of locally ringed spaces  $(f, f^\#)$ , in particular, we get a map  $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$ , where we identify a ring  $R$  with  $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$ .  $\varphi$  must agree with  $f^\#$  on stalks, i.e. we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ I_A \downarrow & & \downarrow I_B \\ A_{f(p)} & \xrightarrow[p]{f^\#} & B_p \end{array} .$$

## Proof.

- ② Given a morphism of locally ringed spaces  $(f, f^\#)$ , in particular, we get a map  $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$ , where we identify a ring  $R$  with  $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$ .  $\varphi$  must agree with  $f^\#$  on stalks, i.e. we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ I_A \downarrow & & \downarrow I_B \\ A_{f(p)} & \xrightarrow[p]{f^\#} & B_p \end{array} .$$

In particular, since the bottom row consists of local rings and a local homomorphism, it must be that

$$\varphi^{-1}(\mathfrak{p}) = \varphi^{-1} \circ I_B^{-1}(\mathfrak{p}B_p) = I_A^{-1} \circ f^\#(\mathfrak{p}B_p) = f(\mathfrak{p}).$$

Thus  $f = \varphi^{-1}$ .

## Example

Let  $R = \overline{\mathbb{F}_q}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ .

## Example

Let  $R = \overline{\mathbb{F}_q}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ .  
 $T = \text{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi^{-1}$ .



## Example

Let  $R = \overline{\mathbb{F}_q}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ .  
 $T = \text{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi^{-1}$ .

- The associated map  $f : \text{Spec}(R) \rightarrow \text{Spec}(R)$  is the identity map.

## Example

Let  $R = \overline{\mathbb{F}_q}[[t]]$ , and  $\varphi : R \rightarrow R$  defined by  $\varphi(\sum a_i t^i) = \sum a_i^q t^i$ .  
 $T = \operatorname{Spec}(R)$  has two points with  $\mathfrak{p} = (t)$ , a closed point, and  $(0)$ , which is open and dense. Both points are invariant under  $\varphi^{-1}$ .

- The associated map  $f : \operatorname{Spec}(R) \rightarrow \operatorname{Spec}(R)$  is the identity map.
- Given  $V \subseteq \operatorname{Spec}(R)$  and  $s \in \mathcal{O}_{\operatorname{Spec}(R)}(V)$ , we have

$$f^\#(s)(\cdot) = (s(\cdot))^q.$$

## Corollary (of the proposition)

*Let  $A, B$  be rings. Then  $A$  and  $B$  are isomorphic if and only if  $\operatorname{Spec}(A)$  and  $\operatorname{Spec}(B)$  are isomorphic as locally ringed spaces.*

## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

A *scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  for which  $X$  is covered by open sets  $X = \bigcup U_\alpha$ , such that  $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$  is an affine scheme.

## Definition

An *affine scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  which is isomorphic to  $\mathrm{Spec}(A)$  for some ring  $A$ .

A *scheme* is a locally ringed space  $(X, \mathcal{O}_X)$  for which  $X$  is covered by open sets  $X = \bigcup U_\alpha$ , such that  $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$  is an affine scheme.

Here  $\mathcal{O}_X|_{U_\alpha}$  denotes the restricted sheaf  $\mathcal{O}_X|_{U_\alpha}(V) = \mathcal{O}_X(V)$  for  $V \subseteq U_\alpha$  open.