

Algebraic Geometry 2

Tutorial session 6

Lecturer: Rami Aizenbud
TA: Shai Shechter

June 12, 2020

Functor of points

Definition

A functor $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \underline{\mathbf{Sets}}$ is said to be *representable* if there exists $A \in \underline{\mathbf{C}}$ such that $F = h^A$; i.e. $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$ for all $B \in \underline{\mathbf{C}}$.

Definition

A functor $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \underline{\mathbf{Sets}}$ is said to be *representable* if there exists $A \in \underline{\mathbf{C}}$ such that $F = h^A$; i.e. $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$ for all $B \in \underline{\mathbf{C}}$.

There is also an analogous definition for covariant functors.

Definition

A functor $F : \underline{\mathbf{C}}^{\text{op}} \rightarrow \mathbf{Sets}$ is said to be *representable* if there exists $A \in \underline{\mathbf{C}}$ such that $F = h^A$; i.e. $F(B) \simeq \text{Mor}_{\underline{\mathbf{C}}}(B, A)$ for all $B \in \underline{\mathbf{C}}$.

There is also an analogous definition for covariant functors. There $F : \underline{\mathbf{C}} \rightarrow \mathbf{Set}$ is said to be representable if

$$F = h_A = (B \mapsto \text{Mor}_{\underline{\mathbf{C}}}(A, B)) \text{ for some } A \in \underline{\mathbf{C}}.$$

New definition of (affine) schemes

Given a scheme (X, \mathcal{O}_X) , we can define a functor $\mathbf{Ring} \rightarrow \mathbf{Set}$ by

$$F_X(R) = \mathrm{Hom}_{\mathbf{Sch}}(\mathrm{Spec}(R), X).$$

New definition of (affine) schemes

Given a scheme (X, \mathcal{O}_X) , we can define a functor $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$ by

$$F_X(R) = \mathrm{Hom}_{\underline{\mathbf{Sch}}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of X . By an exercise, a scheme is affine iff F_X is representable; specifically, $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$.

New definition of (affine) schemes

Given a scheme (X, \mathcal{O}_X) , we can define a functor $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$ by

$$F_X(R) = \mathrm{Hom}_{\underline{\mathbf{Sch}}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of X . By an exercise, a scheme is affine iff F_X is representable; specifically, $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$.

Definition (New definition of affine scheme)

An affine scheme is a representable functor $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$.

New definition of (affine) schemes

Given a scheme (X, \mathcal{O}_X) , we can define a functor $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$ by

$$F_X(R) = \mathrm{Hom}_{\underline{\mathbf{Sch}}}(\mathrm{Spec}(R), X).$$

This is called the *functor of points* of X . By an exercise, a scheme is affine iff F_X is representable; specifically, $F_X(\cdot) = \mathrm{Hom}(\mathcal{O}_X(X), \cdot)$.

Definition (New definition of affine scheme)

An affine scheme is a representable functor $\underline{\mathbf{Ring}} \rightarrow \underline{\mathbf{Set}}$.

A general scheme, in this setting, would be a functor which is “locally representable”.

Example

Consider the ring $A = \mathbb{Z}[t, t^{-1}]$.

Example

Consider the ring $A = \mathbb{Z}[t, t^{-1}]$. Then $\operatorname{Spec}(A) = \operatorname{Spec}(\mathbb{Z}[t]) \setminus V(t)$, which is, a-priori, not easy to describe.

Example

Consider the ring $A = \mathbb{Z}[t, t^{-1}]$. Then $\operatorname{Spec}(A) = \operatorname{Spec}(\mathbb{Z}[t]) \setminus V(t)$, which is, a-priori, not easy to describe.

The associated functor of points, on the other hand, is very easy. Namely-

$$F_{\operatorname{Spec}(A)}(R) = \operatorname{Hom}_{\underline{\mathbf{Rings}}}(\mathbb{Z}[t, t^{-1}], R) = R^{\times},$$

since choosing a homomorphism $\mathbb{Z}[t, t^{-1}] \rightarrow R$ amounts to choosing the image of t , which is necessarily in R^{\times} .

Example

Consider the ring $A = \mathbb{Z}[t, t^{-1}]$. Then $\mathrm{Spec}(A) = \mathrm{Spec}(\mathbb{Z}[t]) \setminus V(t)$, which is, a-priori, not easy to describe.

The associated functor of points, on the other hand, is very easy. Namely-


$$F_{\mathrm{Spec}(A)}(R) = \mathrm{Hom}_{\mathbf{Rings}}(\mathbb{Z}[t, t^{-1}], R) = R^{\times},$$

since choosing a homomorphism $\mathbb{Z}[t, t^{-1}] \rightarrow R$ amounts to choosing the image of t , which is necessarily in R^{\times} .

In fact, we can also encode the group laws of R^{\times} in terms natural transformations of $F_{\mathrm{Spec}(A)}$, giving rise to $F_{\mathrm{Spec}(A)}$ as a *group scheme*. For example, the inversion in R^{\times} is “encoded” in the map

$$t \mapsto t^{-1} : A \rightarrow A.$$

A very nice example is Mumford's doodle of $\mathrm{Spec}(\mathbb{Z}[t])$:



See <http://www.neverendingbooks.org/grothendiecks-functor-of-points> for more information.

Products and fibered products

Exercise

Let k be an algebraically closed field.

Exercise

Let k be an algebraically closed field.

- 1 Show that the map of sets, $\mathbb{A}_k^1 \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^2$, defined by $((x - a), (y - b)) \mapsto (x - a, y - b)$ is not surjective.

Exercise

Let k be an algebraically closed field.

- 1 Show that the map of sets, $\mathbb{A}_k^1 \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^2$, defined by $((x - a), (y - b)) \mapsto (x - a, y - b)$ is not surjective.
- 2 Show that it *is* a bijection of closed points.

Exercise

Let k be an algebraically closed field.

- 1 Show that the map of sets, $\mathbb{A}_k^1 \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^2$, defined by $((x - a), (y - b)) \mapsto (x - a, y - b)$ is not surjective.
- 2 Show that it *is* a bijection of closed points.

Solution.

- 2 By Nullstellensatz, any proper ideal I of $k[x, y]$ admits a point $(a, b) \in k^2$ such that $f(a, b) = 0$ for all $f \in I$.

Exercise

Let k be an algebraically closed field.

- 1 Show that the map of sets, $\mathbb{A}_k^1 \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^2$, defined by $((x - a), (y - b)) \mapsto (x - a, y - b)$ is not surjective.
- 2 Show that it *is* a bijection of closed points.

Solution.

- 2 By Nullstellensatz, any proper ideal I of $k[x, y]$ admits a point $(a, b) \in k^2$ such that $f(a, b) = 0$ for all $f \in I$. In particular, $I \subseteq (x - a, y - b)$, and any maximal ideal of $k[x, y]$ is of the form $(x - a, y - b)$.



Exercise

Let k be an algebraically closed field.

- 1 Show that the map of sets, $\mathbb{A}_k^1 \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^2$, defined by $((x - a), (y - b)) \mapsto (x - a, y - b)$ is not surjective.
- 2 Show that it *is* a bijection of closed points.

Solution.

- 2 By Nullstellensatz, any proper ideal I of $k[x, y]$ admits a point $(a, b) \in k^2$ such that $f(a, b) = 0$ for all $f \in I$. In particular, $I \subseteq (x - a, y - b)$, and any maximal ideal of $k[x, y]$ is of the form $(x - a, y - b)$.
- 1 For example, the ideal (x) is prime and not in the image of this map.



The last exercise is intended in order to stress that, unlike classical AG, where the difference between the cartesian product and the product in the category of varieties was manifested in the topology, in the case of schemes, the two are different even as sets.

The last exercise is intended in order to stress that, unlike classical AG, where the difference between the cartesian product and the product in the category of varieties was manifested in the topology, in the case of schemes, the two are different even as sets.


On the other hand, taking the functor of points approach, we will have that, for example, $\mathbb{A}^1(R) \times \mathbb{A}^1(R) \simeq \mathbb{A}^2(R)$ for any ring R .

Definition

Let $\underline{\mathbf{C}}$ be a category and X, Y, Z objects with morphisms $X \rightarrow Y$ and $Z \rightarrow Y$.

Definition

Let $\underline{\mathbf{C}}$ be a category and X, Y, Z objects with morphisms $X \rightarrow Y$ and $Z \rightarrow Y$. A fibered product $X \times_Y Z$ is an object of $\underline{\mathbf{C}}$ along with morphisms to X and Z such that the universal property described by the following diagram holds.



The fibered product

Exercise


Show that, if Y is a terminal object in $\underline{\mathbf{C}}$, then $X \times_Y Z \simeq X \times Y$.

The fibered product

Exercise

Show that, if Y is a terminal object in $\underline{\mathbf{C}}$, then $X \times_Y Z \simeq X \times Z$.

Solution.




The fibered product

Exercise

Show that, if Y is a terminal object in $\underline{\mathbf{C}}$, then $X \times_Y Z \simeq X \times Y$.


Solution.



- 1 The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.

The fibered product


Solution.



- ① The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.

The fibered product


Solution.



- 1 The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.
- 2 Since Y is terminal, we have that $f \circ \pi_X = g \circ \pi_Z$ and both equal the unique morphism $X \times Z \rightarrow Y$.

The fibered product


Solution.



- 1 The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.
- 2 Since Y is terminal, we have that $f \circ \pi_X = g \circ \pi_Z$ and both equal the unique morphism $X \times Z \rightarrow Y$. Get $X \times_Y Z \xrightarrow{\psi} X \times Y$.

The fibered product


Solution.



- 1 The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.
- 2 Since Y is terminal, we have that $f \circ \pi_X = g \circ \pi_Z$ and both equal the unique morphism $X \times Z \rightarrow Y$. Get $X \times_Y Z \xrightarrow{\psi} X \times Y$.
- 3 $\varphi \circ \psi$ is the unique map $X \times Z$ which commutes with the projection maps;

The fibered product


Solution.



- ① The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.
- ② Since Y is terminal, we have that $f \circ \pi_X = g \circ \pi_Z$ and both equal the unique morphism $X \times Z \rightarrow Y$. Get $X \times_Y Z \xrightarrow{\psi} X \times Y$.
- ③ $\varphi \circ \psi$ is the unique map $X \times Z$ which commutes with the projection maps; i.e. it is $\mathbf{1}_{X \times Z}$.

The fibered product

Solution.



- 1 The map $X \times_Y Z \xrightarrow{\varphi} X \times Z$ always exists, by UP of product.
- 2 Since Y is terminal, we have that $f \circ \pi_X = g \circ \pi_Z$ and both equal the unique morphism $X \times Z \rightarrow Y$. Get $X \times_Y Z \xrightarrow{\psi} X \times Y$.
- 3 $\varphi \circ \psi$ is the unique map $X \times Z$ which commutes with the projection maps; i.e. it is $\mathbf{1}_{X \times Z}$. Similarly, $\psi \circ \varphi = \mathbf{1}_{X \times_Y Z}$.

Fibered products

Theorem

Fibered products exist in the category of schemes.

Fibered products

Theorem

Fibered products exist in the category of schemes.

In the case where X , Y and Z are all affine, the existence of fibered product is easy to prove.

Fibered products

Theorem

Fibered products exist in the category of schemes.

In the case where X , Y and Z are all affine, the existence of fibered product is easy to prove. Essentially, it follows from the UP of the tensor product. In particular

$$\mathrm{Spec}(A) \times_{\mathrm{Spec}(B)} \mathrm{Spec}(C) \simeq \mathrm{Spec}(A \otimes_B C),$$

for A and C algebras over B .

Example

$\mathbb{A}_k^1(R) \times \mathbb{A}_k^1(R) \simeq \mathbb{A}_k^2(R)$ for any ring k and k -algebra R .

Indeed, we have that

$$\begin{aligned}\mathbb{A}^2(R) &= \operatorname{Hom}_k(k[x, y], R) \simeq \operatorname{Hom}_k(k[x] \otimes_k k[y], R) \\ &\simeq \operatorname{Hom}(k[x], R) \times \operatorname{Hom}(k[y], R) = \mathbb{A}_k^1(R) \times \mathbb{A}_k^1(R)\end{aligned}$$

where the second-to-last isomorphism is the UP of \otimes .

Some facts about tensor products

Let $\varphi : A \rightarrow B$ be a ring homomorphism.

Some facts about tensor products

Let $\varphi : A \rightarrow B$ be a ring homomorphism.

- 1 Given an ideal $I \triangleleft A$, we have a natural isomorphism $(A/I) \otimes_A B \simeq B/(\varphi(I))$.

Some facts about tensor products

Let $\varphi : A \rightarrow B$ be a ring homomorphism.

- 1 Given an ideal $I \triangleleft A$, we have a natural isomorphism $(A/I) \otimes_A B \simeq B/(\varphi(I))$.
- 2 Given a multiplicatively closed subset $S \subseteq A$, we have a natural isomorphism $(S^{-1}A) \otimes_A B \simeq \varphi(S)^{-1}B$.

Proof of item (1)

Key tool

Tensor products are right-exact.

Proof of item (1)

Key tool

Tensor products are right-exact.

Proof.

Given a ses of modules $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$ and another module M consider

$$A \otimes M \xrightarrow{\alpha \otimes 1_M} B \otimes M \xrightarrow{\beta \otimes 1_M} C \otimes M \rightarrow 0.$$

Proof of item (1)

Key tool

Tensor products are right-exact.

Proof.

Given a ses of modules $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$ and another module M consider

$$A \otimes M \xrightarrow{\alpha \otimes 1_M} B \otimes M \xrightarrow{\beta \otimes 1_M} C \otimes M \rightarrow 0.$$

Proof of item (1)

Key tool

Tensor products are right-exact.

Proof.

Given a ses of modules $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$ and another module M consider

$$A \otimes M \xrightarrow{\alpha \otimes \mathbf{1}_M} B \otimes M \xrightarrow{\beta \otimes \mathbf{1}_M} C \otimes M \rightarrow 0.$$

Showing $\beta \otimes \mathbf{1}_M$ is surjective and $\text{Im}(\alpha \otimes \mathbf{1}_M) \subseteq \text{Ker}(\beta \otimes \mathbf{1}_M)$ is easy.

Proof of item (1)

Key tool

Tensor products are right-exact.

Proof.

Given a ses of modules $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$ and another module M consider

$$A \otimes M \xrightarrow{\alpha \otimes \mathbf{1}_M} B \otimes M \xrightarrow{\beta \otimes \mathbf{1}_M} C \otimes M \rightarrow 0.$$

Showing $\beta \otimes \mathbf{1}_M$ is surjective and $\text{Im}(\alpha \otimes \mathbf{1}_M) \subseteq \text{Ker}(\beta \otimes \mathbf{1}_M)$ is easy. Setting $I = \text{Im}(\alpha \otimes \mathbf{1}_M)$, we get a map

$$B \otimes M / I \rightarrow C \otimes M.$$

Check that $(c, m) \mapsto \beta^{-1}(c) \otimes m + I$ is a well-defined inverse. □

We can also prove exactness using the more general fact that $\cdot \otimes M$ has a left adjoint (namely- $\text{hom}(M, \cdot)$), hence is right exact. This type of proof is neater, but beyond the scope of this tutorial.

Proof of item (1) - contd

We can apply this to the short exact sequence (of A -modules)

$$I \rightarrow A \rightarrow A/I \rightarrow 0$$

and with the functor $\cdot \otimes_A B$, we get

$$\begin{array}{ccccccc} I \otimes B & \longrightarrow & A \otimes B & \longrightarrow & (A/I) \otimes B & \longrightarrow & 0 \\ \parallel \wr & & \parallel \wr & & \vdots \wr & & \\ \varphi(I)B & \longrightarrow & B & \longrightarrow & B/\varphi(I)B & \longrightarrow & 0 \end{array}$$

where the first two columns are the multiplication map.

Proof of item (2)

We have a natural map

$$S^{-1}A \otimes_A B \rightarrow \varphi(S)^{-1}B$$

given on generators by

$$\frac{a}{s} \otimes b \mapsto \frac{\varphi(a)b}{\varphi(s)}.$$

Proof of item (2)

We have a natural map

$$S^{-1}A \otimes_A B \rightarrow \varphi(S)^{-1}B$$

given on generators by

$$\frac{a}{s} \otimes b \mapsto \frac{\varphi(a)b}{\varphi(s)}.$$

Surjectivity is clear, because the image contains all fractions of the form $\frac{b}{\varphi(s)}$ for $b \in B$ and $s \in S$.

Proof of item (2) - contd

For injectivity, first consider the case of a simple tensor $\frac{a}{s} \otimes b$.

Proof of item (2) - contd

For injectivity, first consider the case of a simple tensor $\frac{a}{s} \otimes b$. If $\frac{\varphi(a)b}{\varphi(s)} = 0$ in $\varphi(S)^{-1}B$ then $\exists t = \varphi(t') \in \varphi(S)$ such that $t\varphi(a)b = 0$, or, equivalently, such that $(t'a) \otimes b$ is in the kernel of the map $A \otimes_A B \rightarrow B$.

Proof of item (2) - contd

For injectivity, first consider the case of a simple tensor $\frac{a}{s} \otimes b$. If $\frac{\varphi(a)b}{\varphi(s)} = 0$ in $\varphi(S)^{-1}B$ then $\exists t = \varphi(t') \in \varphi(S)$ such that $t\varphi(a)b = 0$, or, equivalently, such that $(t'a) \otimes b$ is in the kernel of the map $A \otimes_A B \rightarrow B$. But this map is an isomorphism, hence $(t'a) \otimes b = 0$, hence $\frac{a}{s} \otimes b = 0$ in $(S^{-1}A) \otimes B$.

Proof of item (2) - contd

For injectivity, first consider the case of a simple tensor $\frac{a}{s} \otimes b$. If $\frac{\varphi(a)b}{\varphi(s)} = 0$ in $\varphi(S)^{-1}B$ then $\exists t = \varphi(t') \in \varphi(S)$ such that $t\varphi(a)b = 0$, or, equivalently, such that $(t'a) \otimes b$ is in the kernel of the map $A \otimes_A B \rightarrow B$. But this map is an isomorphism, hence $(t'a) \otimes b = 0$, hence $\frac{a}{s} \otimes b = 0$ in $(S^{-1}A) \otimes B$.

The general case is similar- if $\sum_i \frac{a_i}{s_i} \otimes b_i$ is mapped to zero, then

$$\sum_i \frac{\varphi(a_i)b_i}{\varphi(s_i)} = \frac{\overbrace{\sum_i \varphi(a_i) \cdot \left(\prod_{j \neq i} \varphi(s_j)\right)}^{=\varphi(a'_i)} \cdot b_i}{\prod_i \varphi(s_i)} = 0.$$

Put $a'_i = a_i \sum_{j \neq i} s_j$. Then, for some $t' \in S$, we have

$$t' \left(\sum_i a'_i \otimes b_i \right) \in \text{Ker}(A \otimes B \rightarrow B)$$

Corollary

All examples of morphisms of affine schemes discussed in the previous tutorial are preserved under base change.

Corollary

All examples of morphisms of affine schemes discussed in the previous tutorial are preserved under base change. More generally- closed embeddings, open embeddings and localizations of affine schemes are preserved under base change.

Models of schemes

Let X be a \mathbb{Q} -scheme, i.e., a scheme with a morphism $X \rightarrow \operatorname{Spec}(\mathbb{Q})$.

Definition

A \mathbb{Z} -model for X is a scheme X' (over \mathbb{Z}) such that $X' \times_{\mathbb{Z}} \mathbb{Q} \simeq X$.

Models of schemes

Let X be a \mathbb{Q} -scheme, i.e, a scheme with a morphism $X \rightarrow \operatorname{Spec}(\mathbb{Q})$.

Definition

A \mathbb{Z} -model for X is a scheme X' (over \mathbb{Z}) such that $X' \times_{\mathbb{Z}} \mathbb{Q} \simeq X$.

Exercise

Assume X is affine. Show that, by composing $X \rightarrow \operatorname{Spec}(\mathbb{Q})$ with the map $\operatorname{Spec}(\mathbb{Q}) \rightarrow \operatorname{Spec}(\mathbb{Z})$, we get a \mathbb{Z} -model for X .

Models of schemes

Let X be a \mathbb{Q} -scheme, i.e., a scheme with a morphism $X \rightarrow \operatorname{Spec}(\mathbb{Q})$.

Definition

A \mathbb{Z} -model for X is a scheme X' (over \mathbb{Z}) such that $X' \times_{\mathbb{Z}} \mathbb{Q} \simeq X$.

Exercise

Assume X is affine. Show that, by composing $X \rightarrow \operatorname{Spec}(\mathbb{Q})$ with the map $\operatorname{Spec}(\mathbb{Q}) \rightarrow \operatorname{Spec}(\mathbb{Z})$, we get a \mathbb{Z} -model for X . Does this generalize to any morphism of rings $R \rightarrow S$? Any inclusion?

Models for schemes

Solution.

Let $A = \Gamma(X, \mathcal{O}_X)$. The claim that $X \rightarrow \operatorname{Spec}(\mathbb{Z})$ is a \mathbb{Z} -model is equivalent to the following:

Lemma

Let A be a \mathbb{Q} -algebra, and let A' be its underlying ring (i.e. A , considered as a \mathbb{Z} -algebra). Then $A' \otimes_{\mathbb{Z}} \mathbb{Q} \simeq A$.

Models for schemes

Solution.

Let $A = \Gamma(X, \mathcal{O}_X)$. The claim that $X \rightarrow \operatorname{Spec}(\mathbb{Z})$ is a \mathbb{Z} -model is equivalent to the following:

Lemma

Let A be a \mathbb{Q} -algebra, and let A' be its underlying ring (i.e. A , considered as a \mathbb{Z} -algebra). Then $A' \otimes_{\mathbb{Z}} \mathbb{Q} \simeq A$.

The lemma holds because $\mathbb{Q} = S^{-1}\mathbb{Z}$, for $S = \mathbb{Z} \setminus \{0\}$, and by a previous exercise:

$$A' \otimes_{\mathbb{Z}} \mathbb{Q} = A' \otimes_{\mathbb{Z}} S^{-1}\mathbb{Z} = S^{-1}A' = A,$$

since A' is already closed under multiplication by \mathbb{Q} . □

Models for schemes

Solution– contd.

Regarding generalizations of this statement for a ring homomorphism $R \rightarrow S$, and the case where we have an S -scheme X and we seek an R -model, the same argument will hold in the cases where:

Models for schemes

Solution– contd.

Regarding generalizations of this statement for a ring homomorphism $R \rightarrow S$, and the case where we have an S -scheme X and we seek an R -model, the same argument will hold in the cases where:

- 1 S is a localization of R by some multiplicatively closed set, and the homomorphism is the natural map $R \rightarrow S$; or

Models for schemes

Solution– contd.

Regarding generalizations of this statement for a ring homomorphism $R \rightarrow S$, and the case where we have an S -scheme X and we seek an R -model, the same argument will hold in the cases where:

- 1 S is a localization of R by some multiplicatively closed set, and the homomorphism is the natural map $R \rightarrow S$; or
- 2 $S = R/I$ for some ideal $I \triangleleft R$.

Models for schemes

Solution– contd.

Regarding generalizations of this statement for a ring homomorphism $R \rightarrow S$, and the case where we have an S -scheme X and we seek an R -model, the same argument will hold in the cases where:

- ① S is a localization of R by some multiplicatively closed set, and the homomorphism is the natural map $R \rightarrow S$; or
- ② $S = R/I$ for some ideal $I \triangleleft R$.

The argument **does not** hold for arbitrary inclusions.



Example

Let $X = \operatorname{Spec}(\mathbb{C})$, considered as a scheme over \mathbb{C} . Then X is not an \mathbb{R} -model of itself.

Example

Let $X = \operatorname{Spec}(\mathbb{C})$, considered as a scheme over \mathbb{C} . Then X is not an \mathbb{R} -model of itself.

Indeed,

$$X \times_{\operatorname{Spec}(\mathbb{R})} \operatorname{Spec}(\mathbb{C}) = \operatorname{Spec}(\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}) = \operatorname{Spec}(\mathbb{C} \times \mathbb{C}),$$

where the last equality holds since

$$\begin{aligned}\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} &\simeq \mathbb{C} \otimes_{\mathbb{R}} (\mathbb{R}[t]/(t^2 + 1)) \\ &= \mathbb{C}[t]/(t^2 + 1) \\ &\simeq (\mathbb{C}[t]/(t - i)) \times (\mathbb{C}[t]/(t + i)) = \mathbb{C} \times \mathbb{C}.\end{aligned}$$

Thus $X \times_{\operatorname{Spec}(\mathbb{R})} \operatorname{Spec}(\mathbb{C})$ has two points, while X has only one.

Definition (The fiber of a morphism)

Let $X \rightarrow Y$ be a morphism of schemes, and $y \in Y$ a point. The fiber of f at y is defined by the fibered product

$$\begin{array}{ccc} X \times_Y \operatorname{Spec}(k(y)) & \longrightarrow & X \\ \downarrow & & \downarrow \\ \operatorname{Spec}(k(y)) & \longrightarrow & Y \end{array}$$

where $\operatorname{Spec}(k(y)) \rightarrow Y$ is the canonical map with image $\{y\}$.

Exercise

Compute the fibers of $\mathrm{Spec}(\mathbb{Q}[x, y]/(y^2 - x)) \rightarrow \mathbb{A}_{\mathbb{Q}}^1$ at the maximal ideals $(x - 1)$, (x) and $(x + 1)$. Also at (0) .

Exercise

Compute the fibers of $\mathrm{Spec}(\mathbb{Q}[x, y]/(y^2 - x)) \rightarrow \mathbb{A}_{\mathbb{Q}}^1$ at the maximal ideals $(x - 1)$, (x) and $(x + 1)$. Also at (0) .

- 1 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x - 1) \simeq \mathbb{Q}[y]/(y^2 - 1) \simeq \mathbb{Q} \times \mathbb{Q}$,
and the spectrum has two points.

Exercise

Compute the fibers of $\mathrm{Spec}(\mathbb{Q}[x, y]/(y^2 - x)) \rightarrow \mathbb{A}_{\mathbb{Q}}^1$ at the maximal ideals $(x - 1)$, (x) and $(x + 1)$. Also at (0) .

- 1 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x - 1) \simeq \mathbb{Q}[y]/(y^2 - 1) \simeq \mathbb{Q} \times \mathbb{Q}$, and the spectrum has two points.
- 2 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x) \simeq \mathbb{Q}[y]/(y^2)$, and the spectrum has one *double* point.

Exercise

Compute the fibers of $\mathrm{Spec}(\mathbb{Q}[x, y]/(y^2 - x)) \rightarrow \mathbb{A}_{\mathbb{Q}}^1$ at the maximal ideals $(x - 1)$, (x) and $(x + 1)$. Also at (0) .

- 1 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x - 1) \simeq \mathbb{Q}[y]/(y^2 - 1) \simeq \mathbb{Q} \times \mathbb{Q}$, and the spectrum has two points.
- 2 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x) \simeq \mathbb{Q}[y]/(y^2)$, and the spectrum has one *double* point.
- 3 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x + 1) \simeq \mathbb{Q}[y]/(y^2 + 1) \simeq \mathbb{Q}[i]$, and the spectrum has one point, but with a larger field.

Exercise

Compute the fibers of $\mathrm{Spec}(\mathbb{Q}[x, y]/(y^2 - x)) \rightarrow \mathbb{A}_{\mathbb{Q}}^1$ at the maximal ideals $(x - 1)$, (x) and $(x + 1)$. Also at (0) .

- 1 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x - 1) \simeq \mathbb{Q}[y]/(y^2 - 1) \simeq \mathbb{Q} \times \mathbb{Q}$, and the spectrum has two points.
- 2 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x) \simeq \mathbb{Q}[y]/(y^2)$, and the spectrum has one *double* point.
- 3 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}[x]/(x + 1) \simeq \mathbb{Q}[y]/(y^2 + 1) \simeq \mathbb{Q}[i]$, and the spectrum has one point, but with a larger field.
- 4 $\mathbb{Q}[x, y]/(y^2 - x) \otimes_{\mathbb{Q}[x]} \mathbb{Q}(x) \simeq \mathbb{Q}[y] \otimes \mathbb{Q}(y^2) \simeq \mathbb{Q}(y)$

Questions?