

Algebraic Geometry 2

Tutorial session 4

Lecturer: Rami Aizenbud
TA: Shai Shechter

May 17, 2020

The pullback and direct image - recollections

Let $f : X \rightarrow Y$ be a continuous map of topological space, and let \mathcal{F} be a sheaf on X and \mathcal{G} a sheaf on Y .

The pullback and direct image - recollections

Let $f : X \rightarrow Y$ be a continuous map of topological space, and let \mathcal{F} be a sheaf on X and \mathcal{G} a sheaf on Y . Recall that we defined the pullback of \mathcal{F} to be the sheaf $f^*\mathcal{F}$ on Y , defined by

$$f^*\mathcal{F}(V) = \mathcal{F}(f^{-1}(V)) \quad \text{for } V \subseteq Y \text{ open.}$$

The pullback and direct image - recollections

Let $f : X \rightarrow Y$ be a continuous map of topological space, and let \mathcal{F} be a sheaf on X and \mathcal{G} a sheaf on Y . Recall that we defined the pullback of \mathcal{F} to be the sheaf $f_*\mathcal{F}$ on Y , defined by

$$f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}(V)) \quad \text{for } V \subseteq Y \text{ open.}$$

We also defined the direct image sheaf $f^{-1}\mathcal{G}$ on X , by sheafification of the presheaf defined by

$$f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U) \text{ open}} \mathcal{G}(V) \quad \text{for } U \subseteq X \text{ open.}$$

Adjointness of pullback and direct image

Exercise

Let $f : X \rightarrow Y$ be a continuous map, \mathcal{F} a sheaf on X and \mathcal{G} a sheaf on Y . Then there exists a natural bijection

$$\mathrm{Hom}_{\mathbf{Sh}(X)}(f^{-1}\mathcal{G}, \mathcal{F}) \simeq \mathrm{Hom}_{\mathbf{Sh}(Y)}(\mathcal{G}, f_*\mathcal{F}).$$

Adjontness of pullback and direct image

Exercise

Let $f : X \rightarrow Y$ be a continuous map, \mathcal{F} a sheaf on X and \mathcal{G} a sheaf on Y . Then there exists a natural bijection

$$\mathrm{Hom}_{\underline{\mathbf{Sh}}(X)}(f^{-1}\mathcal{G}, \mathcal{F}) \simeq \mathrm{Hom}_{\underline{\mathbf{Sh}}(Y)}(\mathcal{G}, f_*\mathcal{F}).$$

Proof.

We construct natural maps going in both directions, and verify that their compositions are equivalent to identity (some details are left as exercises).

Proof.

- $F : \operatorname{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \operatorname{Hom}(\mathcal{G}, f_*\mathcal{F})$:

Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$: Let $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$. We construct $F(\varphi)_V$ for V in a basis for the topology on Y . Recall from exercise, that Y has a basis of subsets V such that for $U \subseteq f^{-1}(V) \subseteq X$ open, we have $f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U)} \mathcal{G}(V)$.

Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$: Let $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$. We construct $F(\varphi)_V$ for V in a basis for the topology on Y . Recall from exercise, that Y has a basis of subsets V such that for $U \subseteq f^{-1}(V) \subseteq X$ open, we have $f^{-1}\mathcal{G}(U) = \lim_{V \supseteq f(U)} \mathcal{G}(V)$. Let $V \subseteq Y$ be such, and $U = f^{-1}(V)$.

Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$: Let $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$. We construct $F(\varphi)_V$ for V in a basis for the topology on Y . Recall from exercise, that Y has a basis of subsets V such that for $U \subseteq f^{-1}(V) \subseteq X$ open, we have $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$. Let $V \subseteq Y$ be such, and $U = f^{-1}(V)$. By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$: Let $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$. We construct $F(\varphi)_V$ for V in a basis for the topology on Y . Recall from exercise, that Y has a basis of subsets V such that for $U \subseteq f^{-1}(V) \subseteq X$ open, we have $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$. Let $V \subseteq Y$ be such, and $U = f^{-1}(V)$. By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

In particular, $\text{Hom}(f^{-1}\mathcal{G}(U), \mathcal{F}(U)) = \text{Hom}(\mathcal{G}(V), f_*\mathcal{F}(V))$, and we can define $F(\varphi)_V = \varphi_{f^{-1}(V)}$.

Proof.

- $F : \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{G}, f_*\mathcal{F})$: Let $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$. We construct $F(\varphi)_V$ for V in a basis for the topology on Y . Recall from exercise, that Y has a basis of subsets V such that for $U \subseteq f^{-1}(V) \subseteq X$ open, we have $f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V')$. Let $V \subseteq Y$ be such, and $U = f^{-1}(V)$. By definition

$$f^{-1}\mathcal{G}(U) = \lim_{V' \supseteq f(U)} \mathcal{G}(V') = \mathcal{G}(V) \quad \text{and} \quad \mathcal{F}(U) = f_*\mathcal{F}(V).$$

In particular, $\text{Hom}(f^{-1}\mathcal{G}(U), \mathcal{F}(U)) = \text{Hom}(\mathcal{G}(V), f_*\mathcal{F}(V))$, and we can define $F(\varphi)_V = \varphi_{f^{-1}(V)}$. This gives rise to a sheaf morphism $F(\varphi) \in \text{Hom}(\mathcal{G}, f_*\mathcal{F})$ (Ex).



Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$: Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$,
 $U \subseteq X$ open and $f(U) \subseteq V \subseteq Y$ open, we have a map
 $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$, given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$: Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$, $U \subseteq X$ open and $f(U) \subseteq V \subseteq Y$ open, we have a map $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$, given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative.

Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$: Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$, $U \subseteq X$ open and $f(U) \subseteq V \subseteq Y$ open, we have a map $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$, given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative. The maps $(g_U)_{U \subseteq X \text{ open}}$ comprise a morphism of presheaves from the presheaf $(U \mapsto \lim_{V \supseteq f(U)} \mathcal{G}(V))$ to \mathcal{F} (Ex).

Proof.

- $G : \text{Hom}(\mathcal{G}, f_*\mathcal{F}) \rightarrow \text{Hom}(f^{-1}\mathcal{G}, \mathcal{F})$: Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$, $U \subseteq X$ open and $f(U) \subseteq V \subseteq Y$ open, we have a map $g_{V,U} : \mathcal{G}(V) \rightarrow \mathcal{F}(U)$, given by the composition

$$\mathcal{G}(V) \xrightarrow{\psi_V} f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}V) \xrightarrow{\text{res}_{f^{-1}(V),U}} \mathcal{F}(U).$$

The universal property of direct limit then gives a map

$$g_U = \lim_{V \supseteq f(U)} g_{V,U} : \lim_{V \supseteq f(U)} \mathcal{G}(V) \rightarrow \mathcal{F}(U),$$

which makes the suitable diagram commutative. The maps $(g_U)_{U \subseteq X \text{ open}}$ comprise a morphism of presheaves from the presheaf $(U \mapsto \lim_{V \supseteq f(U)} \mathcal{G}(V))$ to \mathcal{F} (Ex). Sheafifying, we set $G(\psi) = \tilde{g} : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$.



Proof.

We need to show $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$ and $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$.

- Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ and $V \subseteq Y$ open, we have $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$, where the RHS is given by a direct limit over open sets containing $f(f^{-1}(V)) = V$.

Proof.

We need to show $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$ and $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$.

- Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ and $V \subseteq Y$ open, we have $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$, where the RHS is given by a direct limit over open sets containing $f(f^{-1}(V)) = V$. Hence it is a filtered limit with a terminal object, and equals to its value at V .

Proof.

We need to show $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$ and $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$.

- Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ and $V \subseteq Y$ open, we have $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$, where the RHS is given by a direct limit over open sets containing $f(f^{-1}(V)) = V$. Hence it is a filtered limit with a terminal object, and equals to its value at V . It follows that $(FG\psi)_V = (G\psi)_{f^{-1}(V)} = \psi_V$.



Proof.

We need to show $FG \sim \mathbf{1}_{\mathrm{Hom}(\mathcal{G}, f_*\mathcal{F})}$ and $GF \sim \mathbf{1}_{\mathrm{Hom}(f^{-1}\mathcal{G}, \mathcal{F})}$.

- Given $\psi : \mathcal{G} \rightarrow f_*\mathcal{F}$ and $V \subseteq Y$ open, we have $(FG\psi)_V = (G\psi)_{f^{-1}(V)}$, where the RHS is given by a direct limit over open sets containing $f(f^{-1}(V)) = V$. Hence it is a filtered limit with a terminal object, and equals to its value at V . It follows that $(FG\psi)_V = (G\psi)_{f^{-1}(V)} = \psi_V$.
- The equivalence $(GF\varphi)_U = \varphi_U$ for all $\varphi : f^{-1}\mathcal{G} \rightarrow \mathcal{F}$ follows similarly, by unfolding the definitions (Ex).



Schemes

The spectrum of a ring

Let A be a commutative unital ring.

Definition

The spectrum of R is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

The spectrum of a ring

Let A be a commutative unital ring.

Definition

The spectrum of R is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

Examples

- 1 $\operatorname{Spec}(k) = \{*\}$ for any field k .

The spectrum of a ring

Let A be a commutative unital ring.

Definition

The spectrum of R is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

Examples

- 1 $\operatorname{Spec}(k) = \{*\}$ for any field k .
- 2 $\operatorname{Spec}(k[x]) =: \mathbb{A}_k^1 \approx \{\text{irreducible monic polynomials in } x\} \sqcup \{0\}$

The spectrum of a ring

Let A be a commutative unital ring.

Definition

The spectrum of R is the set

$$\operatorname{Spec}(A) = \{\mathfrak{p} \triangleleft A : \mathfrak{p} \text{ prime}\}.$$

Examples

- 1 $\operatorname{Spec}(k) = \{*\}$ for any field k .
- 2 $\operatorname{Spec}(k[x]) =: \mathbb{A}_k^1 \approx \{\text{irreducible monic polynomials in } x\} \sqcup \{0\}$
- 3 $\operatorname{Spec}(\mathbb{Z}) = \{\langle 0 \rangle\} \sqcup \{\langle p \rangle : p \text{ prime}\}.$

The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

- 1 $V((0)) = A$ and $V(A) = \emptyset$.

The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

- 1 $V((0)) = A$ and $V(A) = \emptyset$.
- 2 $V(IJ) = V(I) \cup V(J)$.

The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

- 1 $V((0)) = A$ and $V(A) = \emptyset$.
- 2 $V(IJ) = V(I) \cup V(J)$.
- 3 Given a collection $\{I_\alpha\}$ of ideals, $V(\sum I_\alpha) = \bigcap V(I_\alpha)$.

The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \operatorname{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

- 1 $V((0)) = A$ and $V(A) = \emptyset$.
- 2 $V(IJ) = V(I) \cup V(J)$.
- 3 Given a collection $\{I_\alpha\}$ of ideals, $V(\sum I_\alpha) = \bigcap V(I_\alpha)$.

Proof.

Home exercise.



The spectrum of a ring - topology

Given $I \triangleleft A$, define $V(I) := \{\mathfrak{p} \in \text{Spec}(A) : I \subseteq \mathfrak{p}\}$.

Exercise

- 1 $V((0)) = A$ and $V(A) = \emptyset$.
- 2 $V(IJ) = V(I) \cup V(J)$.
- 3 Given a collection $\{I_\alpha\}$ of ideals, $V(\sum I_\alpha) = \bigcap V(I_\alpha)$.

Proof.

Home exercise.



The collection $\{V(I) : I \triangleleft R\}$ is the set of closed sets for a topology on $\text{Spec}(R)$, which is known as the *Zariski Topology* of R .

Exercise

Let A be a ring.

- 1 Show that $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$, for all $\mathfrak{p} \in \operatorname{Spec}(A)$ and, in particular, that $\{\mathfrak{p}\}$ is closed iff \mathfrak{p} is maximal.
- 2 Show that, if A is a domain, then $\{(0)\}$ is a dense point.

Exercise

Let A be a ring.

- 1 Show that $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$, for all $\mathfrak{p} \in \operatorname{Spec}(A)$ and, in particular, that $\{\mathfrak{p}\}$ is closed iff \mathfrak{p} is maximal.
- 2 Show that, if A is a domain, then $\{(0)\}$ is a dense point.

Solution.



Exercise

Let A be a ring.

- 1 Show that $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$, for all $\mathfrak{p} \in \operatorname{Spec}(A)$ and, in particular, that $\{\mathfrak{p}\}$ is closed iff \mathfrak{p} is maximal.
- 2 Show that, if A is a domain, then $\{(0)\}$ is a dense point.

Solution.

- 1 By definition, and by the previous exercise:

$$\overline{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in F \text{ closed}} F = \bigcap_{\substack{I \triangleleft R \\ I \subseteq \mathfrak{p}}} V(I) = V\left(\sum_{I \subseteq \mathfrak{p}} I\right) = V(\mathfrak{p}).$$

In particular, $\{\mathfrak{p}\}$ is closed iff $\{\mathfrak{p}\} = V(\mathfrak{p})$ which occurs iff \mathfrak{p} is maximal (o/w, take $\mathfrak{m} \supsetneq \mathfrak{p}$ maximal).



Exercise

Let A be a ring.

- 1 Show that $\overline{\{\mathfrak{p}\}} = V(\mathfrak{p})$, for all $\mathfrak{p} \in \operatorname{Spec}(A)$ and, in particular, that $\{\mathfrak{p}\}$ is closed iff \mathfrak{p} is maximal.
- 2 Show that, if A is a domain, then $\{(0)\}$ is a dense point.

Solution.

- 1 By definition, and by the previous exercise:

$$\overline{\mathfrak{p}} = \bigcap_{\mathfrak{p} \in F \text{ closed}} F = \bigcap_{\substack{I \triangleleft R \\ I \subseteq \mathfrak{p}}} V(I) = V\left(\sum_{I \subseteq \mathfrak{p}} I\right) = V(\mathfrak{p}).$$

In particular, $\{\mathfrak{p}\}$ is closed iff $\{\mathfrak{p}\} = V(\mathfrak{p})$ which occurs iff \mathfrak{p} is maximal (o/w, take $\mathfrak{m} \supsetneq \mathfrak{p}$ maximal).

- 2 Note: $(0) \in \operatorname{Spec}(R)$ iff R is a domain, in which case $V(0) = R$.



Exercise

Show that $\text{Spec}(A)$ is irreducible iff A is a domain.

Exercise

Show that $\text{Spec}(A)$ is irreducible iff A is a domain.

Solution.

Assume A is domain, and $0 \neq I_1, I_2 \triangleleft A$ proper ideals such that $\text{Spec}(A) = V(I_1) \cup V(I_2)$. Let $f_i \in I_i$ ($i = 1, 2$). Then, since

$$V(I_1) \cup V(I_2) \subseteq V((f_1)) \cup V((f_2)) = V((f_1 f_2)),$$

we have that $f_1 f_2$ is necessarily nilpotent, and since A is a domain, either $f_1 = 0$ or $f_2 = 0$. By fixing $0 \neq f_1 \in I_1$ (wlog, assuming such exists) and letting $f_2 \in I_2$ vary we deduce that $I_2 = (0)$ and hence $V(I_2) = \text{Spec}(A)$.

Exercise

Show that $\text{Spec}(A)$ is irreducible iff A is a domain.

Solution.

Assume A is domain, and $0 \neq I_1, I_2 \triangleleft A$ proper ideals such that $\text{Spec}(A) = V(I_1) \cup V(I_2)$. Let $f_i \in I_i$ ($i = 1, 2$). Then, since

$$V(I_1) \cup V(I_2) \subseteq V((f_1)) \cup V((f_2)) = V((f_1 f_2)),$$

we have that $f_1 f_2$ is necessarily nilpotent, and since A is a domain, either $f_1 = 0$ or $f_2 = 0$. By fixing $0 \neq f_1 \in I_1$ (wlog, assuming such exists) and letting $f_2 \in I_2$ vary we deduce that $I_2 = (0)$ and hence $V(I_2) = \text{Spec}(A)$.

For the converse implication, assume $\text{Spec}(A)$ is irreducible and let $f_1, f_2 \in A$ non-nilpotents. By assumption

$V((f_1)) \cup V((f_2)) = V((f_1 f_2)) \subsetneq \text{Spec}(A)$, and hence there exists $\mathfrak{p} \in \text{Spec}(A)$ such that $f_1 f_2 \notin \mathfrak{p}$. In particular, $f_1 f_2 \neq 0$. □

The last two exercises imply the following:

The last two exercises imply the following:

Corollary

If $\text{Spec}(A)$ is irreducible then there exists $\xi \in \text{Spec}(A)$ such that $\overline{\xi} = \text{Spec}(A)$.

The last two exercises imply the following:

Corollary

If $\text{Spec}(A)$ is irreducible then there exists $\xi \in \text{Spec}(A)$ such that $\overline{\xi} = \text{Spec}(A)$. Such an element is called a generic point of $\text{Spec}(A)$.

Ringed spaces and locally ringed spaces

Definition

Ringed spaces and locally ringed spaces

Definition

- 1 A *ringed space* is a pair (X, \mathcal{O}_X) consisting of a topological space and a sheaf of rings \mathcal{O}_X on X .

Ringed spaces and locally ringed spaces

Definition

- 1 A *ringed space* is a pair (X, \mathcal{O}_X) consisting of a topological space and a sheaf of rings \mathcal{O}_X on X .
- 2 A *morphism of ringed spaces* is a pair $(f, f^\#)$, in which f is a continuous map $X \rightarrow Y$ and $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is a morphism of sheaves of rings on Y .

Ringed spaces and locally ringed spaces

Definition

- 1 A *ringed space* is a pair (X, \mathcal{O}_X) consisting of a topological space and a sheaf of rings \mathcal{O}_X on X .
- 2 A *morphism of ringed spaces* is a pair $(f, f^\#)$, in which f is a continuous map $X \rightarrow Y$ and $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is a morphism of sheaves of rings on Y .
- 3 (X, \mathcal{O}_X) is a *locally ringed space* if, in addition, the stalk $\mathcal{O}_{X,x}$ is a local ring for all $x \in X$.

Ringed spaces and locally ringed spaces

Definition

- 1 A *ringed space* is a pair (X, \mathcal{O}_X) consisting of a topological space and a sheaf of rings \mathcal{O}_X on X .
- 2 A *morphism of ringed spaces* is a pair $(f, f^\#)$, in which f is a continuous map $X \rightarrow Y$ and $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is a morphism of sheaves of rings on Y .
- 3 (X, \mathcal{O}_X) is a *locally ringed space* if, in addition, the stalk $\mathcal{O}_{X,x}$ is a local ring for all $x \in X$.
- 4 A morphism of locally ringed spaces $(f, f^\#)$ is a morphism of ringed spaces with the **added requirement** that $f^\#_x : \mathcal{O}_{Y,f(x)} \rightarrow f_{*}\mathcal{O}_{X,x}$ is a local homomorphism (i.e. preimage of the maximal ideal is maximal) for any $x \in X$.

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

- for any $\mathfrak{p} \in \mathrm{Spec}(A)$, the stalk $\mathcal{O}_{\mathfrak{p}}$ of \mathcal{O} at \mathfrak{p} is isomorphic to the local ring $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$, and

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

- for any $\mathfrak{p} \in \mathrm{Spec}(A)$, the stalk $\mathcal{O}_{\mathfrak{p}}$ of \mathcal{O} at \mathfrak{p} is isomorphic to the local ring $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$, and
- for any $f \in A$, let $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$; this is an open set of $\mathrm{Spec}(A)$. Then $\mathcal{O}(D(f)) \simeq f^{-1}A$.

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

- for any $\mathfrak{p} \in \mathrm{Spec}(A)$, the stalk $\mathcal{O}_{\mathfrak{p}}$ of \mathcal{O} at \mathfrak{p} is isomorphic to the local ring $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$, and
- for any $f \in A$, let $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$; this is an open set of $\mathrm{Spec}(A)$. Then $\mathcal{O}(D(f)) \simeq f^{-1}A$.

Thus $(\mathrm{Spec}(A), \mathcal{O})$ is a locally ring space, with $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$.

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

- for any $\mathfrak{p} \in \mathrm{Spec}(A)$, the stalk $\mathcal{O}_{\mathfrak{p}}$ of \mathcal{O} at \mathfrak{p} is isomorphic to the local ring $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$, and
- for any $f \in A$, let $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$; this is an open set of $\mathrm{Spec}(A)$. Then $\mathcal{O}(D(f)) \simeq f^{-1}A$.

Thus $(\mathrm{Spec}(A), \mathcal{O})$ is a locally ring space, with $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$.

Explicit construction

The sheaf \mathcal{O} is defined by sheafifying the presheaf on $\mathrm{Spec}(A)$ whose stalks are given by $A_{\mathfrak{p}} = \lim_{\mathfrak{p} \in D(f), f \in A} A_f$.

The sheaf of rings on $\mathrm{Spec}(A)$

Let A be a ring. The spectrum of A has an associated sheaf of rings \mathcal{O} , which has the following properties:

- for any $\mathfrak{p} \in \mathrm{Spec}(A)$, the stalk $\mathcal{O}_{\mathfrak{p}}$ of \mathcal{O} at \mathfrak{p} is isomorphic to the local ring $A_{\mathfrak{p}} = (A - \mathfrak{p})^{-1}A$, and
- for any $f \in A$, let $D(f) = \{\mathfrak{p} \in \mathrm{Spec}(A) : f \notin \mathfrak{p}\}$; this is an open set of $\mathrm{Spec}(A)$. Then $\mathcal{O}(D(f)) \simeq f^{-1}A$.

Thus $(\mathrm{Spec}(A), \mathcal{O})$ is a locally ring space, with $\mathcal{O}(\mathrm{Spec}(A)) \simeq A$.

Explicit construction

The sheaf \mathcal{O} is defined by sheafifying the presheaf on $\mathrm{Spec}(A)$ whose stalks are given by $A_{\mathfrak{p}} = \lim_{\mathfrak{p} \in D(f), f \in A} A_f$. Explicitly, $\mathcal{O}(U)$ consists of functions $s : U \rightarrow \bigsqcup_{\mathfrak{p} \in U} A_{\mathfrak{p}}$ for which there exists an open cover $U = \bigcup_{\alpha} V_{\alpha}$ such that, given α , $\exists a_{\alpha}, b_{\alpha} \in A$ with $V_{\alpha} \subseteq D(b_{\alpha})$, such that $s(\mathfrak{q}) =$ “the image of a_{α}/b_{α} in $A_{\mathfrak{q}}$ ”, for all $\mathfrak{q} \in V_{\alpha}$.

Proposition (2.3 in Hartshorne)

- ① *Let $\varphi : A \rightarrow B$ be a homomorphism of rings. Then φ induces a natural morphism of locally ringed spaces*

$$(f, f^\#) : (\mathrm{Spec}(B), \mathcal{O}_{\mathrm{Spec}(B)}) \rightarrow (\mathrm{Spec}(A), \mathcal{O}_{\mathrm{Spec}(A)}).$$

- ② *Conversely, any morphism of locally ring spaces $(f, f^\#)$ as above is induced from a ring homomorphism $\varphi : A \rightarrow B$.*

Example

Let R be a dvr and $K = \text{Frac}(R)$, its field of fractions.

Example

Let R be a dvr and $K = \text{Frac}(R)$, its field of fractions. The space $T = \text{Spec}(R)$ has two points. One of them, denoted t_0 , is closed and corresponds to the maximal ideal \mathfrak{m} of R . The second, denoted t_0 is open and dense, and corresponds to the zero ideal.

Example

Let R be a dvr and $K = \text{Frac}(R)$, its field of fractions. The space $T = \text{Spec}(R)$ has two points. One of them, denoted t_0 , is closed and corresponds to the maximal ideal \mathfrak{m} of R . The second, denoted t_1 is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$

Example

Let R be a dvr and $K = \text{Frac}(R)$, its field of fractions. The space $T = \text{Spec}(R)$ has two points. One of them, denoted t_0 , is closed and corresponds to the maximal ideal \mathfrak{m} of R . The second, denoted t_1 is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$

The map $f : S = \text{Spec}(K) \rightarrow T$, sending the only point in $\text{Spec}(K)$ to t_1 , is part of a morphism $(f, f^\#)$ of **locally** ringed spaces, since the corresponding stalks are both K .

Example

Let R be a dvr and $K = \text{Frac}(R)$, its field of fractions. The space $T = \text{Spec}(R)$ has two points. One of them, denoted t_0 , is closed and corresponds to the maximal ideal \mathfrak{m} of R . The second, denoted t_1 is open and dense, and corresponds to the zero ideal.

$$\mathcal{O}_{T,t_0} \simeq R_{\mathfrak{m}} = R \quad \text{and} \quad \mathcal{O}_{T,t_1} \simeq R_{(0)} = K.$$

The map $f : S = \text{Spec}(K) \rightarrow T$, sending the only point in $\text{Spec}(K)$ to t_1 , is part of a morphism $(f, f^\#)$ of **locally** ringed spaces, since the corresponding stalks are both K . Therefore, it is induced from a ring homomorphism $\varphi : R \rightarrow K$.

Example

Example - contd On the other hand, if $\text{Im}(f) = \{t_0\}$, and $(f, f^\#)$ is a morphism of ringed spaces, then $f^\#$ induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

Example

Example - contd On the other hand, if $\text{Im}(f) = \{t_0\}$, and $(f, f^\#)$ is a morphism of ringed spaces, then $f^\#$ induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

Assuming $R/\mathfrak{m} \not\cong K$, this is **not** a local homomorphism.

Example

Example - contd On the other hand, if $\text{Im}(f) = \{t_0\}$, and $(f, f^\#)$ is a morphism of ringed spaces, then $f^\#$ induces a homomorphism on stalks

$$f^\# : \mathcal{O}_{\text{Spec}(R), t_0} = R \rightarrow K.$$

Assuming $R/\mathfrak{m} \not\cong K$, this is **not** a local homomorphism. Therefore, in this case, $(f, f^\#)$ is not induced from any morphism $R \rightarrow K$.

Proof of proposition

Proof.

Proof of proposition

Proof.

- 1 Let $\varphi : A \rightarrow B$ be a ring homomorphism. Define $f : \operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$ by $f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \in \operatorname{Spec}(A)$.

Proof of proposition

Proof.

- ① Let $\varphi : A \rightarrow B$ be a ring homomorphism. Define $f : \operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$ by $f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \in \operatorname{Spec}(A)$.

f is continuous. Given $I \triangleleft A$, we have

$$\mathfrak{p} \in f^{-1}(V(I)) \iff f(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p}) \supseteq I \iff \mathfrak{p} \supseteq \varphi(I).$$

Thus $f^{-1}(V(I)) = V(\varphi(I))$.



Proof.

Construction of f^\sharp . For any $\mathfrak{p} \in \operatorname{Spec} B$, we have a localized map $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$, defined by $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$.

Proof.

Construction of $f^\#$. For any $\mathfrak{p} \in \operatorname{Spec} B$, we have a localized map $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$, defined by $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$. Given $V \subseteq \operatorname{Spec}(A)$ open, and $s \in \mathcal{O}_{\operatorname{Spec}(A)}(V)$ define

$$f^\#(s)(\mathfrak{p}) = \varphi_{\mathfrak{p}} \circ s \circ f.$$

Proof.

Construction of $f^\#$. For any $\mathfrak{p} \in \operatorname{Spec} B$, we have a localized map $\varphi_{\mathfrak{p}} : A_{\varphi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$, defined by $\varphi_{\mathfrak{p}}(a/b) = \varphi(a)/\varphi(b)$. Given $V \subseteq \operatorname{Spec}(A)$ open, and $s \in \mathcal{O}_{\operatorname{Spec}(A)}(V)$ define

$$f^\#(s)(\mathfrak{p}) = \varphi_{\mathfrak{p}} \circ s \circ f.$$

Then $f^\#$ is a ring homomorphism. Furthermore, if $U \subseteq V$ is such that there exist $a, b \in A$ with $U \subseteq D(b)$ such that $s(\mathfrak{q}) = a/b \in A_{\mathfrak{q}}$ for all $\mathfrak{q} \in U$, then $f^{-1}(U) \subseteq D(\varphi(b))$ and $f^\#(s)(\mathfrak{q}') = \varphi(a)/\varphi(b) \in B_{\mathfrak{q}'}$ for all $\mathfrak{q}' \in f^{-1}(U)$. It follows that $f^\#(s) \in \mathcal{O}_{\operatorname{Spec}(B)}(f^{-1}(V)) = f_*\mathcal{O}_{\operatorname{Spec}(A)}(V)$.

Proof.

- ② Given a morphism of locally ringed spaces $(f, f^\#)$, in particular, we get a map $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$, where we identify a ring R with $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$.

Proof.

- ② Given a morphism of locally ringed spaces $(f, f^\#)$, in particular, we get a map $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$, where we identify a ring R with $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$. φ must agree with $f^\#$ on stalks, i.e. we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ I_A \downarrow & & \downarrow I_B \\ A_{f(p)} & \xrightarrow[p]{f^\#} & B_p \end{array} .$$

Proof.

- ② Given a morphism of locally ringed spaces $(f, f^\#)$, in particular, we get a map $\varphi := f^\#_{\text{Spec}(A)} : A \rightarrow B$, where we identify a ring R with $\mathcal{O}_{\text{Spec}(R)}(\text{Spec}(R))$. φ must agree with $f^\#$ on stalks, i.e. we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ I_A \downarrow & & \downarrow I_B \\ A_{f(p)} & \xrightarrow[p]{f^\#} & B_p \end{array} .$$

In particular, since the bottom row consists of local rings and a local homomorphism, it must be that

$$\varphi^{-1}(\mathfrak{p}) = \varphi^{-1} \circ I_B^{-1}(\mathfrak{p}B_p) = I_A^{-1} \circ f^\#(\mathfrak{p}B_p) = f(\mathfrak{p}).$$

Thus $f = \varphi^{-1}$.

Example

Let $R = \overline{\mathbb{F}_q}[[t]]$, and $\varphi : R \rightarrow R$ defined by $\varphi(\sum a_i t^i) = \sum a_i^q t^i$.

Example

Let $R = \overline{\mathbb{F}_q}[[t]]$, and $\varphi : R \rightarrow R$ defined by $\varphi(\sum a_i t^i) = \sum a_i^q t^i$.
 $T = \text{Spec}(R)$ has two points with $\mathfrak{p} = (t)$, a closed point, and (0) , which is open and dense. Both points are invariant under φ^{-1} .

Example

Let $R = \overline{\mathbb{F}_q}[[t]]$, and $\varphi : R \rightarrow R$ defined by $\varphi(\sum a_i t^i) = \sum a_i^q t^i$.
 $T = \operatorname{Spec}(R)$ has two points with $\mathfrak{p} = (t)$, a closed point, and (0) , which is open and dense. Both points are invariant under φ^{-1} .

- The associated map $f : \operatorname{Spec}(R) \rightarrow \operatorname{Spec}(R)$ is the identity map.

Example

Let $R = \overline{\mathbb{F}_q}[[t]]$, and $\varphi : R \rightarrow R$ defined by $\varphi(\sum a_i t^i) = \sum a_i^q t^i$.
 $T = \operatorname{Spec}(R)$ has two points with $\mathfrak{p} = (t)$, a closed point, and (0) , which is open and dense. Both points are invariant under φ^{-1} .

- The associated map $f : \operatorname{Spec}(R) \rightarrow \operatorname{Spec}(R)$ is the identity map.
- Given $V \subseteq \operatorname{Spec}(R)$ and $s \in \mathcal{O}_{\operatorname{Spec}(R)}(V)$, we have

$$f^\#(s)(\cdot) = (s(\cdot))^q.$$

Corollary (of the proposition)

Let A, B be rings. Then A and B are isomorphic if and only if $\operatorname{Spec}(A)$ and $\operatorname{Spec}(B)$ are isomorphic as locally ringed spaces.

Definition

An *affine scheme* is a locally ringed space (X, \mathcal{O}_X) which is isomorphic to $\mathrm{Spec}(A)$ for some ring A .

Definition

An *affine scheme* is a locally ringed space (X, \mathcal{O}_X) which is isomorphic to $\mathrm{Spec}(A)$ for some ring A .

A *scheme* is a locally ringed space (X, \mathcal{O}_X) for which X is covered by open sets $X = \bigcup U_\alpha$, such that $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$ is an affine scheme.

Definition

An *affine scheme* is a locally ringed space (X, \mathcal{O}_X) which is isomorphic to $\mathrm{Spec}(A)$ for some ring A .

A *scheme* is a locally ringed space (X, \mathcal{O}_X) for which X is covered by open sets $X = \bigcup U_\alpha$, such that $(U_\alpha, \mathcal{O}_X|_{U_\alpha})$ is an affine scheme.

Here $\mathcal{O}_X|_{U_\alpha}$ denotes the restricted sheaf $\mathcal{O}_X|_{U_\alpha}(V) = \mathcal{O}_X(V)$ for $V \subseteq U_\alpha$ open.