

# On Regular Representations of Groups

Shai Shechter

Ben Gurion University of the Negev, Beer Sheva

June 2016

# Introduction

Jaikin's Example-  $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

Norm-1 Subgroups of Local Division Algebras

Regular Representations

# Motivation

- ▶  $\Gamma$  is a topological group.

# Motivation

- ▶  $\Gamma$  is a topological group.
- ▶  $\text{Irr}(\Gamma)$  is the set of continuous irreducible complex **finite-dimensional** representations of  $\Gamma$ , upto equivalence.

# Motivation

- ▶  $\Gamma$  is a topological group.
- ▶  $\text{Irr}(\Gamma)$  is the set of continuous irreducible complex **finite-dimensional** representations of  $\Gamma$ , upto equivalence.
- ▶ **Mission-** Understand  $\text{Irr}(\Gamma)$ .

# Understand $\text{Irr}(\Gamma)$ ?

- ▶ **Possible Answer -**

# Understand $\text{Irr}(\Gamma)$ ?

- ▶ **Possible Answer -**

- ▶ Parametrize all irreducible representations of  $\Gamma$  according to some known parametrizing space

$$X \leftrightarrow \text{Irr}(\Gamma).$$

- ▶ For each class  $[\rho] \in \text{Irr}(\Gamma)$ , present a complex finite dimensional vector space  $V_\rho$ , and the homomorphism  $\rho : \Gamma \rightarrow \text{Aut}(V_\rho)$ .

# Understand $\text{Irr}(\Gamma)$ ?

## ► Possible Answer -

- Parametrize all irreducible representations of  $\Gamma$  according to some known parametrizing space

$$X \leftrightarrow \text{Irr}(\Gamma).$$

- For each class  $[\rho] \in \text{Irr}(\Gamma)$ , present a complex finite dimensional vector space  $V_\rho$ , and the homomorphism  $\rho : \Gamma \rightarrow \text{Aut}(V_\rho)$ .

## ► Another Option- Describe all characters of $\Gamma$ explicitly

$$\chi_\rho(\gamma) = \text{Tr}(\rho(\gamma)) \quad (\gamma \in \Gamma).$$



# Understand $\text{Irr}(\Gamma)$ ?

- **Asymptotics**- For any  $n \in \mathbb{N}$ , define

$$r_n(\Gamma) := \# \{[\rho] \in \text{Irr}(\Gamma) \mid \dim(\rho) = n\} \in \mathbb{N} \cup \{0, \infty\}.$$

# Understand $\text{Irr}(\Gamma)$ ?

- **Asymptotics**- For any  $n \in \mathbb{N}$ , define

$$r_n(\Gamma) := \# \{[\rho] \in \text{Irr}(\Gamma) \mid \dim(\rho) = n\} \in \mathbb{N} \cup \{0, \infty\}.$$

Describe the asymptotic behaviour of the sequence  $\{r_n(\Gamma)\}$ .

# Understand $\text{Irr}(\Gamma)$ ?

- **Asymptotics**- For any  $n \in \mathbb{N}$ , define

$$r_n(\Gamma) := \# \{[\rho] \in \text{Irr}(\Gamma) \mid \dim(\rho) = n\} \in \mathbb{N} \cup \{0, \infty\}.$$

Describe the asymptotic behaviour of the sequence  $\{r_n(\Gamma)\}$ .  
E.g.- Do there exist  $c_1, c_2 \in \mathbb{R}$  such that

$$r_n(\Gamma) \leq c_1 n^{c_2}, \forall n \in \mathbb{N}?$$

# Understand $\text{Irr}(\Gamma)$ ?

- **Asymptotics**- For any  $n \in \mathbb{N}$ , define

$$r_n(\Gamma) := \# \{[\rho] \in \text{Irr}(\Gamma) \mid \dim(\rho) = n\} \in \mathbb{N} \cup \{0, \infty\}.$$

Describe the asymptotic behaviour of the sequence  $\{r_n(\Gamma)\}$ .  
E.g.- Do there exist  $c_1, c_2 \in \mathbb{R}$  such that

$$r_n(\Gamma) \leq c_1 n^{c_2}, \forall n \in \mathbb{N}?$$

The group  $\Gamma$  is said to be **rigid** if  $r_n(\Gamma) < \infty$  for all  $n$ .

# Arithmetic Groups

Let  $\Gamma$  be an arithmetic group, e.g.  $\Gamma = \mathrm{SL}_3(\mathbb{Z})$ . The group  $\Gamma$  is said to have **the congruence subgroup property** (CSP) if any finite-index normal subgroup of  $\Gamma$  contains a principal congruence subgroup.

For example, the principal congruence subgroups of  $\mathrm{SL}_3(\mathbb{Z})$  are all subgroups of the form  $\mathrm{Ker}(\mathrm{SL}_3(\mathbb{Z}) \rightarrow \mathrm{SL}_3(\mathbb{Z}/n\mathbb{Z}))$ , for some  $n \in \mathbb{N}$ .

# Arithmetic Groups

Let  $\Gamma$  be an arithmetic group, e.g.  $\Gamma = \mathrm{SL}_3(\mathbb{Z})$ . The group  $\Gamma$  is said to have **the congruence subgroup property** (CSP) if any finite-index normal subgroup of  $\Gamma$  contains a principal congruence subgroup.

For example, the principal congruence subgroups of  $\mathrm{SL}_3(\mathbb{Z})$  are all subgroups of the form  $\mathrm{Ker}(\mathrm{SL}_3(\mathbb{Z}) \rightarrow \mathrm{SL}_3(\mathbb{Z}/n\mathbb{Z}))$ , for some  $n \in \mathbb{N}$ .

## Theorem (Bass-Lazard-Serre, '64)

*The group  $\mathrm{SL}_3(\mathbb{Z})$  has the CSP.*

# Arithmetic Groups

Let  $\Gamma$  be an arithmetic group, e.g.  $\Gamma = \mathrm{SL}_3(\mathbb{Z})$ . The group  $\Gamma$  is said to have **the congruence subgroup property** (CSP) if any finite-index normal subgroup of  $\Gamma$  contains a principal congruence subgroup.

For example, the principal congruence subgroups of  $\mathrm{SL}_3(\mathbb{Z})$  are all subgroups of the form  $\mathrm{Ker}(\mathrm{SL}_3(\mathbb{Z}) \rightarrow \mathrm{SL}_3(\mathbb{Z}/n\mathbb{Z}))$ , for some  $n \in \mathbb{N}$ .

## Theorem (Bass-Lazard-Serre, '64)

*The group  $\mathrm{SL}_3(\mathbb{Z})$  has the CSP.*

*The same is true for  $\mathrm{SL}_k(\mathbb{Z})$  for all  $k \geq 3$ .*

# Arithmetic Groups

Let  $\Gamma$  be an arithmetic group, e.g.  $\Gamma = \mathrm{SL}_3(\mathbb{Z})$ . The group  $\Gamma$  is said to have **the congruence subgroup property** (CSP) if any finite-index normal subgroup of  $\Gamma$  contains a principal congruence subgroup.

For example, the principal congruence subgroups of  $\mathrm{SL}_3(\mathbb{Z})$  are all subgroups of the form  $\mathrm{Ker}(\mathrm{SL}_3(\mathbb{Z}) \rightarrow \mathrm{SL}_3(\mathbb{Z}/n\mathbb{Z}))$ , for some  $n \in \mathbb{N}$ .

## Theorem (Bass-Lazard-Serre, '64)

*The group  $\mathrm{SL}_3(\mathbb{Z})$  has the CSP.*

*The same is true for  $\mathrm{SL}_k(\mathbb{Z})$  for all  $k \geq 3$ .*

*The group  $\mathrm{SL}_2(\mathbb{Z})$ , however, does not have CSP.*



# Arithmetic Groups

## Theorem (Lubotzky-Martin, '04)

*An arithmetic group over a number field has polynomial representation growth iff it has CSP.*

# Arithmetic Groups

## Theorem (Lubotzky-Martin, '04)

*An arithmetic group over a number field has polynomial representation growth iff it has CSP.*

The **representation zeta function** of  $\Gamma$  is

$$\zeta_{\Gamma}(s) = \sum_{n=1}^{\infty} r_n(\Gamma) n^{-s} = \sum_{[\rho] \in \text{Irr}(\Gamma)} \dim(\rho)^{-s}, \quad (s \in \mathbb{C}). \quad (*)$$

# Arithmetic Groups

The representation zeta function of  $\Gamma$  admits an Euler product-

$$\zeta_{\mathrm{SL}_3(\mathbb{Z})}(s) = \zeta_{\mathrm{SL}_3(\mathbb{C})}(s) \cdot \prod_{p \text{ is prime}} \zeta_{\mathrm{SL}_3(\mathbb{Z}_p)}(s).$$

# Arithmetic Groups

The representation zeta function of  $\Gamma$  admits an Euler product-

$$\zeta_{\mathrm{SL}_3(\mathbb{Z})}(s) = \zeta_{\mathrm{SL}_3(\mathbb{C})}(s) \cdot \prod_{p \text{ is prime}} \zeta_{\mathrm{SL}_3(\mathbb{Z}_p)}(s).$$

Here-

- ▶  $\zeta_{\mathrm{SL}_3(\mathbb{Z})}$  enumerates arbitrary representations of  $\Gamma$ .
- ▶ The archimedean factor  $\zeta_{\mathrm{SL}_3(\mathbb{C})}$  enumerates rational representations of  $\mathrm{SL}_3(\mathbb{C})$ .

# Arithmetic Groups

The representation zeta function of  $\Gamma$  admits an Euler product-

$$\zeta_{\mathrm{SL}_3(\mathbb{Z})}(s) = \zeta_{\mathrm{SL}_3(\mathbb{C})}(s) \cdot \prod_{p \text{ is prime}} \zeta_{\mathrm{SL}_3(\mathbb{Z}_p)}(s).$$

Here-

- ▶  $\zeta_{\mathrm{SL}_3(\mathbb{Z})}$  enumerates arbitrary representations of  $\Gamma$ .
- ▶ The archimedean factor  $\zeta_{\mathrm{SL}_3(\mathbb{C})}$  enumerates rational representations of  $\mathrm{SL}_3(\mathbb{C})$ .
- ▶ The non-archimedean factors  $\zeta_{\mathrm{SL}_3(\mathbb{Z}_p)}$  enumerate continuous representations.

# $p$ -adic Groups

## Theorem (Jaikin-Zapirain, '07)

*Fix an odd prime  $p \in \mathbb{Z}$ . Let  $G$  be a rigid compact  $p$ -adic analytic group. There exist rational functions  $f_1(t), \dots, f_k(t) \in \mathbb{Q}(t)$  and numbers  $n_1, \dots, n_k \in \mathbb{N}$  such that*

$$\zeta_G(s) = \sum_{j=1}^k n_j^{-s} \cdot f_j(p^{-s}).$$

# $p$ -adic Groups

## Theorem (Jaikin-Zapirain, '07)

*Fix an odd prime  $p \in \mathbb{Z}$ . Let  $G$  be a rigid compact  $p$ -adic analytic group. There exist rational functions  $f_1(t), \dots, f_k(t) \in \mathbb{Q}(t)$  and numbers  $n_1, \dots, n_k \in \mathbb{N}$  such that*

$$\zeta_G(s) = \sum_{j=1}^k n_j^{-s} \cdot f_j(p^{-s}).$$

**Example (AKOV)** Assume  $p > 2$  and let  $G \subseteq \mathrm{SL}_2(\mathbb{Z}_p)$  be given by  $\mathrm{SL}_2(\mathbb{Z}_p) \cap (1 + p\mathrm{M}_2(\mathbb{Z}_p))$ . Then

$$\zeta_G(s) = p^3 \frac{1 - p^{-2-s}}{1 - p^{1-s}}.$$

# Ingredients of the Rationality Theorem



# Ingredients of the Rationality Theorem

1. Kirillov orbit method for  $p$ -adic groups (Howe).

$$\mathrm{Irr}(G) \xleftrightarrow{1-1} \widehat{\mathfrak{g}}/G.$$

# Ingredients of the Rationality Theorem

1. Kirillov orbit method for  $p$ -adic groups (Howe).

$$\mathrm{Irr}(G) \xleftrightarrow{1-1} \widehat{\mathfrak{g}}/G.$$

2. Interpretation of the representation zeta function as an Igusa-type  $p$ -adic integral.

$$\zeta_G(s) \approx \int_X |g(\mathbf{x})|_p^{-s} d\mathbf{x}.$$

# Ingredients of the Rationality Theorem

1. Kirillov orbit method for  $p$ -adic groups (Howe).

$$\mathrm{Irr}(G) \xleftrightarrow{1-1} \widehat{\mathfrak{g}}/G.$$

2. Interpretation of the representation zeta function as an Igusa-type  $p$ -adic integral.

$$\zeta_G(s) \approx \int_X |g(\mathbf{x})|_p^{-s} d\mathbf{x}.$$

3. Model theoretic proof of the rationality of such integrals (Denef).

# Jaikin's Example

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).
- ▶  $\mathfrak{o} \supseteq \mathfrak{p}$  are the valuation ring and maximal ideal,  $\pi$  is a uniformizer. Let  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).
- ▶  $\mathfrak{o} \supseteq \mathfrak{p}$  are the valuation ring and maximal ideal,  $\pi$  is a uniformizer. Let  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$ . We assume  $p = \mathrm{char}(\mathbb{F}_q) > 2$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).
- ▶  $\mathfrak{o} \supseteq \mathfrak{p}$  are the valuation ring and maximal ideal,  $\pi$  is a uniformizer. Let  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$ . We assume  $p = \mathrm{char}(\mathbb{F}_q) > 2$ .
- ▶ Fix  $G = \mathrm{SL}_2(\mathfrak{o})$ .



# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).
- ▶  $\mathfrak{o} \supseteq \mathfrak{p}$  are the valuation ring and maximal ideal,  $\pi$  is a uniformizer. Let  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$ . We assume  $p = \mathrm{char}(\mathbb{F}_q) > 2$ .
- ▶ Fix  $G = \mathrm{SL}_2(\mathfrak{o})$ . The subgroups  $G^m = \mathrm{SL}_2(\mathfrak{o}) \cap (1 + \pi^m \mathrm{M}_2(\mathfrak{o}))$  are called the **principal congruence subgroups** of  $G$ . The sequence  $\{G^m\}$  is a neighbourhood basis at 1.

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

- ▶  $K$  is a local non-archimedean local field (e.g.  $K = \mathbb{Z}_p, \mathbb{F}_q[t]$ ).
- ▶  $\mathfrak{o} \supseteq \mathfrak{p}$  are the valuation ring and maximal ideal,  $\pi$  is a uniformizer. Let  $\mathbb{F}_q = \mathfrak{o}/\mathfrak{p}$ . We assume  $p = \mathrm{char}(\mathbb{F}_q) > 2$ .
- ▶ Fix  $G = \mathrm{SL}_2(\mathfrak{o})$ . The subgroups  $G^m = \mathrm{SL}_2(\mathfrak{o}) \cap (1 + \pi^m \mathrm{M}_2(\mathfrak{o}))$  are called the **principal congruence subgroups** of  $G$ . The sequence  $\{G^m\}$  is a neighbourhood basis at 1.

## Definition

The **level** of a representation  $\rho$  of  $G$  is the minimal  $k \in \mathbb{N}_0$  such that  $G^{k+1} \subseteq \mathrm{Ker}(\rho)$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

## Definition

The **level** of a representation  $\rho$  of  $G$  is the minimal  $k \in \mathbb{N}_0$  such that  $G^{k+1} \subseteq \mathrm{Ker}(\rho)$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

For example, since the map  $G \rightarrow \mathrm{SL}_2(\mathbb{F}_q)$  is surjective, there is a natural identification

$$\{(\text{irr.}) \text{ reps of level } 0\} \longleftrightarrow \{(\text{irr.}) \text{ reps of } \mathrm{SL}_2(\mathbb{F}_q)\}.$$

## Definition

The **level** of a representation  $\rho$  of  $G$  is the minimal  $k \in \mathbb{N}_0$  such that  $G^{k+1} \subseteq \mathrm{Ker}(\rho)$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$

For example, since the map  $G \rightarrow \mathrm{SL}_2(\mathbb{F}_q)$  is surjective, there is a natural identification

$$\{(\text{irr.}) \text{ reps of level } 0\} \longleftrightarrow \{(\text{irr.}) \text{ reps of } \mathrm{SL}_2(\mathbb{F}_q)\}.$$

Put  $\mathrm{Irr}^k(G) := \{\rho \in \mathrm{Irr}(G) \mid \rho \text{ of level } k\}$ . Then

$$\bigcup_{m \leq k} \mathrm{Irr}^m(G) = \mathrm{Irr}(\mathrm{SL}_2(\mathfrak{o}/\mathfrak{p}^{k+1})).$$

## Definition

The **level** of a representation  $\rho$  of  $G$  is the minimal  $k \in \mathbb{N}_0$  such that  $G^{k+1} \subseteq \mathrm{Ker}(\rho)$ .

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$ - Strategy

- ▶ In what follows, we will describe an explicit map between the set  $\mathrm{Irr}^k(G)$  that the space of orbits in a finite  $G$ -space.

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$ - Strategy

- ▶ In what follows, we will describe an explicit map between the set  $\mathrm{Irr}^k(G)$  that the space of orbits in a finite  $G$ -space.
- ▶ Using this map, we can compute the cardinality of  $\mathrm{Irr}^k(G)$ , as well as the dimensions of its elements, thereby allowing us to compute

$$\zeta_{\mathrm{SL}_2(\mathfrak{o}/\mathfrak{p}^{k+1})}(s) = \sum_{\substack{\rho \in \mathrm{Irr}^m(G) \\ m \leq k}} \dim(\rho)^{-s}.$$

# Computing $\zeta_{\mathrm{SL}_2(\mathfrak{o})}$ - Strategy (contd.)

Applying the process above for all  $k \in \mathbb{N}$  we obtain-

## Theorem (Jaikin-Zapirain, '07)

$$\zeta_{\mathrm{SL}_2(\mathfrak{o})}(s) = \zeta_{\mathrm{SL}_2(\mathbb{F}_q)}(s) + \frac{4q \left(\frac{q^2-1}{2}\right)^{-s} + \frac{q^2-1}{2}(q^2-q)^{-s} + \left(\frac{q-1}{2}\right)^2 (q^2+q)^{-s}}{1-q^{1-s}}.$$



# General Notation

- ▶  $G^r$ - superscript indices stand for congruence subgroups.
- ▶  $G_m$ - subscript indices stand for congruence quotients- i.e. quotients of  $G$  by a subgroup  $G^m$ .
- ▶  $G_m^r$ - both super- and subscript indices signify congruence subquotients,  $G^r/G^m$ .

# General Notation

- ▶  $G^r$ - superscript indices stand for congruence subgroups.
- ▶  $G_m$ - subscript indices stand for congruence quotients- i.e. quotients of  $G$  by a subgroup  $G^m$ .
- ▶  $G_m^r$ - both super- and subscript indices signify congruence subquotients,  $G^r/G^m$ .

Similar notation will be used for Lie-algebras.

# General Notation

- ▶  $G^r$ - superscript indices stand for congruence subgroups.
- ▶  $G_m$ - subscript indices stand for congruence quotients- i.e. quotients of  $G$  by a subgroup  $G^m$ .
- ▶  $G_m^r$ - both super- and subscript indices signify congruence subquotients,  $G^r/G^m$ .

Similar notation will be used for Lie-algebras. From here on we fix  $K \supseteq \mathfrak{o} \supseteq \mathfrak{p} = \pi\mathfrak{o}$ .

# Construction of the Map

- ▶ Let  $\rho \in \mathrm{Irr}^{k-1}(G)$  be fixed, with  $k > 1$ .

# Construction of the Map

- ▶ Let  $\rho \in \mathrm{Irr}^{k-1}(G)$  be fixed, with  $k > 1$ . Throughout, we will consider  $\rho$  as an element of  $\mathrm{Irr}(G_k)$ , which is non-trivial on the subgroup  $G_k^{k-1}$ .
- ▶ Put  $r = \lceil \frac{k}{2} \rceil$ , and consider the restriction of  $\rho$  to the group  $G_k^r$ .

# Construction of the Map

- ▶ Let  $\rho \in \mathrm{Irr}^{k-1}(G)$  be fixed, with  $k > 1$ . Throughout, we will consider  $\rho$  as an element of  $\mathrm{Irr}(G_k)$ , which is non-trivial on the subgroup  $G_k^{k-1}$ .
- ▶ Put  $r = \lceil \frac{k}{2} \rceil$ , and consider the restriction of  $\rho$  to the group  $G_k^r$ .

# Construction of the Map

- ▶ Let  $\rho \in \mathrm{Irr}^{k-1}(G)$  be fixed, with  $k > 1$ . Throughout, we will consider  $\rho$  as an element of  $\mathrm{Irr}(G_k)$ , which is non-trivial on the subgroup  $G_k^{k-1}$ .
- ▶ Put  $r = \lceil \frac{k}{2} \rceil$ , and consider the restriction of  $\rho$  to the group  $G_k^r$ . The following hold
  - ▶ The group  $G_k^r$  is abelian.
  - ▶ The map  $1 + \pi^r x \mapsto x$  is an isomorphism of abelian groups between  $G^r/G^k$  and the finite Lie-ring  $\mathfrak{g}_{k-r} := \mathfrak{sl}_2(\mathfrak{o}/\mathfrak{p}^{k-r})$  of traceless  $2 \times 2$  matrices over  $\mathfrak{o}/\mathfrak{p}^{k-r}$ .
  - ▶ There is a  $G$ -equivariant bijection of  $\mathfrak{g}_{k-r}$  with its Pontryagin dual  $\widehat{\mathfrak{g}_{k-r}}$ ,

$$y \mapsto \phi_y(x) := \chi(\mathrm{Tr}(xy)).$$

# Construction of the Map

Get a  $G$ -equivariant bijection

$$\mathfrak{g}_{k-r} \xrightarrow{1-1} \widehat{G_k^r} = \mathrm{Irr}(G_k^r)$$

$$y \mapsto \theta_y(1 + \pi^r x) = \phi_y(x).$$



# Construction of the Map

Get a  $G$ -equivariant bijection

$$\begin{aligned} \mathfrak{g}_{k-r} &\xrightarrow{1-1} \widehat{G_k^r} = \mathrm{Irr}(G_k^r) \\ y &\mapsto \theta_y(1 + \pi^r x) = \phi_y(x). \end{aligned}$$

By Clifford Theory, there exists  $y \in \mathfrak{g}_{k-r}$  which is unique up to  $G_k$ -conjugation and a fixed  $e \in \mathbb{N}$ , such that

$$\rho_{G_k^r} = \left( \bigoplus_{y' \in \mathrm{Ad}(G_k)y} \theta_{y'} \right)^{\oplus e}.$$

# Construction of the Map

Get a  $G$ -equivariant bijection

$$\begin{aligned} \mathfrak{g}_{k-r} &\xrightarrow{1-1} \widehat{G_k^r} = \mathrm{Irr}(G_k^r) \\ y &\mapsto \theta_y(1 + \pi^r x) = \phi_y(x). \end{aligned}$$

By Clifford Theory, there exists  $y \in \mathfrak{g}_{k-r}$  which is unique up to  $G_k$ -conjugation and a fixed  $e \in \mathbb{N}$ , such that

$$\rho_{G_k^r} = \left( \bigoplus_{y' \in \mathrm{Ad}(G_k)y} \theta_{y'} \right)^{\oplus e}.$$

We obtain a mapping  $\rho \mapsto \mathrm{Ad}(G_k)y$ .

Put  $S = \mathrm{Stab}^{G_k}(\theta_y) := \{g \in G_k \mid \theta_y(g^{-1}xg) = \theta_y(x), \quad \forall x\}.$

Put  $S = \mathrm{Stab}^{G_k}(\theta_y) := \{g \in G_k \mid \theta_y(g^{-1}xg) = \theta_y(x), \quad \forall x\}$ .

## Proposition

*If  $\theta_y$  can be extended irreducibly to  $S$ , then it induces irreducibly from  $S$  back to  $\rho$ .*

Put  $S = \mathrm{Stab}^{G_k}(\theta_y) := \{g \in G_k \mid \theta_y(g^{-1}xg) = \theta_y(x), \quad \forall x\}$ .

## Proposition

*If  $\theta_y$  can be extended irreducibly to  $S$ , then it induces irreducibly from  $S$  back to  $\rho$ .*

*More generally- Suppose that there exists  $\Theta \in \mathrm{Irr}(S)$  whose restriction to  $G_k^r$  lies above  $\theta_y$  and under  $\rho$ . Then the induced representation  $\Theta^{G_k}$  is irreducible, and furthermore*

$$\rho \simeq \Theta^{G_k}.$$

Put  $S = \mathrm{Stab}^{G_k}(\theta_y) := \{g \in G_k \mid \theta_y(g^{-1}xg) = \theta_y(x), \quad \forall x\}$ .

## Proposition

*If  $\theta_y$  can be extended irreducibly to  $S$ , then it induces irreducibly from  $S$  back to  $\rho$ .*

*More generally- Suppose that there exists  $\Theta \in \mathrm{Irr}(S)$  whose restriction to  $G_k^r$  lies above  $\theta_y$  and under  $\rho$ . Then the induced representation  $\Theta^{G_k}$  is irreducible, and furthermore*

$$\rho \simeq \Theta^{G_k}.$$

*In particular  $\dim(\rho) = |G_k : S| \cdot \dim \Theta$ .*

Put  $S = \mathrm{Stab}^{G_k}(\theta_y) := \{g \in G_k \mid \theta_y(g^{-1}xg) = \theta_y(x), \quad \forall x\}$ .

## Proposition

*If  $\theta_y$  can be extended irreducibly to  $S$ , then it induces irreducibly from  $S$  back to  $\rho$ .*

*More generally- Suppose that there exists  $\Theta \in \mathrm{Irr}(S)$  whose restriction to  $G_k^r$  lies above  $\theta_y$  and under  $\rho$ . Then the induced representation  $\Theta^{G_k}$  is irreducible, and furthermore*

$$\rho \simeq \Theta^{G_k}.$$

*In particular  $\dim(\rho) = |G_k : S| \cdot \dim \Theta$ .*

The question of computing the dimension of  $\rho$  is now converted to analysis of  $S$  and of the existence of extensions of  $\theta_y$  to  $S$ .

# Lucky Facts About $\mathfrak{sl}_2$

- ▶ Let  $y \in \mathfrak{g}_{k-r}$  and  $S = \mathrm{Stab}^{G_k}(\theta_y)$ , as above.



# Lucky Facts About $\mathfrak{sl}_2$

- ▶ Let  $y \in \mathfrak{g}_{k-r}$  and  $S = \mathrm{Stab}^{G_k}(\theta_y)$ , as above.
- ▶ Let  $\mathbf{C}_{G_k}(y) := \{g \in G_k \mid \mathrm{Ad}(g)y = y\}$ .

# Lucky Facts About $\mathfrak{sl}_2$

- ▶ Let  $y \in \mathfrak{g}_{k-r}$  and  $S = \mathrm{Stab}^{G_k}(\theta_y)$ , as above.
- ▶ Let  $\mathbf{C}_{G_k}(y) := \{g \in G_k \mid \mathrm{Ad}(g)y = y\}$ .
- ▶ Note the inclusion

$$\mathbf{C}_{G_k}(y) \cdot (G_k^{k-r}) \subseteq S. \quad (**)$$


# Lucky Facts About $\mathfrak{sl}_2$

- ▶ Let  $y \in \mathfrak{g}_{k-r}$  and  $S = \mathrm{Stab}^{G_k}(\theta_y)$ , as above.
- ▶ Let  $\mathbf{C}_{G_k}(y) := \{g \in G_k \mid \mathrm{Ad}(g)y = y\}$ .
- ▶ Note the inclusion

$$\mathbf{C}_{G_k}(y) \cdot (G_k^{k-r}) \subseteq S. \quad (**)$$


- ▶ The following facts are true-
  - ▶ If  $y$  is non-zero modulo  $\mathfrak{p}$ , then  $\mathbf{C}_{G_k}(y)$  is abelian.
  - ▶ In fact,  $\mathbf{C}_{G_k}(y)$  belongs to one of 3 isomorphism types of abelian subgroups of  $G_k$ .
  - ▶ The inclusion  $(**)$  is an equality.

# Extending $\theta_y$ - Even Level




Assume  $k = 2r$ .

# Extending $\theta_y$ - Even Level



Assume  $k = 2r$ . By gluing  $\theta_y$  to any suitable representation of  $\mathbf{C}_{G_k}(y)$ , we obtain the desired extension  $\Theta \in \mathrm{Irr}(S)$ .


# Extending $\theta_y$ - Odd Level



Assume  $k = 2r + 1$ .

# Extending $\theta_y$ - Odd Level

In order to be able to extend  $\theta_y$  to  $S$ , it is necessary to understand the set of representations of  $G^r/G^k$  lying above  $\theta_y$ . This is done using the method of **Heisenberg lifts**.





# The Norm-One Subgroups of Local Division Algebras

# Division Algebras- General Properties

Let  $D$  be a division algebra, with  $\mathbf{Z}(D) = K$  a local field of odd residual characteristic. Let  $\ell = \deg D = (\dim_K D)^{1/2}$ .

Let  $L/K$  be an unramified field extension of degree  $\ell$ .

# Division Algebras- General Properties

Let  $D$  be a division algebra, with  $\mathbf{Z}(D) = K$  a local field of odd residual characteristic. Let  $\ell = \deg D = (\dim_K D)^{1/2}$ .

Let  $L/K$  be an unramified field extension of degree  $\ell$ .

The algebra  $D$  embeds as a subalgebra of the matrix algebra  $M_\ell(L)$ , where an element  $x = x_0 + \nu x_1 + \dots + \nu^{\ell-1} x_{\ell-1}$  is mapped to the matrix

$$\Lambda_x = \begin{pmatrix} x_0 & \pi\sigma(x_{\ell-1}) & \dots & \pi\sigma^{\ell-1}(x_1) \\ x_1 & \sigma(x_0) & \dots & \pi\sigma^{\ell-1}(x_2) \\ \vdots & & \ddots & \vdots \\ x_\ell & \sigma(x_{\ell-1}) & \dots & \sigma^{\ell-1}(x_0) \end{pmatrix},$$

where  $\sigma : L \rightarrow L$  is a generator of the Galois groups  $\mathbf{Gal}(L/K)$ .

# Division Algebras- General Properties

Let  $D$  be a division algebra, with  $\mathbf{Z}(D) = K$  a local field of odd residual characteristic. Let  $\ell = \deg D = (\dim_K D)^{1/2}$ .

Let  $L/K$  be an unramified field extension of degree  $\ell$ .

The algebra  $D$  embeds as a subalgebra of the matrix algebra  $M_\ell(L)$ , where an element  $x = x_0 + \nu x_1 + \dots + \nu^{\ell-1} x_{\ell-1}$  is mapped to the matrix

$$\Lambda_x = \begin{pmatrix} x_0 & \pi\sigma(x_{\ell-1}) & \dots & \pi\sigma^{\ell-1}(x_1) \\ x_1 & \sigma(x_0) & \dots & \pi\sigma^{\ell-1}(x_2) \\ \vdots & & \ddots & \vdots \\ x_\ell & \sigma(x_{\ell-1}) & \dots & \sigma^{\ell-1}(x_0) \end{pmatrix},$$

where  $\sigma : L \rightarrow L$  is a generator of the Galois groups  $\mathbf{Gal}(L/K)$ .

Define  $\text{Nrd}(x) = \det \Lambda_x$  and  $\text{Trd}(x) = \text{Tr} \Lambda_x$ .

# The Norm-One Subgroup of $D^\times$

Let  $\mathfrak{o}_L$  denote the valuation ring of  $L$ , and let  $\mathcal{O} := D \cap \mathbf{M}_\ell(\mathfrak{o}_L)$ .

# The Norm-One Subgroup of $D^\times$

Let  $\mathfrak{o}_L$  denote the valuation ring of  $L$ , and let  $\mathcal{O} := D \cap \mathbf{M}_\ell(\mathfrak{o}_L)$ . Let

$$G = \mathrm{SL}_1(D) = \{x \in D \mid \mathrm{Nrd}(x) = 1\} \subseteq \mathcal{O}^\times.$$

The congruence subgroups of  $G$  are defined by

$$G^m := G \cap (1 + \nu^m \mathbf{M}_\ell(\mathfrak{o}_L)),$$

where  $\nu$  is the element of  $\mathcal{O}$  which is identified with the matrix

$$\begin{pmatrix} 0 & & \cdots & \pi \\ 1 & 0 & & \vdots \\ & \ddots & \ddots & \\ & & 1 & 0 \end{pmatrix} \in \mathbf{M}_\ell(\mathfrak{o}_L).$$

The **level** of an irreducible representation  $\rho \in \text{Irr}(G)$  is defined as in the case of  $\text{SL}_2(\mathfrak{o})$ , i.e. the minimal  $m \in \mathbb{N}$  such that  $G^{m+1} \subseteq \text{Ker}(\rho)$ .

The **level** of an irreducible representation  $\rho \in \text{Irr}(G)$  is defined as in the case of  $\text{SL}_2(\mathfrak{o})$ , i.e. the minimal  $m \in \mathbb{N}$  such that  $G^{m+1} \subseteq \text{Ker}(\rho)$ .

Owing to the work of H. Koch, the centralizer in  $G_k$  of an element  $y \in G_k^r$  can be effectively described. An explicit description of the centralizer was presented in the work of V. Singh.



The **level** of an irreducible representation  $\rho \in \text{Irr}(G)$  is defined as in the case of  $\text{SL}_2(\mathfrak{o})$ , i.e. the minimal  $m \in \mathbb{N}$  such that  $G^{m+1} \subseteq \text{Ker}(\rho)$ .

Owing to the work of H. Koch, the centralizer in  $G_k$  of an element  $y \in G_k^r$  can be effectively described. An explicit description of the centralizer was presented in the work of V. Singh. In particular- in the case where  $\ell = \deg D$  is a prime number, all such centralizers are of the form

$$((L')^\times \cap G) \cdot G^{m'},$$

where  $L' \supseteq K$  is a field extension, and  $m' \in \mathbb{N}$  is dependent on  $y$ .

The case where  $\ell$  is prime is amenable to a similar treatment as  $\text{SL}_2(\mathfrak{o})$ .

# Results

Let  $D/K$  be a division algebra of prime degree  $\ell$ , and put  $\epsilon = \gcd(\ell, q - 1) = \#\mu_\ell(K)$ .

## Theorem

1. *There exist rational polynomials  $a_m^\epsilon(t), d_m^\epsilon(t) \in \mathbb{Q}(t)$  ( $m \in \mathbb{N}_0$ ), such that for any  $m \in \mathbb{N}_0$ ,  $\mathrm{SL}_1(D)$  has exactly  $a_m^\epsilon(q)$  irreducible representations of level  $m$ . All such representations are of dimension  $d_m^\epsilon(q)$ .*

# Theorem

2.

$$\begin{aligned} \zeta_{\mathrm{SL}_1(D)}(s) &= \zeta_{\mathrm{SL}_1(\mathbb{F}_{q^\ell}|\mathbb{F}_q)}(s) + \sum_{m=1}^{\infty} a_m^\epsilon(q) d_m^\epsilon(q)^{-s} \\ &= \frac{\frac{q^\ell-1}{q-1} \left(1 - q^{-\binom{\ell}{2}s}\right) + \left(\frac{q^\ell-1}{\epsilon(q-1)}\right)^{-s} \epsilon^2 (q-1) \left(\sum_{\lambda=0}^{\ell-2} q^{\lambda(1-\frac{\ell-1}{2}s)}\right)}{1 - q^{(\ell-1)-\binom{\ell}{2}s}}. \end{aligned}$$

# Theorem

2.

$$\begin{aligned} \zeta_{\mathrm{SL}_1(D)}(s) &= \zeta_{\mathrm{SL}_1(\mathbb{F}_{q^\ell}|\mathbb{F}_q)}(s) + \sum_{m=1}^{\infty} a_m^\epsilon(q) d_m^\epsilon(q)^{-s} \\ &= \frac{\frac{q^\ell-1}{q-1} \left(1 - q^{-\binom{\ell}{2}s}\right) + \left(\frac{q^\ell-1}{\epsilon(q-1)}\right)^{-s} \epsilon^2 (q-1) \left(\sum_{\lambda=0}^{\ell-2} q^{\lambda(1-\frac{\ell-1}{2}s)}\right)}{1 - q^{(\ell-1)-\binom{\ell}{2}s}}. \end{aligned}$$

3. *If  $\ell > 2$  then all irreducible representations of  $\mathrm{SL}_1(D)$  are induced from one-dimensional representations of congruence subgroups.*

# Regular Representations

# Generalizing to Other Groups

Let  $\mathbf{G} \subseteq \mathrm{GL}_n$  be a connected reductive algebraic group, which is defined over  $\mathfrak{o}$ . Put  $G = \mathbf{G}(\mathfrak{o})$ .

# Generalizing to Other Groups

Let  $\mathbf{G} \subseteq \mathrm{GL}_n$  be a connected reductive algebraic group, which is defined over  $\mathfrak{o}$ . Put  $G = \mathbf{G}(\mathfrak{o})$ .

The congruence subgroups of  $G$  are defined by

$$G^m := G \cap (1 + \pi^m \mathbf{M}_n(\mathfrak{o})), \quad (m \in \mathbb{N}),$$

and under some mild assumptions on  $p$ , we have that

$$G_m := G/G^m \cong \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m).$$

# Generalizing to Other Groups

Let  $\mathbf{G} \subseteq \mathrm{GL}_n$  be a connected reductive algebraic group, which is defined over  $\mathfrak{o}$ . Put  $G = \mathbf{G}(\mathfrak{o})$ .

The congruence subgroups of  $G$  are defined by

$$G^m := G \cap (1 + \pi^m \mathbf{M}_n(\mathfrak{o})), \quad (m \in \mathbb{N}),$$

and under some mild assumptions on  $p$ , we have that

$$G_m := G/G^m \cong \mathbf{G}(\mathfrak{o}/\mathfrak{p}^m).$$

We retain the notation  $G_m^r$  for the subgroups  $G^r/G^m$ .



# Generalizing to Other Groups

By virtue of  $\mathbf{G}$  being reductive, it is possible to apply the process described above for  $\mathbf{G} = \mathrm{SL}_2$  and to associate any representation  $\rho \in \mathrm{Irr}(G)$  of level  $m$  with an orbit in a finite Lie-ring  $\mathfrak{g}_{m-r}$ , as above.

# Generalizing to Other Groups

By virtue of  $\mathbf{G}$  being reductive, it is possible to apply the process described above for  $\mathbf{G} = \mathrm{SL}_2$  and to associate any representation  $\rho \in \mathrm{Irr}(G)$  of level  $m$  with an orbit in a finite Lie-ring  $\mathfrak{g}_{m-r}$ , as above.

However, the description of stabilizers of the associated characters  $\theta_y \in \mathrm{Irr}(G_m^r)$  is seldom as straightforward as in the previous cases.

# Regular Representations

Let  $\mathfrak{g}$  be the Lie algebra of  $\mathbf{G}(\mathfrak{o})$ , and put  $\mathfrak{g}^m := \pi^m \mathfrak{g}$  and  $\mathfrak{g}_m = \mathfrak{g}/\mathfrak{g}^m$ .

## Definition (Steinberg, Hill)

- ▶ An element  $y \in \mathfrak{g}_1 = \mathfrak{g}(\mathbb{F}_q)$  is called regular if its centralizer  $\mathbf{C}_{\mathbf{G}(\mathbb{F}_q)}(y)$  in  $G_1$  is of minimal dimension.
- ▶ An element  $y \in \mathfrak{g}_m$  is called regular if its image in  $\mathfrak{g}_1$  is regular.
- ▶ A representation  $\rho \in \text{Irr}(G)$  of level  $m$  is called regular, if the orbit associated to it is regular.

# Regular Representations

For example, in the case  $\mathbf{G} = \mathrm{GL}_n$  it is true that an element  $y \in \mathfrak{gl}_n(\mathfrak{o}/\mathfrak{p}^m)$  is regular if and only if its minimal polynomial is equal to its characteristic polynomial.

# Regular Representations

For example, in the case  $\mathbf{G} = \mathrm{GL}_n$  it is true that an element  $y \in \mathfrak{gl}_n(\mathfrak{o}/\mathfrak{p}^m)$  is regular if and only if its minimal polynomial is equal to its characteristic polynomial.

It is also true in this case that the centralizer  $\mathbf{C}_{G_m}(y)$  is an abelian group, and in fact equals to the group

$$(\mathfrak{o}/\mathfrak{p}^m)[y] \cap \mathrm{GL}_n(\mathfrak{o}/\mathfrak{p}^m).$$

# Regular Representations

For example, in the case  $\mathbf{G} = \mathrm{GL}_n$  it is true that an element  $y \in \mathfrak{gl}_n(\mathfrak{o}/\mathfrak{p}^m)$  is regular if and only if its minimal polynomial is equal to its characteristic polynomial.

It is also true in this case that the centralizer  $\mathbf{C}_{G_m}(y)$  is an abelian group, and in fact equals to the group

$$(\mathfrak{o}/\mathfrak{p}^m)[y] \cap \mathrm{GL}_n(\mathfrak{o}/\mathfrak{p}^m).$$

In addition, if  $y$  is taken to be in the orbit associated to a regular representation  $\rho \in \mathrm{Irr}(\mathrm{GL}_n(\mathfrak{o}))$  of level  $m$ , and  $\theta_y \in \mathrm{Irr}(G_m^r)$  is the associated character, then

$$\mathrm{Stab}^{G_m}(\theta_y) = \mathbf{C}_{G_m}(y) \cdot \mathbf{G}_m^{m-r},$$

as in the case of  $\mathrm{SL}_2$ .

# Regular Representation

Using this stabilizer structure, G. Hill managed to produce an explicit description of all regular characters of  $GL_n(\mathfrak{o})$ .

# Regular Representation

Using this stabilizer structure, G. Hill managed to produce an explicit description of all regular characters of  $GL_n(\mathfrak{o})$ .

Recent works (KOS, Stasinsky, Takase, S.) have applied the construction of regular representations to the study of representation growth, and attempted to generalize the construction to other families of reductive groups.



# Regular Representation

Using this stabilizer structure, G. Hill managed to produce an explicit description of all regular characters of  $GL_n(\mathfrak{o})$ .

Recent works (KOS, Stasinsky, Takase, S.) have applied the construction of regular representations to the study of representation growth, and attempted to generalize the construction to other families of reductive groups.

A possible application to the study of regular representation is to the analysis of the regular part of the representation zeta function of  $G$ , in view of the decomposition-

$$\zeta_G(s) = \zeta_{G(\mathbb{F}_q)}(s) + \zeta_G^{\text{reg.}}(s) + \zeta_G^{\text{irreg.}}(s).$$

# Regular Zeta Functions of Groups of Type $A_{n-1}$

Let  $\mathbf{G}_{+1} = \mathrm{GL}_n$  and  $\mathbf{G}_{-1} = \mathrm{U}_n$  the unitary group attached to an unramified quadratic extension  $L/K$ . Let  $\epsilon \in \{\pm 1\}$ .

# Regular Zeta Functions of Groups of Type $A_{n-1}$

Let  $\mathbf{G}_{+1} = \mathrm{GL}_n$  and  $\mathbf{G}_{-1} = \mathrm{U}_n$  the unitary group attached to an unramified quadratic extension  $L/K$ . Let  $\epsilon \in \{\pm 1\}$ .

As was recently shown by Krokovski, Onn and Singla, there exist

- ▶  $\mathcal{A}_n$  a 'nice' set,
- ▶ an explicit map from the regular part of  $\mathrm{Irr}^m(\mathbf{G}_\epsilon(\mathfrak{o}))$  to  $\mathcal{A}_n$ ,

# Regular Zeta Functions of Groups of Type $A_{n-1}$

Let  $\mathbf{G}_{+1} = \mathrm{GL}_n$  and  $\mathbf{G}_{-1} = \mathrm{U}_n$  the unitary group attached to an unramified quadratic extension  $L/K$ . Let  $\epsilon \in \{\pm 1\}$ .

As was recently shown by Krokovski, Onn and Singla, there exist

- ▶  $\mathcal{A}_n$  a 'nice' set,
- ▶ an explicit map from the regular part of  $\mathrm{Irr}^m(\mathbf{G}_\epsilon(\mathfrak{o}))$  to  $\mathcal{A}_n$ ,
- ▶ polynomials  $u_\epsilon^\tau(t)$ , for all  $\tau \in \mathcal{A}_n$

such that, if  $\rho \in \mathrm{Irr}^m(\mathbf{G}_\epsilon(\mathfrak{o}))$  is mapped to  $\tau \in \mathcal{A}_n$  then

$$\dim(\rho) = q^{m\binom{n}{2}} \left( \frac{|\mathbf{G}_\epsilon(\mathbb{F}_q)|}{u_\epsilon^\tau(q)} \right).$$

The number of regular representations of a given type  $\tau \in \mathcal{A}_n$  can also be described, using the polynomial  $w_d(q)$ , which gives the number of irreducible polynomials of degree  $d$  over  $\mathbb{F}_q$ .

The number of regular representations of a given type  $\tau \in \mathcal{A}_n$  can also be described, using the polynomial  $w_d(q)$ , which gives the number of irreducible polynomials of degree  $d$  over  $\mathbb{F}_q$ . From this, one obtains the regular zeta function of  $\mathbf{G}_\epsilon(\mathfrak{o})$ .

Theorem (Krakovski, Onn, Singla, '16)

$$\zeta_{\mathbf{G}_\epsilon(\mathfrak{o})}^{\text{reg.}}(s) = \frac{1}{1 - q^{n - \binom{n}{2}s}} \cdot \sum_{\tau \in \mathcal{A}_n} u_\epsilon^\tau(q) \prod_{i=1}^n \binom{\sum_j \tau_{i,j}}{\tau_{i,1}, \tau_{i,2}, \dots, \tau_{i,n}} \binom{w_d(q)}{\sum_j \tau_{i,j}} \left( \frac{|\mathbf{G}_\epsilon(\mathbb{F}_q)|}{u_\epsilon^\tau(q)} \right)^{-s}.$$

# Regular Zeta Function for Classical Groups

A similar analysis has been thus far accomplished for classical groups defined over  $\mathfrak{o}$ , i.e. linear algebraic groups of the form

$$\{\mathbf{X} = (x_{i,j}) \in M_n \mid \mathbf{X}^T \mathbf{J} \mathbf{X} = \mathbf{J}\},$$

where  $\mathbf{J} \in GL_n(\mathfrak{o})$  is a fixed symmetric or anti-symmetric matrix.

# Regular Zeta Function for Classical Groups

A similar analysis has been thus far accomplished for classical groups defined over  $\mathfrak{o}$ , i.e. linear algebraic groups of the form

$$\{\mathbf{X} = (x_{i,j}) \in M_n \mid \mathbf{X}^T \mathbf{J} \mathbf{X} = \mathbf{J}\},$$

where  $\mathbf{J} \in \mathrm{GL}_n(\mathfrak{o})$  is a fixed symmetric or anti-symmetric matrix. In the specific cases of the symplectic group  $\mathrm{Sp}_{2n}$  and the odd-orthogonal group  $\mathrm{SO}_{2n+1}$ , the regular representation zeta function has been computed.



Questions?

Thank You.