

# Kidney transplantation, rejection prophylaxis

## Adults:

Mycophenolate mofetil: *Oral, IV*:

1 g (range: 500 mg to 1 g) twice daily as part of an appropriate combination regimen.

Mycophenolate sodium, delayed release: *Oral*:

720 mg (range: 360 to 720 mg) twice daily as part of an appropriate combination regimen <sup>[1]</sup>

## Pediatrics:

Mycophenolate mofetil: Infants  $\geq 3$  months and children: *Oral*:

→ Suspension: 600 mg/m<sup>2</sup>/dose twice daily; maximum daily dose: 2,000 mg/day.

→ Tablets or capsules:

- BSA 1.25 m<sup>2</sup> to 1.5 m<sup>2</sup>: 750 mg twice daily.
- BSA  $\geq 1.5$  m<sup>2</sup>: 1,000 mg twice daily.

Mycophenolate sodium delayed release tablets: Children >5 years:

400 mg/m<sup>2</sup>/dose twice daily; maximum daily dose: 1,440 mg/day:

- BSA <1.19 m<sup>2</sup>: Use of this formulation is not recommended.
- BSA 1.19 to 1.58 m<sup>2</sup>: 540 mg twice daily.
- BSA >1.58 m<sup>2</sup>: 720 mg twice daily.

---

[ 1 ] - Hardinger K, Brennen DC. Kidney transplantation in adult: maintenance immunosuppressive therapy. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed May 18, 2021.

