



## Severe Adverse Effects

- → pericardial effusion / Delayed / 0-5.0
- → myocardial infarction / Delayed / 0-5.0
- → arrhythmia exacerbation / Early / 0-5.0
- → atrial fibrillation / Early / 0-5.0
- → cardiac arrest / Early / 0-5.0
- → pericarditis / Delayed / 0-5.0
- → bronchospasm / Rapid / 0-5.0
- → pleural effusion / Delayed / 0-5.0
- → agranulocytosis / Delayed / 0-5.0
- $\rightarrow$  GI bleeding / Delayed / 0-5.0
- → thrombosis / Delayed / 0-5.0

## Moderate Adverse Effects

- → antibody formation / Delayed / 1.0-61.0
- → hyperlipidemia / Delayed / 7.0-7.0
- → hypercholesterolemia / Delayed / 6.0-6.0
- → peripheral edema / Delayed / 0-5.0
- → encephalopathy / Delayed / 0-5.0
- → chest pain (unspecified) / Early / 0-5.0
- → hypertension / Early / 5.0-5.0
- → sinus tachycardia / Rapid / 0-5.0
- $\rightarrow$  palpitations / Early / 0-5.0
- → impaired wound healing / Delayed / 0-5.0
- $\rightarrow$  dyspnea / Early / 0-5.0
- → subdural hematoma / Early / 0-5.0
- → polycythemia / Delayed / 0-5.0
- → cystitis / Delayed / 0-5.0 [1]

