

VERTEBRAL FRACTURES

- Consider Physical Therapy and Exercise
- Calcitonin significantly reduces pain and facilitates earlier mobilization for up to four weeks
- Medications facilitate patient mobility and participation in physical therapy, and should be tapered slowly as pain improves
- consider Physical Therapy and Exercise
- Acetaminophen: 500 to 1,000 mg every four to eight hours (maximum 3 g per day)
- Calcitonin: 200 IU per day intra-nasally
- Lidocaine 5% patch: Apply to affected area for 12 hours
- Muscle relaxants (e.g., cyclobenzaprine): 10 mg every eight hours

First: oral Bisphosphonates

- Alendronate (oral 5 or 10 mg once daily, or 70 mg once weekly),
- Ibandronate 150mg per month
- risedronate (oral 5 mg daily or 35 mg one weekly)

Then consider IV bisphosphonates

- zoledronate (intravenous infusion 5 mg once yearly)
- Clacitonin: 50 to 100 IU per day intramuscularly or 200 IU per day intra-nasally
- calcium (1200 mg/d or 500 mg twice daily)
- Vitamin D analogs: alfacalcidol – calcitriol
- Estrogen agonist/antagonist (raloxifene): 60mg / day



second line treatment or incase patient is intolerant to biphosphonates

- Parathyroid hormone (teriparatide): 20 mcg per day subcutaneously for up to 24 months
- RANKL inhibitor (denosumab): 60 mg per day subcutaneously every six months

Hormone replacement therapy

Hormone replacement therapy and testosterone therapy have therefore been found to increase lumbar spine BMD in hypogonadal patients on glucocorticoid therapy.

-
- [1]- Wong, S., & Mok, C. C. (2020). Management of glucocorticoid-related osteoporotic vertebral fracture. Osteoporosis and sarcopenia, 6(1), 1–7.
<https://doi.org/10.1016/j.afos.2020.02.002>
 - [2]- McCarthy J, Davis A. Diagnosis and Management of Vertebral Compression Fractures. Am Fam Physician. 2016 Jul 1;94(1):44-50. PMID: 27386723.
 - [3]- McDonald FD, Horensten ML, Mayor GB, Turcotte JG, Selezinka W, Schork MA. Effect of alternate-day steroids on renal transplant function. A controlled study. Nephron. 1976;17(6):415-29. doi: 10.1159/000180750. PMID: 796739.
 - [4]- Baker, R., Mark, P., Patel, R., Stevens, K., & Palmer, N. (2017). Renal association clinical practice guideline in post-operative care in the kidney transplant recipient. BMC Nephrology, 18(1). doi: 10.1186/s12882-017-0553-2
 - [5]- Buckley, L., Guyatt, G., Fink, H., Cannon, M., Grossman, J., & Hansen, K. et al. (2017). 2017 American College of Rheumatology Guideline for the Prevention and Treatment of Glucocorticoid-Induced Osteoporosis. Arthritis & Rheumatology, 69(8), 1521-1537. doi: 10.1002/art.40137

