

HYPERTENSION

Pretransplant

- **Beta-adrenergic agonists, clonidine** and beta-blockers are commonly associated with this rebound hypertension , particularly when abruptly stopped.
- Sympathetic overactivity without renin angiotensin system mediation plays a major role in the mechanism of rebound HTN from clonidine .
- Similar to clonidine, **rebound HTN** from beta-blockers **causes elevated BP and heart rate. It also leads to cardiac events including angina, myocardial infarction, or sudden cardiac death in patients with underlying coronary artery disease (CAD) .**
- .Given the cardiac risk with peri-operative beta-blocker withdrawal the 2014 ACC/AHA guidelines give a class I **recommendation** with B level evidence for the **continuation of beta-blockers during the peri-operative period in patients who chronically use them(3)**



Post Renal transplant management

TABLE 2 | Common interventions for post-kidney transplant hypertension.

Blood pressure managements	Interventions	Comments
Lifestyle modifications	Diet Exercise Stress reduction	Required for all patients
Pharmacological therapy	Antihypertensive medication <ul style="list-style-type: none"> • Diuretics <ul style="list-style-type: none"> - Loop - Thiazides • Calcium channel blockers • Beta-blockers • Renin-Angiotensin-Aldosterone System blockade <ul style="list-style-type: none"> - Angiotensin-converting enzyme inhibitors - ARB, angiotensin II receptor blockers - Mineralocorticoid receptor antagonists • Alpha₁ antagonists • Alpha₂ agonists 	Choice of medications depending on: <ul style="list-style-type: none"> • Patients' characteristics, • Tolerability • Medication side effect profiles
Procedural or surgical interventions	Specific treatment modalities <ul style="list-style-type: none"> • Transplant renal artery angioplasty ± stenting • Continuous positive airway pressure (CPAP) • Bilateral native nephrectomy • Native renal denervation 	Etiologies of resistant hypertension <ul style="list-style-type: none"> • Transplant renal artery stenosis • Obstructive sleep apnea (OSA) • Failed native kidney • Sympathetic overactivity



- glucocorticoid-induced hypertension.
- The optimal regimen is certainly **not clear** given the potential impact of glucocorticoids on several hypertensive pathways, including increased vascular tone, increased sodium reabsorption, and centrally mediated mechanisms.
- It is proposed that using **calcium channel blockers** or other vascular relaxants if hypertension is manifest within the first few days of glucocorticoid administration with the eventual addition of a **diuretic** if the hypertension persists or worsens.
- There is potentially a role for centrally acting agents such as **clonidine**, either in addition to other pharmacotherapy or independently, although this is less clear^[2]

- High-dose IV steroids as more than 20 mg of prednisone per day can contribute to HTN during immediate post-transplant period.
- **steroid avoidance or withdrawal (SAW)** maintenance immunosuppressive medication regimens can be considered.

- However, the **effect of SAW** on post-transplant HTN has yielded conflicting data.
- A systematic review and meta-analysis revealed that steroid avoidance or withdrawal significantly decreases CV outcomes including HTN but increases risk of acute rejection .
- It is common for steroids to be reintroduced after a diagnosis of acute rejection in recipients who were initially managed with a SAW regimen.
- A randomized control trial, however, demonstrated no difference in blood pressure change between alternative day and daily prednisone .
- So **SAW protocols** should be considered in select patients, specifically those who would be at greater risk for CV outcomes but be immunologically lower risk of rejection



PHARMACOLOGICAL MANAGERMENTS

Use of Antihypertensive Medications

TABLE 3 | Summarized common antihypertensive medications used in kidney transplant patients.

Antihypertensive classes	Pros	Comments	Cons	Comments
Diuretics				
Loop	<ul style="list-style-type: none"> - Generally, not the first line antihypertensive medication Used in CsA treated recipients - Used for volume control - May use with ACEI or ARB in TRAS 	<ul style="list-style-type: none"> - Renal sodium excretion defect in CsA-induced HTN (198) 	<ul style="list-style-type: none"> - Loop diuretic may worsen renal allograft function from redistribution of decreased renal blood flow at juxtamedullary cortex and outer medulla (199) and ↓ oxygenation in medulla due to decreased cortical vascular resistance diverting medullary perfusion (200). - Electrolyte disturbance 	<ul style="list-style-type: none"> - Hyponatremia - Hypomagnesemia - Hyperuricemia
Thiazide	<ul style="list-style-type: none"> - May consider in hypomagnesemic patients who needs volume control from diuresis (154). - May consider in salt-sensitive HTN from CNIs - May use with ACEI or ARB in TRAS 	<ul style="list-style-type: none"> - Not the first line antihypertensive medication - WNK-SPAK-NCC pathway 	<ul style="list-style-type: none"> - Potential volume depletion - Hyperlipidemia - Electrolyte disturbance 	<ul style="list-style-type: none"> - Hyponatremia - Hypomagnesemia - Hyperuricemia
CCB	<ul style="list-style-type: none"> - May improve renal allograft function (201) and lower DGF but inconclusive 	<ul style="list-style-type: none"> - Afferent arteriolar vasodilatation (202) 	<ul style="list-style-type: none"> - Non-dihydropyridine CCB is CYP450 inhibitor and increases CNI level 	<ul style="list-style-type: none"> - CYP450 3A4 enzyme inhibitor → ↑CNI and mTOR inhibitors level
ACEI/ARB	<ul style="list-style-type: none"> - Anti-proteinuric - Cardioprotection (203, 204) - May use with diuretic in TRAS 		<ul style="list-style-type: none"> - Hyperkalemia - Anemia - Elevated creatinine 	<ul style="list-style-type: none"> - In the setting of volume depletion, TRAS
Beta-blockers	<ul style="list-style-type: none"> - For cardioprotection 		<ul style="list-style-type: none"> - Mask symptoms of hypoglycemia and thyrotoxicosis - Worsening lipid profiles - Hyperkalemia - Potential rebound HTN - Hyperkalemia 	<ul style="list-style-type: none"> - Especially with mTOR inhibitors
Mineralocorticoid receptor antagonists	<ul style="list-style-type: none"> - Systolic dysfunction 	<ul style="list-style-type: none"> - Safe with using ACEI and ARB 		
Alpha ₁ antagonist	<ul style="list-style-type: none"> - Comparable to ACEI for BP control - Generally, not the first line antihypertensive agents 	<ul style="list-style-type: none"> - May need to add other antihypertensive agents 		
Alpha ₂ agonist	<ul style="list-style-type: none"> - Lower plasma renin activity that modulated renal vascular resistance and subsequently lower MAP (140). - No change in GFR and effective renal plasma flow (141) 		<ul style="list-style-type: none"> - Potential rebound HTN 	<ul style="list-style-type: none"> - Need to be slowly tapered off if medication discontinuation is needed.

ACEI, angiotensin-converting enzyme inhibitors; ARB, angiotensin receptor blockers; CCB, calcium channel blockers; CNI, calcineurin inhibitor; CYP, cytochrome; DGF, delayed graft function; GFR, glomerular filtration rate; mTOR, mammalian target of rapamycin; TRAS, transplant renal artery stenosis; WNK-SPAK-NCC, WNK, With-No-K(Lys)—STE20/SPS1-related proline/alanine-rich kinase—Sodium Chloride Cotransporter.



OTHER MANAGEMENT FOR BLOOD PRESSURE FOR RENAL TRANSPLANT RECIPIENTS:

1. Native Nephrectomy and ACES
2. Native Renal Sympathetic Denervation
3. Surgical Renal Denervation by Bilateral Native Nephrectomy

→ ALL IN ALL There is no conclusive BP target for kidney transplant recipients and **therapy targets need to be individualized**. Further research is needed ^[1]

[1]- Tantisattamo, E., Molnar, M., Ho, B., Reddy, U., Dafoe, D., & Ichii, H. et al. (2020). Approach and Management of Hypertension After Kidney Transplantation. *Frontiers In Medicine*, 7. doi: 10.3389/fmed.2020.00229

[2]- Goodwin, J.E., Geller, D.S. Glucocorticoid-induced hypertension. *Pediatr Nephrol* 27, 1059–1066 (2012). <https://doi.org/10.1007/s00467-011-1928-4>

[3]- Tantisattamo, E., Molnar, M., Ho, B., Reddy, U., Dafoe, D., & Ichii, H. et al. (2020). Approach and Management of Hypertension After Kidney Transplantation. *Frontiers In Medicine*, 7. doi: 10.3389/fmed.2020.00229

