

Severe Adverse Effects

- hyperkalemia / Delayed / 7.0-18.0
- lymphopenia / Delayed / 0-18.0
- hyperglycemia / Delayed / 0-17.0
- anemia / Delayed / 0-15.0
- hypertensive crisis / Early / 0-10.0
- atrial fibrillation / Early / 0-10.0
- heart failure / Delayed / 0-10.0
- pulmonary edema / Early / 0-10.0
- pleural effusion / Delayed / 0-10.0
- hypophosphatemia / Delayed / 0-10.0
- renal failure (unspecified) / Delayed / 2.0-10.0
- thromboembolism / Delayed / 0-10.0
- pulmonary embolism / Delayed / 0-10.0
- wound dehiscence / Delayed / 1.0-10.0
- pancytopenia / Delayed / 0-10.0
- osteonecrosis / Delayed / 0-10.0
- prostatic hypertrophy / Delayed / 0-10.0
- stomatitis / Delayed / 0-9.0
- angioedema / Rapid / 0-6.8
- fever / Early / 0-6.0
- hypokalemia / Delayed / 0-6.0
- fatigue / Early / 0-6.0
- agitation / Early / 0-5.0
- anxiety / Delayed / 0-5.0 ^[1]

[1] - Afinitor (everolimus) dose, indications, adverse effects, interactions... from PDR.net;
<https://www.pdr.net/drug-summary/Afinitor-everolimus-416>, accessed: 2021-11-25

