

HEMATOLOGICAL

Monitor

- Monitoring daily CBC , Total white blood cell and platelet counts
- Low counts of platelets and white blood cells (including low counts of lymphocytes and neutrophils) have been identified and are reversible following dose adjustments.
- If available monitor CD3 (+) T-Cell. Keeping CD3 (+) T-Cell $<0.05 \times 10^9/L$ ($<50/\text{microliter}$) has been a useful index to avoid excessive doses
- Also If available measure total lymphocyte count as it can be used. Total lymphocyte count below $0.3 \times 10^9/l$ has been found to be a useful index for using ATG ^[1]

Management

- The THYMOGLOBULIN dose should be reduced by one-half if the white blood cell (WBC) count is between 2,000 and 3,000 cells/mm³ or if the platelet count is between 50,000 and 75,000 cells/mm³. Stopping THYMOGLOBULIN treatment should be considered if the WBC count falls below 2,000 cells/mm³ or if the platelet count falls below 50,000 cells/mm³. ^[1]



HEMOLYSIS

- can usually be detected only in the laboratory. Clinically significant hemolysis has been reported rarely. Appropriate treatment of hemolysis may include transfusion of erythrocytes; if necessary, administer intravenous mannitol, furosemide, sodium bicarbonate, and fluids. Severe and unremitting hemolysis may require discontinuation of therapy with ATGAM ^[2]

THROMBOCYTOPENIA

- Thrombopoietin receptor agonist, romiplostim, and eltrombopag have been successfully used for treatment of thrombocytopenia in chronic idiopathic thrombocytopenic purpura ^[1]
- Thrombocytopenia is usually transient in renal transplant patients; platelet counts generally return to adequate levels without discontinuing therapy with ATGAM. ^[2]

NEUTROPENIA

- If neutropenia persists despite modification of immunosuppressive medications, then one may opt for colony-stimulating factors to increase white cell counts ^[1]
- Granulocyte-Monocyte Colony-Stimulating Factor (GM-CSF) is another stimulating agent which activates neutrophil, monocytes, macrophages, and dendritic cells and it has a proinflammatory profile unlike G-CSF ^[2]
- Platelet transfusions may be necessary in patients with aplastic anemia. ^[2]

^[1]-Muhammad Abdul Mabood Khalil, Muhammad Ashhad Ullah Khalil, Taqi F. Taufeeq Khan, Jackson Tan, "Drug-Induced Hematological Cytopenia in Kidney Transplantation and the Challenges It Poses for Kidney Transplant Physicians", Journal of Transplantation, vol. 2018, Article ID 9429265, 22 pages, 2018. <https://doi.org/10.1155/2018/9429265>

^[2]-Atgam® (lymphocyte immune globulin, anti-thymocyte globulin [equine] sterile solution). (2021). Retrieved 11 November 2021, from <https://www.fda.gov/media/78206/download>

