

## Severe Adverse Effects

- anaphylactoid reactions / Rapid / 0-1.0
- serum sickness / Delayed / 0.2-1.0
- GI obstruction / Delayed / 0.2
- pulmonary edema / Early / 0.2
- bradycardia / Rapid / 0.2
- hemolytic anemia / Delayed / 0.2

## Moderate Adverse Effects

- antibody formation / Delayed / 7.0-62.0
- elevated hepatic enzymes / Delayed / 17.0-51.0
- infusion-related reactions / Rapid / 18.0-20.0
- hypertension / Early / 0-7.0
- neutropenia / Delayed / 7.0-7.0
- candidiasis / Delayed / 5.0-5.0
- hypotension / Rapid / 0.2-1.0
- chest pain (unspecified) / Early / 1.0-1.0
- dyspnea / Early / 1.0-1.0 dehydration / Delayed / 0.2
- constipation / Delayed / 0.2
- edema / Delayed / 0.2
- thrombocytopenia / Delayed / 0.2
- anemia / Delayed / 0.2
- phlebitis / Rapid / 0.2
- leukopenia / Delayed / 0.2
- lymphadenopathy / Delayed / 0.2

---

[ 1 ] - Remicade (infliximab) dose, indications, adverse effects, interactions... from PDR.net

<https://www.pdr.net/drug-summary/Remicade-infliximab-263.4292#12>, accessed: 2021-11-28

