

SEMINAR ON MODULI THEORY

August 27, 2020

NEERAJ DESHMUKH

1. PLAN

One way to go through these sessions is go recall theory of schemes (at level of things in Hartshorne chapter two), by doing a beeline through all the definitions, properties, etc. However, I feel since the point of doing this exercise is to become more comfortable in working with scheme, we will just do lots of examples instead. By this, I mean we will just to prove some things in very concrete situations. This will help you build a concrete picture of the generalities.

List of some things to discuss (just do lots of examples):

- (1) Definition of a scheme.
- (2) Say affine communication lemma *stress this!*
- (3) examples
 - (a) \mathbb{P}^n and it sheaf theory! This already clarifies the $\mathcal{O}(n)$'s

Things to say: finite-generation, valuative criteria for \mathbb{P}^1 (using DVR's - how giving a point and it's specialisation is the same as giving a map $\text{Spec } R \rightarrow \mathbb{P}^1$).

List of examples:

- (1) \mathbb{A}^1 with a double point (what are quasi-coherent sheaves on this?)
- (2) $\mathbb{A}^2 \setminus \{0, 0\}$ (what is the structure sheaf?).
- (3) \mathbb{P}^1 (its structure sheaf!).
- (4) $V_+(x^2 + y^2 + z^2)$ over \mathbb{R} and \mathbb{C} . The point is that there is a change of coordintes which
- (5) Blow-up of \mathbb{A}^2 at a point. (because everyone should know about blow-ups!)
- (6) $\text{Spec } R[x_1, x_2, \dots]$ as an example of something non-noetherian.
- (7) An example of a scheme without a closed point.

List of morphisms:

- (1) $x \mapsto x^2$ (more, generally x^n). This covers ramified, finitely presented, flat.
- (2) a non-quasi-compact open-immersion. Polynomial ring in infinitely many variables and knock off the origin. Also, the origin of is an example of something

2. RIGHT AT THE BEGINNING..

For the sake of completeness we begin by reviewing the definition of a locally ringed space.

Definition 2.1. Locally ringed spaces.

- (1) A *locally ringed space* (X, \mathcal{O}_X) is a pair consisting of a topological space X and a sheaf of rings \mathcal{O}_X all of whose stalks are local rings.
- (2) Given a locally ringed space (X, \mathcal{O}_X) we say that $\mathcal{O}_{X,x}$ is the *local ring of X at x* . We denote $\mathfrak{m}_{X,x}$ or simply \mathfrak{m}_x the maximal ideal of $\mathcal{O}_{X,x}$. Moreover, the *residue field of X at x* is the residue field $\kappa(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$.

- (3) A *morphism of locally ringed spaces* $(f, f^\sharp) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a morphism of ringed spaces such that for all $x \in X$ the induced ring map $\mathcal{O}_{Y, f(x)} \rightarrow \mathcal{O}_{X, x}$ is a local ring map.

We know that affine schemes are locally ringed spaces: we take $\text{Spec } R$ with the zariski topology and for any principal open set $D(f)$ we assign the ring R_f . So, any ring R produces the sheaf \widetilde{R} on $\text{Spec } R$. This is called the tilde construction. (sanity check: if you can do this, then you should be able to construct a sheaf on $\text{Spec } R$ for any R -module M).

Definition 2.2. A *scheme* is a locally ringed space with the property that every point has an open neighbourhood which is an affine scheme. A *morphism of schemes* is a morphism of locally ringed spaces. The category of schemes will be denoted Sch .

Definition 2.3. Let (X, \mathcal{O}_X) be a scheme. A sheaf of modules on X is a sheaf \mathcal{F} on X such that for every open set U , $\mathcal{F}(U)$ is an $\mathcal{O}_X(U)$ -module. We say that a sheaf of modules \mathcal{F} is *quasi-coherent* if for every affine open $U \simeq \text{Spec}(R)$, the sheaf $\mathcal{F}|_U$ on U is of the form \widetilde{M} for some R -module M .

Make special note of the next lemma. This basically lets us reduce problems about schemes to statement about affine schemes (ergo, ring theory), whenever the problem at hand is of a *local* nature. Ravi Vakil calls this *affine communication lemma*.

Lemma 2.4. Let X be a scheme. Let P be a local property of rings. The following are equivalent:

- (1) The scheme X is locally P .
- (2) For every affine open $U \subset X$ the property $P(\mathcal{O}_X(U))$ holds.
- (3) There exists an affine open covering $X = \bigcup U_i$ such that each $\mathcal{O}_X(U_i)$ satisfies P .
- (4) There exists an open covering $X = \bigcup X_j$ such that each open subscheme X_j is locally P .

Moreover, if X is locally P then every open subscheme is locally P .

This is how commutative algebra meets geometry. Often, the properties that we want to consider are “globalised” versions of statements about rings. Some examples, normality, reduced, etc.¹

Examples

- (1) Two ways of gluing $\mathbb{A}^1 \setminus \{0\}$: $x \mapsto x$ or $x \mapsto 1/x$.
 - (a) Double origin: What are global sections? what are quasi-coherent sheaves?
 - (b) \mathbb{P}^1 : What are global sections?
- (2) A normal scheme, a reduced scheme, Noetherian scheme.

¹You can also “globalise” morphisms of rings, but now you have two schemes to work locally on. We’ll do this soon.