

The 1-Resolution Property of Algebraic Stacks

Neeraj Deshmukh

July 12, 2020

Definition

An algebraic stack is said to have the *resolution property* if every quasi-coherent sheaf of finite-type is the quotient of a vector bundle.

The Totaro–Gross Theorem relates the resolution property to quotients by group actions.

The Totaro–Gross Theorem relates the resolution property to quotients by group actions.

Theorem ([Totaro, 2004, Gross, 2017])

Let \mathcal{X} be a quasi-compact quasi-separated algebraic stack. Then the following are equivalent:

- *\mathcal{X} has affine stabilizer groups at closed points and satisfies the resolution property.*
- *$\mathcal{X} = [U/GL_n]$ where U is a quasi-affine scheme with an action of GL_n . In particular, \mathcal{X} has affine diagonal.*

Note that the resolution property always holds for (quasi-)projective schemes.

Note that the resolution property always holds for (quasi-)projective schemes.

So, this property becomes interesting only in the non-projective situation where, however, very little is known about it.

Definition

An algebraic stack \mathcal{X} is said to have the *1-resolution property* if it admits a vector bundle V such that every quasi-coherent sheaf of finite-type on \mathcal{X} is a quotient of $V^{\oplus n}$ for some natural number n .

Definition

An algebraic stack \mathcal{X} is said to have the *1-resolution property* if it admits a vector bundle V such that every quasi-coherent sheaf of finite-type on \mathcal{X} is a quotient of $V^{\oplus n}$ for some natural number n .

We will say that such a V is *special*.

The 1-resolution property descends along finite finitely presented covers.

The 1-resolution property descends along finite finitely presented covers.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a faithfully flat finite and finitely presented morphism of algebraic stacks. If X has the 1-resolution property then so does Y .

The 1-resolution property descends along finite finitely presented covers.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a faithfully flat finite and finitely presented morphism of algebraic stacks. If X has the 1-resolution property then so does Y .

And can be pulled back along quasi-affine morphisms.

The 1-resolution property descends along finite finitely presented covers.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a faithfully flat finite and finitely presented morphism of algebraic stacks. If X has the 1-resolution property then so does Y .

And can be pulled back along quasi-affine morphisms.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a quasi-affine morphism of algebraic stacks. If Y has the 1-resolution property then so does X .

The 1-resolution property descends along finite finitely presented covers.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a faithfully flat finite and finitely presented morphism of algebraic stacks. If X has the 1-resolution property then so does Y .

And can be pulled back along quasi-affine morphisms.

Lemma ([Gross, 2017])

Let $X \rightarrow Y$ be a quasi-affine morphism of algebraic stacks. If Y has the 1-resolution property then so does X .

These results also hold for the resolution property.

Hall and Rydh considered this property for algebraic stacks and posed the following question ([Hall and Rydh, 2017, 7.4]):

Question

Does every algebraic stack with the 1-resolution property admit a finite flat covering by a quasi-affine scheme? Moreover, is every algebraic space with the 1-resolution property quasi-affine?

Hall and Rydh considered this property for algebraic stacks and posed the following question ([Hall and Rydh, 2017, 7.4]):

Question

Does every algebraic stack with the 1-resolution property admit a finite flat covering by a quasi-affine scheme? Moreover, is every algebraic space with the 1-resolution property quasi-affine?

In this talk we will try and answer this question under moderate hypotheses. We have the following theorem (joint with Amit Hogadi and Siddharth Mathur):

Theorem ([DHM20])

Let \mathcal{X} be a Noetherian, quasi-excellent and normal algebraic stack whose stabilizers at closed points are affine. Then we have:

- *\mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.*

A result of David Rydh gives us the following corollary.

Corollary

An algebraic space X (Noetherian, quasi-excellent, and normal) has the 1-resolution property if and only if it is quasi-affine.

Some Examples

Some Examples

- For finite group G , BG has the 1-resolution property.

Some Examples

- For finite group G , BG has the 1-resolution property.
- BG_m and BG_a do not have the 1-resolution property.

Some Examples

- For finite group G , BG has the 1-resolution property.
- BG_m and BG_a do not have the 1-resolution property.
- A DVR with a doubled point does not have the 1-resolution property.

Some Examples

- For finite group G , BG has the 1-resolution property.
- BG_m and BG_a do not have the 1-resolution property.
- A DVR with a doubled point does not have the 1-resolution property.
- Any projective variety over a field with the 1-resolution property is a set of points.

We can take the example of the DVR with a doubled point further.

We can take the example of the DVR with a doubled point further. Let X be a scheme with two maps from a DVR $f_1, f_2 : \operatorname{Spec} R \rightarrow X$ which agree on the quotient field $K \supset R$.

We can take the example of the DVR with a doubled point further. Let X be a scheme with two maps from a DVR $f_1, f_2 : \operatorname{Spec} R \rightarrow X$ which agree on the quotient field $K \supset R$.

We can glue the maps f_1 and f_2 along $\operatorname{Spec} K$, and get map f from $\operatorname{Spec} R$ with a doubled point to X .

We can take the example of the DVR with a doubled point further. Let X be a scheme with two maps from a DVR $f_1, f_2 : \operatorname{Spec} R \rightarrow X$ which agree on the quotient field $K \supset R$.

We can glue the maps f_1 and f_2 along $\operatorname{Spec} K$, and get map f from $\operatorname{Spec} R$ with a doubled point to X .

Using the pullback property, we see that if a scheme X has the 1-resolution property, it must be separated.

More generally, we have the following:

More generally, we have the following:

Lemma ([DHM20])

Let \mathcal{X} be a qcqs algebraic stack whose stabilizer groups at closed points are affine. If \mathcal{X} has the 1-resolution property then \mathcal{X} is separated, and hence has finite diagonal.

Strategy of Proof

Strategy of Proof

Theorem ([DHM, 2020])

Let \mathcal{X} be a Noetherian, quasi-excellent and normal algebraic stack whose stabilizers at closed points are affine. Then we have \mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.

Strategy of Proof

Theorem ([DHM, 2020])

Let \mathcal{X} be a Noetherian, quasi-excellent and normal algebraic stack whose stabilizers at closed points are affine. Then we have \mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.

- Prove that if \mathcal{X} is a scheme, it is quasi-affine.

Strategy of Proof

Theorem ([DHM, 2020])

Let \mathcal{X} be a Noetherian, quasi-excellent and normal algebraic stack whose stabilizers at closed points are affine. Then we have \mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.

- Prove that if \mathcal{X} is a scheme, it is quasi-affine.
- Starting from a proper flat covering, use the slicing strategy of Kresch-Vistoli as in [Kresch and Vistoli, 2004] to produce a finite flat scheme covering, and conclude from the scheme case.

The proof in the scheme case is an argument by Noetherian induction which requires openness of the regular locus. This is ensured by the quasi-excellence hypothesis.

The proof in the scheme case is an argument by Noetherian induction which requires openness of the regular locus. This is ensured by the quasi-excellence hypothesis.

The original argument of Kresch and Vistoli works for any Deligne-Mumford stack over a field.

The proof in the scheme case is an argument by Noetherian induction which requires openness of the regular locus. This is ensured by the quasi-excellence hypothesis.

The original argument of Kresch and Vistoli works for any Deligne-Mumford stack over a field.

The following result extends their argument to the arithmetic setting of stacks with finite diagonal over $\mathrm{Spec} Z$.

Theorem ([DHM20])

Let \mathcal{X} be a quasi-compact algebraic stack with finite diagonal and denote by X the corresponding coarse moduli space. Suppose that \mathcal{X} has the resolution property and that X admits an ample line bundle. Then \mathcal{X} admits a finite flat cover $Z \rightarrow \mathcal{X}$ where Z is a scheme with an ample line bundle.

Theorem ([DHM20])

Let \mathcal{X} be a quasi-compact algebraic stack with finite diagonal and denote by X the corresponding coarse moduli space. Suppose that \mathcal{X} has the resolution property and that X admits an ample line bundle. Then \mathcal{X} admits a finite flat cover $Z \rightarrow \mathcal{X}$ where Z is a scheme with an ample line bundle.

Observe that this technique needs the existence of an ample line bundle on the coarse space.

Theorem ([DHM20])

Let \mathcal{X} be a quasi-compact algebraic stack with finite diagonal and denote by X the corresponding coarse moduli space. Suppose that \mathcal{X} has the resolution property and that X admits an ample line bundle. Then \mathcal{X} admits a finite flat cover $Z \rightarrow \mathcal{X}$ where Z is a scheme with an ample line bundle.

Observe that this technique needs the existence of an ample line bundle on the coarse space.

Our hypotheses (Noetherian, normal and quasi-excellent) on \mathcal{X} ensure this.

Approximation

Approximation

We can use limit arguments to eliminate the Noetherian, and quasi-excellence hypotheses on \mathcal{X} .

Approximation

We can use limit arguments to eliminate the Noetherian, and quasi-excellence hypotheses on \mathcal{X} .

This, however, only works in characteristic zero, since it uses the fact that GL_n is linearly reductive in characteristic zero.

Approximation

Approximation

The argument roughly goes as follows:

Approximation

The argument roughly goes as follows:

- \mathcal{X} can be written as a limit $\lim_{\lambda} \mathcal{X}_{\lambda}$ where each \mathcal{X}_{λ} is an algebraic stack of finite type over $\mathrm{Spec} \mathbb{Q}$ and all the maps $\mathcal{X} \rightarrow \mathcal{X}_{\lambda}$ are affine and schematically dominant.

Approximation

The argument roughly goes as follows:

- \mathcal{X} can be written as a limit $\lim_{\lambda} \mathcal{X}_{\lambda}$ where each \mathcal{X}_{λ} is an algebraic stack of finite type over $\mathrm{Spec} \mathbb{Q}$ and all the maps $\mathcal{X} \rightarrow \mathcal{X}_{\lambda}$ are affine and schematically dominant.
- The special vector bundle on \mathcal{X} descend to a special vector bundle at some finite stage \mathcal{X}_{α} for which the result has been proved.

Approximation

The argument roughly goes as follows:

- \mathcal{X} can be written as a limit $\lim_{\lambda} \mathcal{X}_{\lambda}$ where each \mathcal{X}_{λ} is an algebraic stack of finite type over $\mathrm{Spec} \mathbb{Q}$ and all the maps $\mathcal{X} \rightarrow \mathcal{X}_{\lambda}$ are affine and schematically dominant.
- The special vector bundle on \mathcal{X} descend to a special vector bundle at some finite stage \mathcal{X}_{α} for which the result has been proved.
- Base change the finite flat cover to \mathcal{X} , which is quasi-affine because the map $\mathcal{X} \rightarrow \mathcal{X}_{\alpha}$ is affine.

This gives us the following strengthening of our theorem.

This gives us the following strengthening of our theorem.

Theorem ([DHM20])

Let $\mathcal{X}/\mathrm{Spec} \mathbb{Q}$ be a qcqs integral normal algebraic stack whose stabilizers at closed points are affine. Then we have:

- *\mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.*

This gives us the following strengthening of our theorem.

Theorem ([DHM20])

Let $\mathcal{X}/\mathrm{Spec} \mathbb{Q}$ be a qcqs integral normal algebraic stack whose stabilizers at closed points are affine. Then we have:

- *\mathcal{X} has the 1-resolution property if and only if there exists a finite flat covering $U \rightarrow \mathcal{X}$ with U a quasi-affine scheme.*

In particular, if such an \mathcal{X} is an algebraic space then it must already be quasi-affine.

Beyond the normal

Beyond the normal

- As it stands, answering Question 6 in complete generality is beyond the scope of our methods and would require new and innovative techniques.

Beyond the normal

- As it stands, answering Question 6 in complete generality is beyond the scope of our methods and would require new and innovative techniques.
- Another direction is to try and remove the normality hypothesis. It is not clear how to do this even in the scheme case. To that, we ask the following questions. A positive answer to these question will help generalise the above theorems to the non-normal setting.

Question

Let X, Y, Z be quasi-affine schemes which are finite-type over an excellent Noetherian scheme. Suppose we have a closed immersion $Y \rightarrow X$ and a finite morphism $Y \rightarrow Z$, then does the resulting pushout $X \cup_Y Z$ admit an ample family of line bundles?

Question

Let X, Y, Z be quasi-affine schemes which are finite-type over an excellent Noetherian scheme. Suppose we have a closed immersion $Y \rightarrow X$ and a finite morphism $Y \rightarrow Z$, then does the resulting pushout $X \cup_Y Z$ admit an ample family of line bundles?

Question

Let \mathcal{X} be an algebraic stack which is finite-type over an excellent base and suppose \mathcal{X} has the 1-resolution property. Does the coarse moduli space of \mathcal{X} have the 1-resolution property?

References I



Deshmukh, N., Hogadi, A., and Mathur, S. (2020). Quasi-Affineness and the 1-Resolution Property. *International Mathematics Research Notices*. rnaa125.



Gross, P. (2017).
Tensor generators on schemes and stacks.
Algebr. Geom., 4(4):501–522.



Hall, J. and Rydh, D. (2017).
Perfect complexes on algebraic stacks.
Compos. Math., 153(11):2318–2367.

References II



Kresch, A. and Vistoli, A. (2004).

On coverings of Deligne-Mumford stacks and surjectivity of the Brauer map.

Bull. London Math. Soc., 36(2):188–192.



Totaro, B. (2004).

The resolution property for schemes and stacks.

J. Reine Angew. Math., 577:1–22.