

Laplacians as Spatial Operators: From $\nabla \cdot \nabla$ to Two-Layer RelWeights in Spatial Econometrics

Executive summary

The Laplacian is fundamentally an **operator that converts local differences (“gradients”) into net imbalance (“divergence”)**. In continuous space, it is defined by $\Delta = \nabla \cdot \nabla$, and it sits at the core of elliptic and parabolic PDEs—especially **Poisson/Laplace** problems, **heat/diffusion** dynamics, and **eigenvalue/eigenfunction** decompositions that define “spatial frequencies” subject to boundary conditions. Green’s identities formalize how Laplacians interact with boundaries and underpin weak/variational formulations used across analysis and numerical methods. ¹

In discrete settings (lattices, graphs, networks, spatial weights matrices), the Laplacian becomes a matrix operator $L = D - W$ (and normalized variants), with a hallmark quadratic form

$$x^\top Lx = \frac{1}{2} \sum_{i,j} w_{ij} (x_i - x_j)^2,$$

which makes “roughness/smoothness” precise as a weighted sum of squared differences. This identity is the discrete counterpart of the Dirichlet energy $\int |\nabla f|^2$, and it enables clean interpretations in clustering, diffusion, and regularization. ²

In spatial econometrics, the key object is not the Laplacian per se but the **spatial weights matrix** W as a spatial interaction operator (spatial lag Wy , spatially lagged covariates WX , and spatially structured errors). However, the Laplacian emerges immediately when you ask: “What does it mean for a spatial signal to be smooth relative to W ?”—it becomes a **penalty operator**, a **precision matrix** in CAR/GMRF priors, and a **spectral basis** for eigenvector spatial filtering. Two-layer RelWeights constructions—where an overlay/incidence matrix B links observational units α to support/exposure units β —naturally induce a one-mode projection $R = BB^\top$ and thereby an induced Laplacian $L_R = D_R - R$. This provides a principled way to encode **support-mediated spatial externalities** (e.g., watersheds, aquifers, school zones) and connects directly to change-of-support and areal interpolation/misalignment issues. ³

Continuous Laplacian as a differential operator

Definition and local meaning. On a domain $\Omega \subset \mathbb{R}^d$, for sufficiently smooth u ,

$$\Delta u \equiv \nabla \cdot (\nabla u) = \sum_{k=1}^d \frac{\partial^2 u}{\partial x_k^2}.$$

The gradient ∇u is a vector field of local directional change; the divergence $\nabla \cdot v$ measures local net outflow of a vector field. Composing them ($\nabla \cdot \nabla u$) measures, heuristically, how much u differs from its local neighborhood average (capturing curvature/imbalance). ⁴

Green's identities (integration by parts for Δ). For smooth enough u, v on Ω with outward unit normal ν on $\partial\Omega$, Green's first identity:

$$\int_{\Omega} v \Delta u \, dx = - \int_{\Omega} \nabla u \cdot \nabla v \, dx + \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, dS,$$

and Green's second identity:

$$\int_{\Omega} (v \Delta u - u \Delta v) \, dx = \int_{\partial\Omega} \left(v \frac{\partial u}{\partial \nu} - u \frac{\partial v}{\partial \nu} \right) \, dS.$$

These identities are the precise statement that Laplacians are “divergence-of-gradient” operators whose action can be transferred from interior to boundary terms—crucial for uniqueness, solvability conditions, and weak formulations. ⁵

Boundary conditions as part of the operator. The Laplacian is not “complete” until you specify how it interacts with the boundary: - **Dirichlet:** $u|_{\partial\Omega} = g$ (values fixed on the boundary). - **Neumann:** $\frac{\partial u}{\partial \nu}|_{\partial\Omega} = g$ (normal flux fixed). - **Robin/Mixed:** $au + b \frac{\partial u}{\partial \nu} = g$ on $\partial\Omega$ (value-flux combination), and combinations on different boundary segments are common. ⁶

A key structural example: Green's identity implies a **compatibility condition** for pure Neumann problems of Laplace's equation. If $\Delta u = 0$ in Ω and $\partial_{\nu} u = g$ on $\partial\Omega$, then necessarily $\int_{\partial\Omega} g \, dS = 0$. This is one of the canonical ways boundary conditions constrain solvability. ⁵

Canonical PDE roles. - **Heat/diffusion equation:** $u_t = \Delta u$ (or $u_t = \kappa \Delta u$). The Laplacian is the generator of a diffusion semigroup $T(t) = e^{t\Delta}$, meaning solutions can be written (in appropriate function spaces) as $u(t) = e^{t\Delta} u(0)$ and satisfy the semigroup property $e^{(s+t)\Delta} = e^{s\Delta} e^{t\Delta}$. This formalizes the intuition that Laplacian-driven dynamics smooth spatial heterogeneity over time. ⁷ - **Poisson/Laplace equations:** Poisson is typically written $\Delta u = f$ and Laplace is the harmonic case $\Delta u = 0$, posed with boundary conditions (Dirichlet/Neumann/Robin). These are the basic elliptic operators for equilibrium states (steady heat, potentials, etc.). ⁸ - **Eigenproblems (“spatial frequencies”):** find (λ, ϕ) such that

$$-\Delta \phi = \lambda \phi \quad \text{in } \Omega, \quad \phi|_{\partial\Omega} = 0 \text{ (Dirichlet) or } \partial_{\nu} \phi|_{\partial\Omega} = 0 \text{ (Neumann)}.$$

In many standard settings, $\lambda \geq 0$; with full Dirichlet conditions the smallest eigenvalue is strictly positive, while with full Neumann conditions a zero eigenvalue appears (constant eigenfunction), reflecting “no-change” modes. ⁹

Discrete and graph Laplacians as divergence–gradient operators

From spatial weights to a Laplacian matrix. Let $W = [w_{ij}]$ be a symmetric nonnegative weight (adjacency/similarity) matrix on n spatial units (nodes), with $w_{ii} = 0$. Define the degree/strength matrix $D = \text{diag}(d_i)$, $d_i = \sum_j w_{ij}$. The (combinatorial) graph Laplacian is

$$L \equiv D - W.$$

This is the discrete analogue of $-\Delta$ under many finite-difference style interpretations, and it is the foundational Laplacian used in spectral graph theory and graph-based spatial modeling. ¹⁰

Entrywise operator meaning (“divergence of gradient” on a graph). For a node-valued function $x \in \mathbb{R}^n$,

$$(Lx)_i = \sum_{j=1}^n w_{ij}(x_i - x_j).$$

This is exactly the “net outflow” of pairwise differences: each edge (i, j) carries a **discrete gradient** $x_i - x_j$, and the Laplacian sums (“diverges”) these along all edges incident to i . ¹¹

Quadratic form and roughness energy. The graph Laplacian’s quadratic form is the discrete Dirichlet energy:

$$x^\top Lx = \frac{1}{2} \sum_{i,j} w_{ij}(x_i - x_j)^2 \geq 0.$$

This identity makes the “smoothness” interpretation rigorous: small $x^\top Lx$ means $x_i \approx x_j$ whenever w_{ij} is large; large $x^\top Lx$ means many (or heavily weighted) sharp contrasts. ¹²

Incidence matrices and discrete calculus. If you orient the edges and build an oriented incidence matrix Q (vertex–edge), then L can be written as an incidence product. A classical statement is:

$$L = QQ^\top,$$

and this identity is independent of the arbitrary orientation (because it cancels out in $QQ^\top$). This explicitly realizes L as “divergence $\circ$ gradient” under the identification $Q^\top x = \text{edge differences}$. ¹³

Mermaid schematic linking the discrete objects:

```

flowchart LR
    x["node signal x (values on regions)"] --> g["edge gradients: (Q^T x)_e = x_i - x_j"]
    g --> div["node divergence: (Q g)_i = sum of signed edge flows"]
    div --> Lx["Lx = Q(Q^T x) = QQ^T x"]
    Lx --> energy["x^T L x = ||Q^T x||^2 (weighted)"]

```

Normalized Laplacians. Several normalizations matter in practice, especially when degrees vary: - Random-walk normalized Laplacian:

$$L_{\text{rw}} = I - D^{-1}W.$$

- Symmetric normalized Laplacian:

$$L_{\text{sym}} = I - D^{-1/2} W D^{-1/2}.$$

These are central in diffusion and clustering because they normalize out degree heterogeneity (and connect directly to Markov chains/random walks). ¹⁴

Spectrum, diffusion, and clustering

Eigenvalues/eigenvectors as spatial frequencies. For symmetric L , the eigenpairs $Lu_k = \lambda_k u_k$ form an orthonormal basis. Key structural facts: - L is symmetric positive semidefinite, so $\lambda_k \geq 0$. - The smallest eigenvalue is 0 with eigenvector $\mathbf{1}$ (constant mode). - The multiplicity of $\lambda = 0$ equals the number of connected components. ¹²

The constant eigenvector is the discrete analogue of a “flat field.” All other eigenvectors represent departures from constancy with increasing “roughness” as λ_k increases, because $u_k^\top Lu_k = \lambda_k \|u_k\|^2$ and the quadratic form equals a sum of squared differences. ¹²

Fiedler vector and algebraic connectivity. The eigenvector associated with the **second-smallest** eigenvalue (often called the Fiedler vector) plays a special role in partitioning: its signs or quantiles define an approximate minimum-cut/sparsest separation in many settings, and the corresponding eigenvalue measures a kind of connectivity (“algebraic connectivity”). ¹⁵

Cheeger inequality (geometry $\leftrightarrow$ spectrum). Cheeger-type inequalities relate: - a combinatorial/geometric notion of bottleneck (conductance/isoperimetric ratio), - to the second eigenvalue of a (normalized) Laplacian.

A standard formulation (one of several equivalent variants) is that the second eigenvalue of the normalized Laplacian controls—and is controlled by—graph conductance up to squared factors, providing theoretical justification for spectral clustering and fast mixing in diffusion/random walks. ¹⁶

Diffusion and heat kernels on graphs. Graph diffusion mirrors continuous diffusion:

$$\frac{dx(t)}{dt} = -Lx(t) \quad \Rightarrow \quad x(t) = e^{-tL}x(0).$$

Spectrally, if $L = U\Lambda U^\top$, then $e^{-tL} = Ue^{-t\Lambda}U^\top$, i.e., each eigenmode decays at rate λ_k . This is the discrete analogue of $e^{t\Delta}$ in PDE theory and is central to interpretations of smoothing, clustering, and mixing times. ¹⁷

Laplacians and spatial weights in spatial econometrics

Spatial weights W as the core spatial operator. In spatial econometrics, the weights matrix is a **nonstochastic, exogenously specified** representation of the neighborhood structure. Row-standardization is commonly used so that Wy is interpretable as a weighted neighbor average (a “spatial smoother”), though it is not required mathematically and can break symmetry. ¹⁸

Spatial lag, spatial error, and SLX/SDM models as operator statements. The standard taxonomy is operator-first:

Model family	Canonical form	“Where the spatial operator enters”	Typical interpretation
Spatial lag (SAR/SLM)	$y = \rho W y + X\beta + \varepsilon$	outcome	substantive spillovers; simultaneity/endogeneity of $W y$
Spatial error (SEM)	$y = X\beta + u, u = \lambda W u + \varepsilon$	disturbance	nuisance dependence / spatially correlated shocks
SLX	$y = X\beta + W X\theta + \varepsilon$	covariates	contextual spillovers, reduced-form neighbor covariate effects
SDM	$y = \rho W y + X\beta + W X\theta + \varepsilon$	outcome + covariates	rich spillovers; nonlinear marginal effects

These specifications—and interpretive distinctions between “substantive” vs “nuisance” dependence—are standard in the spatial econometrics literature. ¹⁹

Parameter-space restrictions link to spectra. For SAR-type processes, avoiding singularity/explosiveness imposes bounds on ρ in terms of eigenvalues of W , e.g. $\rho \in (1/\omega_{\min}, 1/\omega_{\max})$. For row-standardized weights, $\omega_{\max} = 1$ while $\omega_{\min} > -1$ (so the lower bound is less than -1). This makes the spectrum operational for estimation feasibility and stability. ²⁰

Moran’s I as a quadratic form (and its Laplacian complement). Global Moran’s I can be expressed as a ratio of quadratic forms and can be visualized as the slope of $W x$ regressed on x (in deviations from the mean). ²¹

If W is **row-standardized** and $z = x - \bar{x}\mathbf{1}$, then (with $S_0 = n$) Moran’s I simplifies to

$$I = \frac{z^\top W z}{z^\top z}.$$

Now define the random-walk Laplacian $L_{\text{rw}} = I - W$. Then the normalized “Laplacian energy ratio” satisfies

$$\frac{z^\top L_{\text{rw}} z}{z^\top z} = 1 - I.$$

So Moran’s I and Laplacian energy are dual diagnostics: Moran’s I increases with positive autocorrelation, while Laplacian energy decreases as the field becomes smooth relative to the weight structure. ²²

Eigenvector spatial filtering (ESF/MEM) as Laplacian-weight spectral regression. Eigenvector spatial filtering constructs spatial basis functions from eigenvectors of a transformed weights operator (often involving projection matrices like $M = I - X(X^\top X)^{-1}X^\top$), and includes selected eigenvectors as regressors to absorb spatial dependence in residuals. This links “spatial pattern” directly to the spectrum of matrices derived from W and Moran-type quadratic forms. ²³

CAR/GMRF priors: Laplacian as a precision matrix. In conditional autoregressive (CAR) models and Gaussian Markov random fields (GMRFs) on areal units, the **precision matrix** Q is sparse and encodes adjacency; typical constructions set $Q_{ij} = -1$ for neighbors, and Q_{ii} equals the number of neighbors, forming a Laplacian-like operator (intrinsic CAR corresponds to a rank-deficient Laplacian). This is the Bayesian smoothing analogue of penalizing $x^\top Lx$. ²⁴

A modern bridge between continuous and discrete spatial modeling is the SPDE approach connecting continuously indexed Gaussian fields to GMRFs, where differential operators involving Δ generate covariance structure while sparse precision matrices enable computation. ²⁵

Two-layer RelWeights as induced spatial operators

Derivation: an overlay/incidence matrix induces a one-mode projection

Let α denote the observational units (e.g., municipalities) and β denote a second spatial support/exposure system (e.g., watershed catchments, aquifer polygons, school attendance zones). Define an $|\alpha| \times |\beta|$ matrix B whose entries encode overlap or exposure—common choices include: - binary incidence $B_{im} \in \{0, 1\}$ (does α_i intersect β_m ?), - proportional overlap $B_{im} = \frac{\text{area}(\alpha_i \cap \beta_m)}{\text{area}(\alpha_i)}$, - population-weighted overlap, etc., following standard areal interpolation logic for transferring data between misaligned polygon systems. ²⁶

Then the induced “RelWeights” adjacency on α is a **one-mode projection**:

$$R = BB^\top.$$

This is the standard bipartite-to-one-mode construction: two α -units are linked if they share β -neighbors, with link strength increasing with the amount/number of shared β -support. ²⁷

Finally define the induced Laplacian on α :

$$L_R = D_R - R, \quad (D_R)_{ii} = \sum_j R_{ij}.$$

This turns your two-layer exposure geometry into a **spatial operator on the observational units**.

Change-of-support interpretation

RelWeights is best viewed as a **mechanism-driven weights matrix** arising from a change-of-support / misalignment problem:

- Data and processes often live on different supports (points vs polygons, old census tracts vs new tracts, administrative vs ecological boundaries). “Change of support” formalizes inference when the target support differs from the observed support. ²⁸
- Overlay matrices B (area- or population-weighted) are ubiquitous in areal interpolation: intersect source and target polygons, compute weights, and reaggregate. ²⁹

In this view, $R = BB^\top$ is the induced “neighborhood” structure on α that arises because α -units share the same β -support (shared exposure pathways), rather than because α -units are geographically adjacent.

Examples of support-mediated externalities

RelWeights is especially natural when the externality mechanism is **not adjacency** but **shared membership in a spatial system**:

- **Watersheds / river basins**: upstream/downstream or shared subcatchments induce interaction beyond adjacency.
- **Aquifers**: extraction or contamination spreads through shared hydrogeological units.
- **School zones**: demographic or policy effects propagate through shared attendance/assignment regions.
- **Risk/hazard zones** (floodplains, fire risk, industrial buffers): correlation and spillovers reflect shared exposure supports rather than contiguity.

These examples are instances of the broader areal-interpolation/change-of-support setting in which exposures and outcomes are aggregated differently. ³⁰

Implications for identification and estimation in spatial econometrics

RelWeights interacts with core spatial econometric identification issues in distinctive ways:

Exogeneity of weights. Standard spatial econometric practice treats W as fixed and exogenous; RelWeights inherits this requirement—if B is constructed from administrative/hydrogeological boundaries independent of the outcome model, that aligns with the exogeneity assumption. If B depends on endogenous behavior (e.g., zones drawn in response to outcomes), the “weights are exogenous” premise becomes questionable. ³¹

Collinearity and reduced-form similarity. Operators like WX can become highly collinear with X (and Wy can mimic unobserved heterogeneity) depending on how B aggregates information. This mirrors known sensitivity of estimation and testing to weights specifications and motivates explicit diagnostics and robustness checks over alternative B definitions. ³²

Spectrum changes the feasible parameter range. Because parameter bounds depend on the eigenvalues of the lag operator, moving from adjacency-based W to exposure-based R changes the spectral radius and thus the feasible region for ρ (and the interpretation of spatial multipliers). This is not a minor technicality: it changes stability, identification strength, and the implied “range” of global spillovers. ³³

Computational recipes, diagnostics, and toy examples

Tables comparing operators, models, and interpretations

Continuous vs discrete Laplacians.

Concept	Continuous operator	Discrete/graph analogue	What it measures
Gradient	∇u	edge differences $x_i - x_j$ (via incidence $Q^\top x$)	local change
Divergence	$\nabla \cdot v$	nodewise net outflow (via $Q(\cdot)$)	imbalance/flux
Laplacian	$\Delta u = \nabla \cdot \nabla u$	$Lx = (D - W)x$	net deviation from neighbors
Energy	$\int_{\Omega} \nabla u ^2 dx$	$x^\top Lx = \frac{1}{2} \sum w_{ij} (x_i - x_j)^2$	roughness penalty

These correspondences are standard in discrete calculus, spectral graph theory, and Laplacian-based learning. ³⁴

Spatial econometrics models as operator placements.

Operator object	Where it shows up	Typical role	Related Laplacian view
Wy	SAR/SLM	endogenous interaction (spillovers)	“smoothing” of outcome; $I - W$ as roughness
Wu	SEM	nuisance dependence	precision-like structure if re-expressed
WX	SLX/SDM	contextual spillovers	smoothing of covariate fields
$L = D - W$	CAR/GMRF, regularization	prior precision / penalty	explicit smoothness enforcement
eigenvectors of W -derived matrices	ESF/MEM	spatial pattern basis	“spatial harmonics”

35

Practical construction of W , B , R , and Laplacians

Constructing adjacency-based W (single-layer). Common workflows define neighbors by contiguity, distance thresholds, or k-nearest neighbors; GIS tools explicitly implement these constructions. ³⁶

Constructing overlay/exposure B (two-layer). The standard areal interpolation workflow is “intersect $\rightarrow$ compute overlap weights $\rightarrow$ reaggregate,” producing a matrix of weights between source and target polygon systems. ²⁹

Forming $R = BB^\top$. This is the induced one-mode projection; it is symmetric positive semidefinite by construction and encodes shared-support similarity. ³⁷

Row-standardization and normalization choices. Row-standardization of weights is common in spatial econometrics to interpret Wy as a neighbor average; normalized Laplacians remove degree scaling and are often preferred in spectral methods when degrees vary. ³⁸

Eigenpairs and numerical computation

When n is small/moderate, dense eigendecomposition is fine. **When n is large**, Laplacians are sparse and you typically compute only a few extremal eigenpairs (e.g., smallest nonzero eigenvalues for clustering / smoothing). ARPACK is a classic library for “a few eigenpairs of large sparse matrices,” and SciPy documents the same matrix-free sparse-eigsolver approach. ³⁹

Numerical stability notes specific to Laplacians. - Graph Laplacians are singular whenever the graph has at least one connected component (there is always a constant nullspace mode on each component), so inverses are replaced by Moore–Penrose pseudoinverses or constrained solves (e.g., sum-to-zero constraints in ICAR). ⁴⁰ - In Bayesian ICAR/BYM2 workflows, explicit scaling uses the pseudoinverse diagonal of the Laplacian to standardize marginal variances—illustrating how singularity is handled in practice. ⁴¹

Toy example: your 4×4 RelWeights Laplacian

You gave

$$L_R = \begin{bmatrix} 3 & -2 & -1 & 0 \\ -2 & 5 & -2 & -1 \\ -1 & -2 & 5 & -2 \\ 0 & -1 & -2 & 3 \end{bmatrix}.$$

Recover the induced relational adjacency R and degrees D_R . Since $L_R = D_R - R$ and D_R is the diagonal of L_R ,

$$D_R = \text{diag}(3, 5, 5, 3), \quad R = \begin{bmatrix} 0 & 2 & 1 & 0 \\ 2 & 0 & 2 & 1 \\ 1 & 2 & 0 & 2 \\ 0 & 1 & 2 & 0 \end{bmatrix}.$$

Interpreting R_{ij} : it is the strength of relational coupling between nodes i and j .

Mermaid sketch of the induced weighted graph:

```
graph LR
  n1((1)) --"2"--> n2((2))
  n1 --"1"--> n3((3))
  n2 --"2"--> n3
  n2 --"1"--> n4((4))
  n3 --"2"--> n4
```

Compute $L_R x$ as divergence-of-gradient. For any $x = [x_1, x_2, x_3, x_4]^\top$,

$$(L_R x)_i = \sum_j R_{ij}(x_i - x_j).$$

This is the nodewise “net outflow” of edge differences. ¹¹

Example signal (chosen to create a left-right contrast):

$$x = \begin{bmatrix} 1 \\ 2 \\ 5 \\ 6 \end{bmatrix}.$$

Then (computed directly),

$$L_R x = \begin{bmatrix} -6 \\ -8 \\ 8 \\ 6 \end{bmatrix}.$$

Interpretation: nodes 1–2 are far **below** their relational neighborhood average (negative divergence), while nodes 3–4 are far **above** (positive divergence).

Quadratic energy (“roughness”). For the same x ,

$$x^\top L_R x = 54.$$

Because $x^\top L_R x$ equals $\frac{1}{2} \sum_{i,j} R_{ij}(x_i - x_j)^2$, the large value comes from large differences across heavily weighted ties (notably between $\{1, 2\}$ and $\{3, 4\}$). ¹²

Eigenpairs (computed from the matrix). The eigenvalues are:

$$\lambda \in \{0, 5 - \sqrt{5}, 6, 5 + \sqrt{5}\} \approx \{0, 2.7639, 6, 7.2361\}.$$

A convenient eigenbasis is:

Eigenvalue λ	Exact eigenvector v (one choice)	Spatial interpretation
0	$[1, 1, 1, 1]^\top$	perfectly smooth / constant mode
$5 - \sqrt{5}$	$[-1, 2 - \sqrt{5}, -2 + \sqrt{5}, 1]^\top$	smoothest nonconstant “gradient” (left→right)
6	$[1, -1, -1, 1]^\top$	outer-vs-inner contrast
$5 + \sqrt{5}$	$[-1, 2 + \sqrt{5}, -(2 + \sqrt{5}), 1]^\top$	highest-frequency oscillation in the induced geometry

This is the concrete spectral picture behind “Laplacians encode smoothness and clustering”: low- λ eigenvectors are smooth relative to R ; high- λ eigenvectors oscillate sharply and are penalized strongly in $x^\top L_R x$. ⁴²

Moran’s I vs Laplacian energy in this toy case. Using the row-standardized operator $W_{\text{rw}} = D_R^{-1}R$ and the centered vector $z = x - \bar{x}\mathbf{1}$, Moran’s I becomes $I = (z^\top W_{\text{rw}} z) / (z^\top z)$ (since $S_0 = n$ for row-standardized weights). For $x = [1, 2, 5, 6]^\top$, this yields

$$I \approx 0.1294.$$

Meanwhile $z^\top L_R z = 54$ (constant shifts do not change Laplacian energy). This illustrates a practical diagnostic pairing: Moran’s I is modestly positive (some relational clustering), but Laplacian energy is large because there are several strong relational ties across which the signal jumps sharply. ²²

Pseudocode recipes (language-agnostic)

Construct adjacency-based W (single-layer).

```
Input: polygons or points with coordinates
Choose neighborhood rule: contiguity OR distance band OR kNN
For each pair (i,j):
    set w_ij = 1 if neighbors else 0    (or distance-decay weight)
Optionally: row-standardize
Return W
```

(Implemented in common GIS and spatial-statistics toolchains.) ⁴³

Construct overlay matrix B and induced RelWeights R .

```
Input: source polygons alpha (observations), support polygons beta (exposure/
support)
Compute intersections alpha_i n beta_m
Set B[i,m] = area(alpha_i n beta_m) / area(alpha_i)    (or other exposure weight)
Compute R = B * B^T
Set D_R[i,i] = sum_j R[i,j]
Compute L_R = D_R - R
Return R, L_R
```

This is the matrix form of areal-interpolation weighting followed by bipartite one-mode projection. ⁴⁴

Compute a few eigenpairs for clustering / filtering (large n).

```
Input: sparse symmetric L (or L_sym), target k eigenpairs
Use sparse eigensolver (Lanczos/IRLM) to compute smallest nonzero eigenvalues:
```

```

compute (lambda_2,...,lambda_{k+1}) and vectors (u_2,...,u_{k+1})
Use u_2 (Fiedler) for 2-way cut; use span(u_2,...,u_{k+1}) for k-way clustering
Return eigenpairs

```

Large-scale sparse eigencomputations are a standard ARPACK use-case; SciPy and similar systems document this interface. ⁴⁵

Prioritized sources (core references aligned to your request)

The following sources are particularly central for rigorous, operator-focused work that connects PDE, graphs, and spatial modeling:

Area	High-priority references	Why they matter
Graph Laplacians & spectra	Spectral Graph Theory ⁴⁶ ; von Luxburg tutorial	definitions, quadratic forms, normalized Laplacians, spectral clustering foundations ⁴⁷
Laplacian-based learning & regularization	Belkin & Niyogi (Laplacian eigenmaps); manifold regularization	explicit Laplacian $\leftrightarrow$ heat/diffusion $\leftrightarrow$ geometry connection; Laplacian penalties in statistical learning ⁴⁸
Discrete calculus viewpoint	Discrete Calculus: Applied Analysis on Graphs for Computational Science ⁴⁹ ; discrete exterior calculus notes	makes “divergence of gradient” on graphs explicit and operational ⁵⁰
Spatial econometrics & weights	Spatial Econometrics: Methods and Models ⁵¹ ; Anselin overview; LeSage overview	clarifies the operator role of W , row-standardization, SAR/SEM/SDM/SLX modeling, parameter bounds tied to eigenvalues ⁵²
Spatial autocorrelation & filtering	Anselin (LISA); ESF/MEM materials	Moran’s I as quadratic form; eigenvectors as map-pattern basis functions ⁵³
CAR/GMRF priors & smoothing	Julian Besag ⁵⁴ ; Finn Lindgren et al.; Paciorek vignette ⁵⁵	Laplacian-like precision matrices; links between Δ and sparse GMRF computation ⁵⁶
Change of support & areal interpolation	Alan E. Gelfand ⁵⁷ et al.; areal interpolation references	formal framing for B (overlap weights) as support transformation ⁵⁸

Timeline-style orientation of “where Laplacians entered” (very high-level):

```

timeline
  title Laplacian as an operator across fields
  1800s : Laplacian formalized in mathematical physics; potentials and PDEs

```

1900s : Boundary value problems, Green's identities, spectral theory for PDE eigenproblems
 1970s : Graph Laplacians, algebraic connectivity, early spectral partitioning ideas
 1980s : Spatial econometrics formalizes W-based operators (spatial lag/error)
 1990s : Laplacian spectra tied to cuts, conductance; diffusion/random-walk interpretations
 2000s : Graph Laplacians in ML (spectral clustering, Laplacian eigenmaps, regularization)
 2010s : Laplacian/precision matrices in Bayesian spatial models; SPDE-GMRF links

1 4 5 34 46 <https://users.math.msu.edu/users/yanb/849-full-note.pdf>

<https://users.math.msu.edu/users/yanb/849-full-note.pdf>

2 11 12 15 42 47 49 51 57 https://graphics.stanford.edu/courses/cs233-25-spring/ReferencedPapers/Luxburg07_spectral_clustering_tutorial_4488.pdf

https://graphics.stanford.edu/courses/cs233-25-spring/ReferencedPapers/Luxburg07_spectral_clustering_tutorial_4488.pdf

3 18 19 20 31 33 35 38 52 https://web.pdx.edu/~crkl/WISE/SEAUG/papers/anselin01_CTE14.pdf

https://web.pdx.edu/~crkl/WISE/SEAUG/papers/anselin01_CTE14.pdf

6 https://www.math.ucdavis.edu/~hunter/pdes/pde_notes.pdf

https://www.math.ucdavis.edu/~hunter/pdes/pde_notes.pdf

7 17 https://www.math.ucdavis.edu/~hunter/m218a_09/ch5

https://www.math.ucdavis.edu/~hunter/m218a_09/ch5

8 https://math.nyu.edu/faculty/tabak/PDEs/Laplace_and_Poisson_equations.pdf

https://math.nyu.edu/faculty/tabak/PDEs/Laplace_and_Poisson_equations.pdf

9 <https://math.umd.edu/~petersd/462/sepvar.pdf>

<https://math.umd.edu/~petersd/462/sepvar.pdf>

10 13 <https://www.math.ucdavis.edu/~saito/data/graphlap/merris-graphlap-eigvals.pdf>

<https://www.math.ucdavis.edu/~saito/data/graphlap/merris-graphlap-eigvals.pdf>

14 https://fanchung.ucsd.edu/teach/262/notes/paul/10_3_notes.pdf

https://fanchung.ucsd.edu/teach/262/notes/paul/10_3_notes.pdf

16 <https://www.cs.yale.edu/homes/spielman/561/lect11-18.pdf>

<https://www.cs.yale.edu/homes/spielman/561/lect11-18.pdf>

21 22 53 https://dces.webhosting.cals.wisc.edu/wp-content/uploads/sites/128/2013/08/W4_Anselin1995.pdf

https://dces.webhosting.cals.wisc.edu/wp-content/uploads/sites/128/2013/08/W4_Anselin1995.pdf

23 <https://r-spatial.github.io/spatialreg/articles/SpatialFiltering.html>

<https://r-spatial.github.io/spatialreg/articles/SpatialFiltering.html>

24 55 <https://www.stat.berkeley.edu/~paciorek/research/techVignettes/techVignette5.pdf>

<https://www.stat.berkeley.edu/~paciorek/research/techVignettes/techVignette5.pdf>

- 25 https://joeid.folk.ntnu.no/StatGr/Lindgren_etal.pdf
https://joeid.folk.ntnu.no/StatGr/Lindgren_etal.pdf
- 26 <https://walker-data.com/census-r/spatial-analysis-with-us-census-data.html>
<https://walker-data.com/census-r/spatial-analysis-with-us-census-data.html>
- 27 37 <https://web.cs.ucdavis.edu/~raissa/Classes/253-S25/lecture12-2025.pdf>
<https://web.cs.ucdavis.edu/~raissa/Classes/253-S25/lecture12-2025.pdf>
- 28 54 58 <https://academic.oup.com/biostatistics/article-pdf/2/1/31/654877/020031.pdf>
<https://academic.oup.com/biostatistics/article-pdf/2/1/31/654877/020031.pdf>
- 29 44 <https://chris-prener.github.io/areal/articles/areal-weighted-interpolation.html>
<https://chris-prener.github.io/areal/articles/areal-weighted-interpolation.html>
- 30 <https://pro.arcgis.com/en/pro-app/latest/help/analysis/geostatistical-analyst/what-is-areal-interpolation.htm>
<https://pro.arcgis.com/en/pro-app/latest/help/analysis/geostatistical-analyst/what-is-areal-interpolation.htm>
- 32 <https://www.sciencedirect.com/science/article/pii/S1056819023014343>
<https://www.sciencedirect.com/science/article/pii/S1056819023014343>
- 36 43 <https://pro.arcgis.com/en/pro-app/latest/tool-reference/spatial-statistics/generate-spatial-weights-matrix.htm>
<https://pro.arcgis.com/en/pro-app/latest/tool-reference/spatial-statistics/generate-spatial-weights-matrix.htm>
- 39 45 <https://www.arpack.org/>
<https://www.arpack.org/>
- 40 41 <https://mrp-interface.github.io/shinymrp/articles/spatial-prior.html>
<https://mrp-interface.github.io/shinymrp/articles/spatial-prior.html>
- 48 <https://www2.imm.dtu.dk/projects/manifold/Papers/Laplacian.pdf>
<https://www2.imm.dtu.dk/projects/manifold/Papers/Laplacian.pdf>
- 50 <https://download.e-bookshelf.de/download/0000/0082/11/L-G-0000008211-0002342081.pdf>
<https://download.e-bookshelf.de/download/0000/0082/11/L-G-0000008211-0002342081.pdf>
- 56 <https://stat.duke.edu/~scs/Courses/Stat376/Papers/GibbsFieldEst/BesagJRSSB1974.pdf>
<https://stat.duke.edu/~scs/Courses/Stat376/Papers/GibbsFieldEst/BesagJRSSB1974.pdf>