

Proof of a conjecture of Lin and Ma on 0012-avoiding inversion sequences

Shane Chern

Abstract. The study of pattern avoidance in inversion sequences recently attracts extensive research interests. In particular, Zhicong Lin and Jun Ma conjectured a formula that counts the number of inversion sequences avoiding the length-four pattern 0012. Our object of the paper is to confirm this conjecture.

Keywords. Inversion sequence, pattern avoidance, generating function, kernel method.

2010MSC. 05A05, 05A15.

1. Introduction

An *inversion sequence* of length n is a sequence $e = e_1e_2 \cdots e_n$ such that $0 \leq e_i \leq i - 1$ for each $1 \leq i \leq n$. We denote by \mathbf{I}_n the set of inversion sequences of length n . Given any word $w \in \{0, 1, \dots, n-1\}^n$ of length n , we define its *reduction* by the word obtained via replacing the k -th smallest entries of e with $k - 1$. For instance, the reduction of 0023252 is 0012131. We say that an inversion sequence e *contains* a given pattern p if there exists a subsequence of e such that its reduction is the same as p ; otherwise, we say that e *avoids* the pattern p . For instance, 0023252 has a subsequence 022 whose reduction is 011 — hence, 0023252 contains the pattern 011. On the other hand, none of the length 3 subsequences of 0023252 have the reduction 110 — hence, 0023252 avoids the pattern 110.

Let p_1, p_2, \dots, p_m be given patterns. We denote by $\mathbf{I}_n(p_1, p_2, \dots, p_m)$ the set of inversion sequences of length n that avoid all of the patterns p_1, p_2, \dots, p_m . Recently, the study of pattern avoidance in inversion sequences attracts extensive research interests. See [1–8, 10–12, 14–16, 19, 20] for several instances of work on this topic. Among these work, one particular interesting problem is about the enumeration of inversion sequences that avoids fixed patterns. For example, in a pioneering work of Corteel, Martinez, Savage and Weselcouch [7], it was shown that

$$|\mathbf{I}_n(011)| = B_n \quad \text{and} \quad |\mathbf{I}_n(021)| = S_n$$

where B_n is the n -th Bell number (OEIS [18], A000110) and S_n is the n -th large Schröder number (OEIS, A006318).

In a recent paper [19], Yan and Lin proved a conjecture due to Martinez and Savage [16] that claims

$$|\mathbf{I}_n(021, 120)| = 1 + \sum_{i=1}^{n-1} \binom{2i}{i-1}. \quad (1.1)$$

This sequence is registered as A279561 in OEIS. Lin and Yan also showed that this sequence as well enumerates $|\mathbf{I}_n(102, 110)|$ and $|\mathbf{I}_n(102, 120)|$. This therefore

establishes the Wilf-equivalences

$$\mathbf{I}_n(021, 120) \sim \mathbf{I}_n(102, 110) \sim \mathbf{I}_n(102, 120). \quad (1.2)$$

At the end of [19], a conjecture of Zhicong Lin and Jun Ma discovered in 2019 is recorded.

Conjecture 1.1 (Lin and Ma). For $n \geq 1$,

$$|\mathbf{I}_n(0012)| = 1 + \sum_{i=1}^{n-1} \binom{2i}{i-1}. \quad (1.3)$$

In other words, one may extend the balanced Wilf-equivalences (1.2) as the following unbalanced ones:

$$\mathbf{I}_n(0012) \sim \mathbf{I}_n(021, 120) \sim \mathbf{I}_n(102, 110) \sim \mathbf{I}_n(102, 120).$$

It is also worth pointing out that the consideration of length-four pattern avoidance in inversion sequences appears novel in the literature.

Our object of this paper is to confirm the above conjecture of Lin and Ma.

Theorem 1.1. *Conjecture 1.1 is true.*

Let us fix some notation. Given $e = e_1 e_2 \cdots e_n \in \mathbf{I}_n(0012)$, we define

$$\mathcal{R}(e) := \{m : \exists i \neq j \text{ such that } e_i = e_j = m\}.$$

In other words, $\mathcal{R}(e)$ is the set of letters that appear more than once in e . We further define

$$\text{srpt}(e) := \min \mathcal{R}(e),$$

that is, the smallest number in $\mathcal{R}(e)$ — here srpt stands for “smallest repeated.” Notice that there is only one sequence $01 \cdots (n-1)$ in which none of the letters repeat. For this sequence, we assign that

$$\text{srpt}(01 \cdots (n-1)) := n-1.$$

Finally, we define

$$\text{last}(e) := e_n,$$

the last entry of e .

2. Combinatorial observations

We collect some combinatorial observations about inversion sequences in $\mathbf{I}_n(0012)$.

Lemma 2.1. *For $n \geq 1$ and $e \in \mathbf{I}_n(0012)$, if $\text{srpt}(e) = k$, then for $1 \leq i \leq k+1$, we have $e_i = i-1$.*

Proof. If $\text{srpt}(e) = n-1$, then $e = 01 \cdots (n-1)$ and hence the lemma is true. Let $\text{srpt}(e) \neq n-1$. If in this case the lemma is not true, then since $0 \leq e_i \leq i-1$ for each i , there must exist some $k_1 < k = \text{srpt}(e)$ that appears more than once among e_1, e_2, \dots, e_{k+1} . This violates the assumption that $\text{srpt}(e) = k$. \square

Lemma 2.2. *For $n \geq 2$ and $e = e_1 e_2 \cdots e_n \in \mathbf{I}_n(0012)$, let $\gamma(e) = e_1 e_2 \cdots e_{n-1}$. We further assume that $e \neq 01 \cdots (n-1)$. Then*

(a). *if $\text{last}(e) > \text{srpt}(\gamma(e))$, then*

$$\text{srpt}(e) = \text{srpt}(\gamma(e));$$

(b). if $\text{last}(e) \leq \text{srpt}(\gamma(e))$, then

$$\text{srpt}(e) = \text{last}(e).$$

Proof. A simple observation is that $\gamma(e) \in \mathbf{I}_{n-1}(0012)$. Below let us assume that $\text{last}(e) = \ell$, $\text{srpt}(e) = k$ and $\text{srpt}(\gamma(e)) = k'$.

First, if $\mathcal{R}(\gamma(e)) = \emptyset$, then for each $0 \leq i \leq n-1$, $e_i = i-1$. Since $e \neq 01 \cdots (n-1)$, we have $\text{last}(e) = \ell \leq n-2 = \text{srpt}(\gamma(e))$. This fits into Case (b). Further, we find that $\mathcal{R}(e) = \{\ell\}$ and hence $\text{srpt}(e) = \ell$. This implies that $\text{srpt}(e) = \text{last}(e)$.

Now we assume that $\mathcal{R}(\gamma(e)) \neq \emptyset$. Notice that Case (a) is trivial. For Case (b), we first deduce from $\mathcal{R}(\gamma(e)) \neq \emptyset$ that $k' \leq n-3$. By Lemma 2.1, we find that for $1 \leq i \leq k'+1$, $e_i = i-1$. If $\text{last}(e) = \ell \leq k'$, then we know that $e_{\ell+1} = \ell = e_n$. Also, we notice that the indices satisfy $\ell+1 \leq k'+1 \leq n-2 < n$. Hence, $\ell \in \mathcal{R}(e)$. Therefore, $\text{srpt}(e) = \min\{\ell, k'\} = \ell = \text{last}(e)$. \square

Corollary 2.3. For $e \in \mathbf{I}_n(0012)$,

$$0 \leq \text{srpt}(e) \leq \text{last}(e) \leq n-1.$$

Proof. If $e = 01 \cdots (n-1)$, the above inequalities are trivial since $\text{srpt}(e) = \text{last}(e) = n-1$. If $e \neq 01 \cdots (n-1)$, the inequalities are direct consequences of Lemma 2.2 and the fact that $\text{srpt}(e) \geq 0$ and $\text{last}(e) \leq n-1$. \square

Lemma 2.4. For $n \geq 2$ and $e = e_1 e_2 \cdots e_n \in \mathbf{I}_n(0012)$, let e be such that $\text{srpt}(e) = \text{last}(e) = k$ with $0 \leq k \leq n-2$. Then

(a). for $1 \leq i \leq k+1$,

$$e_i = i-1;$$

(b). if we denote $e' = e'_1 e'_2 \cdots e'_{n-k}$ by the sequence obtained via $e'_i = e_{k+i} - k$ for each $1 \leq i \leq n-k$, then $e' \in \mathbf{I}_{n-k}(0012)$ such that

$$\text{srpt}(e') = \text{last}(e') = 0.$$

Proof. Part (a) simply comes from Lemma 2.1. Also, we know from Part (a) that for $k+1 \leq i \leq n$, it holds that $e_i \geq k$. On the other hand, $e_i \leq i-1$. Hence, e' is still an inversion sequence. Further, it is trivial to see that e' still avoids the pattern 0012. Finally, we have $e'_1 = e_{k+1} - k = k - k = 0$ and $\text{last}(e') = e'_{n-k} = e_n - k = k - k = 0$. Since $n-k \geq 2 > 1$, we have $0 \in \mathcal{R}(e')$ and hence $\text{srpt}(e') = 0$. \square

3. Recurrences

Let

$$f_n(k, \ell) := \left\{ \begin{array}{l} \text{the number of sequences } e \in \mathbf{I}_n(0012) \text{ with} \\ \text{srpt}(e) = k \text{ and } \text{last}(e) = \ell \end{array} \right\}.$$

We will establish the following recurrences.

Lemma 3.1. We have

(a). for $n \geq 1$,

$$f_n(n-1, n-1) = 1;$$

(b). for $n \geq 2$,

$$f_n(n-2, n-1) = 0;$$

(c). for $n \geq 2$ and $0 \leq k \leq n-3$,

$$f_n(k, n-1) = \sum_{k'=k}^{n-2} f_{n-1}(k', n-2);$$

(d). for $n \geq 2$ and $0 \leq \ell \leq n-2$,

$$f_n(\ell, \ell) = \sum_{\ell'=\ell}^{n-2} \sum_{k'=\ell}^{\ell'} f_{n-1}(k', \ell');$$

(e). for $n \geq 2$ and $0 \leq k < \ell \leq n-2$,

$$f_n(k, \ell) = \sum_{k'=k}^{\ell} f_{n-1}(k', \ell) + \sum_{\ell'=\ell}^{n-2} f_{n-1}(k, \ell').$$

Proof. Cases (a) and (b) are trivial. In particular, Case (a) enumerates the only inversion sequence $01 \cdots (n-1)$ in which none of the letters repeat. Below we always assume that $e = e_1 e_2 \cdots e_n \in \mathbf{I}_n(0012)$. Let $\gamma(e)$ be as in Lemma 2.2.

For Case (c), let e be such that $\text{srpt}(e) = k \leq n-3$ and $\text{last}(e) = n-1$. We first notice that $e_{n-1} = \text{last}(\gamma(e)) \geq \text{srpt}(\gamma(e))$ by Corollary 2.3. Also, it is easy to see that $\text{srpt}(\gamma(e)) = \text{srpt}(e) = k$ since $\text{last}(e) = n-1 > k$. Now we claim that $e_{n-1} = k$. Otherwise, namely, if $e_{n-1} > k$, we may find $i < j < n-1$ such that $e_i = e_j = k$. Hence, $e_i e_j e_{n-1} e_n$ has the reduction 0012, which contradicts the assumption that $e \in \mathbf{I}_n(0012)$. We therefore have a bijection

$$e = e_1 e_2 \cdots e_{n-2}(k)(n-1) \longleftrightarrow e_1 e_2 \cdots e_{n-2}(n-2) = e'.$$

Notice that e' is still an inversion sequence avoiding the pattern 0012. Also, $\text{srpt}(e') \geq k$. Otherwise, there exists some $k' < k$ that appears more than once among e_1, e_2, \dots, e_{n-2} and therefore $\text{srpt}(e) < k$, which leads to a contradiction. Finally, to prove Case (c), it suffices to show that e' could be any inversion sequence in $\mathbf{I}_{n-1}(0012)$ with $\text{last}(e') = n-2$ (which is of course true) and $\text{srpt}(e') \geq k$. Let e' be such a sequence and assume that $\text{srpt}(e') = k' \geq k$. By Lemma 2.1, we have $e_{k+1} = k$. Pulling back to e , we have $e_{k+1} = e_{n-1} = k$ with the indices $k+1 \leq n-2 < n-1$. Therefore, for this e , we have $k \in \mathcal{R}(e)$ and hence $\text{srpt}(e) = \min\{k', k\} = k$.

For Case (d), let e be such that $\text{srpt}(e) = \text{last}(e) = \ell$ with $0 \leq \ell \leq n-2$. We first find that $\text{srpt}(\gamma(e)) \geq \text{srpt}(e) = \ell$. On the other hand, let $e' = e'_1 e'_2 \cdots e'_{n-1} \in \mathbf{I}_{n-1}(0012)$ be such that $\text{srpt}(e') \geq \ell$. By Lemma 2.1, $e'_{\ell+1} = \ell$. Hence, by appending ℓ to the end of e' , we obtain a sequence with both srpt and last equal to ℓ . We therefore arrive at a bijection between e and e' ,

$$e = e_1 e_2 \cdots e_{n-1}(\ell) \longleftrightarrow e_1 e_2 \cdots e_{n-1} = e',$$

and the desired relation follows.

For Case (e), let e be such that $\text{srpt}(e) = k$ and $\text{last}(e) = \ell$ with $0 \leq k < \ell \leq n-2$. Notice that $e_{n-1} \geq k$. Otherwise, we assume that $e_{n-1} = k' < k$. Then by Lemma 2.1, $e_{k'+1} = k' = e_{n-1}$. However, $k'+1 < k+1 < n-1$ and hence $k' \in \mathcal{R}(e)$. But this violates the fact that $k = \min \mathcal{R}(e)$. Now we have two cases.

- $e_{n-1} < e_n$. We claim that $e_{n-1} = k$. Otherwise, we may find $i < j < n-1$ such that $e_i = e_j = k$. Hence, $e_i e_j e_{n-1} e_n$ has the reduction 0012, which violates

the assumption that $e \in \mathbf{I}_n(0012)$. Now we have a bijection between e and $e' \in \mathbf{I}_{n-1}(0012)$ such that $\text{srpt}(e') \geq k$ and $\text{last}(e') = \ell$ by

$$e = e_1 e_2 \cdots e_{n-2}(k)(\ell) \longleftrightarrow e_1 e_2 \cdots e_{n-2}(\ell) = e'.$$

The argument is similar to that for Case (c). This bijection leads to the first term in the right-hand side of the recurrence relation in Case (e).

- $e_{n-1} \geq e_n$. We have a bijection between e and $e' \in \mathbf{I}_{n-1}(0012)$ such that $\text{srpt}(e') = k$ and $\text{last}(e') \geq \ell$ by

$$e = e_1 e_2 \cdots e_{n-1}(\ell) \longleftrightarrow e_1 e_2 \cdots e_{n-1} = e'.$$

The argument is similar to that for Case (d). This bijection leads to the second term in the right-hand side of the recurrence relation in Case (e).

The proof of the lemma is therefore complete. \square

We may therefore determine the support of $f_n(k, \ell)$.

Corollary 3.2. *For $n \geq 1$, $f_n(k, \ell)$ is supported on*

$$\{(k, \ell) \in \mathbb{N}^2 : 0 \leq k \leq \ell \leq n-1\} \setminus \{(n-2, n-1)\}.$$

Proof. By Corollary 2.3, $f_n(k, \ell) = 0$ if

$$(k, \ell) \notin \{(k, \ell) \in \mathbb{N}^2 : 0 \leq k \leq \ell \leq n-1\}.$$

Also, $f_n(n-2, n-1) = 0$ by Lemma 3.1(b). Finally, for the remaining (k, ℓ) , we have $f_n(k, \ell) \neq 0$ with the help of the recurrences in Lemma 3.1. \square

Finally, we have another recurrence.

Lemma 3.3. *We have, for $n \geq 2$ and $0 \leq k \leq n-2$,*

$$f_n(k, k) = f_{n-k}(0, 0).$$

Proof. This is an immediate consequence of Lemma 2.4. \square

In the sequel, we require three auxiliary functions with q within a sufficiently small neighborhood of 0:

$$\begin{aligned} \mathcal{L}(x; q) &= \sum_{n \geq 1} L_n(x) q^n := \sum_{n \geq 1} \left(\sum_{k=0}^{n-1} f_n(k, n-1) x^k \right) q^n, \\ \mathcal{D}(x; q) &= \sum_{n \geq 1} D_n(x) q^n := \sum_{n \geq 1} \left(\sum_{\ell=0}^{n-2} f_n(\ell, \ell) x^\ell \right) q^n, \\ \mathcal{F}(x, y; q) &= \sum_{n \geq 1} F_n(x, y) q^n := \sum_{n \geq 1} \left(\sum_{\ell=0}^{n-1} \sum_{k=0}^{\ell} f_n(k, \ell) x^k y^\ell \right) q^n. \end{aligned}$$

Notice that $L_1(x) = 1$, $D_1(x) = 0$ and $F_1(x, y) = 1$. Also, since $f_n(n-1, n-1) = 1$, we have

$$\sum_{\ell=0}^{n-1} f_n(\ell, \ell) x^\ell = D_n(x) + x^{n-1}.$$

4. 001-Avoidance and a result of Corteel et al.

The following result on 001-avoidance was shown by Corteel et al. [7].

Theorem 4.1 (Corteel et al.). *For $n \geq 1$,*

$$|\mathbf{I}_n(001)| = 2^{n-1}. \quad (4.1)$$

One readily observes that, for $n \geq 2$, there is a natural bijection between 001-avoiding inversion sequences of length $n-1$ and 0012-avoiding inversion sequences of length n in which the last entry equals $n-1$. Such a bijection could be simply constructed by appending $n-1$ to the end of the 001-avoiding inversion sequences. Therefore, we have an enumeration result as follows.

Corollary 4.2. *For $n \geq 1$,*

$$|\{e \in \mathbf{I}_n(0012) : \text{last}(e) = n-1\}| = \begin{cases} 1 & \text{if } n = 1, \\ 2^{n-2} & \text{if } n \geq 2. \end{cases} \quad (4.2)$$

Notice that Corollary 4.2 is equivalent to

$$\begin{aligned} \mathcal{L}(1; q) &= \sum_{n \geq 1} \left(\sum_{k=0}^{n-1} f_n(k, n-1) \right) q^n \\ &= q + q^2 + 2q^3 + 4q^4 + 8q^5 + 16q^6 + \cdots \\ &= \frac{q(1-q)}{1-2q}. \end{aligned}$$

Now we prove a bivariate strengthening of the above that will be utilized in our proof of Theorem 1.1.

Theorem 4.3. *We have*

$$\mathcal{L}(x; q) = \frac{q(1-q)^2}{(1-2q)(1-xq)}. \quad (4.3)$$

Proof. For $n \geq 2$, it follows from (a), (b) and (c) of Lemma 3.1 that

$$\begin{aligned} \sum_{k=0}^{n-1} f_n(k, n-1)x^k &= x^{n-1} + \sum_{k=0}^{n-3} \sum_{k'=k}^{n-2} f_{n-1}(k', n-2)x^k \\ &= x^{n-1} + \sum_{k'=0}^{n-3} f_{n-1}(k', n-2) \sum_{k=0}^{k'} x^k + f_{n-1}(n-2, n-2) \sum_{k=0}^{n-3} x^k \\ &= x^{n-1} + \sum_{k'=0}^{n-3} f_{n-1}(k', n-2) \frac{1-x^{k'+1}}{1-x} + \frac{1-x^{n-2}}{1-x}. \end{aligned}$$

Therefore,

$$L_n(x) = x^{n-1} + \frac{1}{1-x} (L_{n-1}(1) - xL_{n-1}(x)) - \frac{1-x^{n-1}}{1-x} + \frac{1-x^{n-2}}{1-x}.$$

Multiplying the above by q^n and summing over $n \geq 2$, we have

$$\mathcal{L}(x; q) - q = \frac{q}{1-x} \mathcal{L}(1; q) - \frac{xq}{1-x} \mathcal{L}(x; q) - \frac{q^2(1-x)}{1-xq},$$

or

$$(1 - xq)(1 - x + xq)\mathcal{L}(x; q) = q(1 - xq)\mathcal{L}(1; q) + q(1 - q)(1 - x). \quad (4.4)$$

Applying the kernel method (see [9, Exercice 4, §2.2.1, p. 243] or [17]) yields

$$\begin{cases} 1 - x + xq = 0, \\ q(1 - xq)\mathcal{L}(1; q) + q(1 - q)(1 - x) = 0. \end{cases}$$

Solving the first equation of the system for x gives

$$x = \frac{1}{1 - q}.$$

Substituting the above into the second equation of the system, we have

$$\mathcal{L}(1; q) = \frac{q(1 - q)}{1 - 2q}.$$

Substituting the above back to (4.4), we arrive at (4.3). \square

5. Proof of Theorem 1.1

We first establish two relations concerning $\mathcal{D}(x; q)$.

Lemma 5.1. *We have*

$$\mathcal{D}(x; q) = \frac{1}{1 - xq}\mathcal{D}(0; q) \quad (5.1)$$

$$= \frac{q}{1 - xq}\mathcal{F}(1, 1; q). \quad (5.2)$$

Proof. We know from Lemma 3.3 that

$$\begin{aligned} \sum_{n \geq 2} \sum_{k=0}^{n-2} f_n(k, k) x^k q^n &= \sum_{n \geq 2} \sum_{k=0}^{n-2} f_{n-k}(0, 0) x^k q^n \\ &\stackrel{(\text{with } n' = n - k)}{=} \sum_{n' \geq 2} \sum_{n \geq n'} f_{n'}(0, 0) x^{n-n'} q^n \\ &= \sum_{n' \geq 2} f_{n'}(0, 0) x^{-n'} \sum_{n \geq n'} (xq)^n \\ &= \frac{1}{1 - xq} \sum_{n' \geq 2} f_{n'}(0, 0) q^{n'}. \end{aligned}$$

Noticing that $D_1(x) = 0$, we have

$$\mathcal{D}(x; q) = \frac{1}{1 - xq}\mathcal{D}(0; q),$$

which is the first part of the lemma. For the second part, we deduce from Lemma 3.1(d) that

$$\begin{aligned} \mathcal{D}(0; q) &= \sum_{n \geq 2} f_n(0, 0) q^n \\ &= \sum_{n \geq 2} \sum_{\ell'=0}^{n-2} \sum_{k'=0}^{\ell'} f_{n-1}(k', \ell') q^n \\ &= q\mathcal{F}(1, 1; q). \end{aligned}$$

Therefore, (5.2) follows. \square

Next, we show a relation between $\mathcal{F}(x, 1; q)$ and $\mathcal{F}(1, 1; q)$.

Lemma 5.2. *We have*

$$\mathcal{F}(x, 1; q) = \frac{1-q}{1-xq} \mathcal{F}(1, 1; q). \quad (5.3)$$

Proof. For $n \geq 2$, it follows from Lemma 3.1(d) that

$$\begin{aligned} D_n(x) &= \sum_{\ell=0}^{n-2} f_n(\ell, \ell) x^\ell \\ &= \sum_{\ell=0}^{n-2} \sum_{\ell'=\ell}^{n-2} \sum_{k'=\ell}^{\ell'} f_{n-1}(k', \ell') x^\ell \\ &= \sum_{\ell'=0}^{n-2} \sum_{k'=0}^{\ell'} f_{n-1}(k', \ell') \sum_{k=0}^{k'} x^\ell \\ &= \sum_{\ell'=0}^{n-2} \sum_{k'=0}^{\ell'} f_{n-1}(k', \ell') \frac{1-x^{k'+1}}{1-x} \\ &= \frac{1}{1-x} (F_{n-1}(1, 1) - xF_{n-1}(x, 1)). \end{aligned}$$

Therefore,

$$\mathcal{D}(x; q) = \frac{q}{1-x} (\mathcal{F}(1, 1; q) - x\mathcal{F}(x, 1; q)).$$

Substituting (5.2) into the above yields

$$\frac{q}{1-xq} \mathcal{F}(1, 1; q) = \frac{q}{1-x} (\mathcal{F}(1, 1; q) - x\mathcal{F}(x, 1; q)),$$

from which (5.3) follows. \square

We then construct a functional equation for $\mathcal{F}(x, y; q)$.

Lemma 5.3. *We have*

$$\begin{aligned} &\left(1 + \frac{xq}{1-x} + \frac{yq}{1-y}\right) \mathcal{F}(x, y; q) \\ &= \frac{q}{1-x} \mathcal{F}(1, y; q) + \frac{q(1-q)}{(1-y)(1-xyq)} \mathcal{F}(1, 1; q) + \frac{q(1-q-2yq+2yq^2+y^2q^2)}{(1-2yq)(1-xyq)}. \end{aligned} \quad (5.4)$$

Proof. We first observe that

$$\begin{aligned} \sum_{\ell=0}^{n-2} f_n(\ell, \ell) x^\ell y^\ell + \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} f_n(k, \ell) x^k y^\ell &= F_n(x, y) - \sum_{k=0}^{n-1} f_n(k, n-1) x^k y^{n-1} \\ &= F_n(x, y) - y^{n-1} L_n(x). \end{aligned} \quad (5.5)$$

Notice also that

$$\sum_{\ell=0}^{n-2} f_n(\ell, \ell) x^\ell y^\ell = D_n(xy). \quad (5.6)$$

Now, by Lemma 3.1(e), we may separate

$$\begin{aligned} \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} f_n(k, \ell) x^k y^\ell &= \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{k'=k}^{\ell} f_{n-1}(k', \ell) x^k y^\ell \\ &\quad + \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{\ell'=\ell}^{n-2} f_{n-1}(k, \ell') x^k y^\ell. \end{aligned}$$

We further notice that the first term on the right-hand side can be separated as

$$\sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{k'=k}^{\ell} f_{n-1}(k', \ell) x^k y^\ell = \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{k'=k}^{\ell-1} f_{n-1}(k', \ell) x^k y^\ell + \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} f_{n-1}(\ell, \ell) x^k y^\ell.$$

We have

$$\begin{aligned} &\sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{k'=k}^{\ell-1} f_{n-1}(k', \ell) x^k y^\ell \\ &= \sum_{\ell=1}^{n-2} \sum_{k'=0}^{\ell-1} f_{n-1}(k', \ell) y^\ell \sum_{k=0}^{k'} x^k \\ &= \sum_{\ell=1}^{n-2} \sum_{k'=0}^{\ell-1} f_{n-1}(k', \ell) y^\ell \frac{1 - x^{k'+1}}{1 - x} \\ &= \sum_{\ell=0}^{n-2} \sum_{k'=0}^{\ell} f_{n-1}(k', \ell) y^\ell \frac{1 - x^{k'+1}}{1 - x} - \sum_{\ell=0}^{n-2} f_{n-1}(\ell, \ell) y^\ell \frac{1 - x^{\ell+1}}{1 - x} \\ &= \frac{1}{1 - x} (F_{n-1}(1, y) - x F_{n-1}(x, y)) \\ &\quad - \frac{1}{1 - x} (D_{n-1}(y) + y^{n-2} - x D_{n-1}(xy) - x^{n-1} y^{n-2}). \end{aligned}$$

Also,

$$\begin{aligned} \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} f_{n-1}(\ell, \ell) x^k y^\ell &= \sum_{\ell=1}^{n-2} f_{n-1}(\ell, \ell) y^\ell \frac{1 - x^\ell}{1 - x} \\ &= \sum_{\ell=0}^{n-2} f_{n-1}(\ell, \ell) y^\ell \frac{1 - x^\ell}{1 - x} \\ &= \frac{1}{1 - x} (D_{n-1}(y) + y^{n-2} - D_{n-1}(xy) - x^{n-2} y^{n-2}). \end{aligned}$$

On the other hand,

$$\begin{aligned} \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} \sum_{\ell'=\ell}^{n-2} f_{n-1}(k, \ell') x^k y^\ell &= \sum_{\ell'=1}^{n-2} \sum_{k=0}^{\ell'-1} f_{n-1}(k, \ell') x^k \sum_{\ell=k+1}^{\ell'} y^\ell \\ &= \sum_{\ell'=1}^{n-2} \sum_{k=0}^{\ell'-1} f_{n-1}(k, \ell') x^k \frac{y^{k+1} - y^{\ell'+1}}{1 - y} \\ &= \sum_{\ell'=0}^{n-2} \sum_{k=0}^{\ell'} f_{n-1}(k, \ell') x^k \frac{y^{k+1} - y^{\ell'+1}}{1 - y} \end{aligned}$$

$$= \frac{y}{1-y} (F_{n-1}(xy, 1) - F_{n-1}(x, y)).$$

Therefore,

$$\begin{aligned} & \sum_{\ell=1}^{n-2} \sum_{k=0}^{\ell-1} f_n(k, \ell) x^k y^\ell \\ &= \frac{1}{1-x} (F_{n-1}(1, y) - xF_{n-1}(x, y)) + \frac{y}{1-y} (F_{n-1}(xy, 1) - F_{n-1}(x, y)) \\ & \quad - D_{n-1}(xy) - x^{n-2}y^{n-2}. \end{aligned} \quad (5.7)$$

It follows from (5.5), (5.6) and (5.7) that

$$\begin{aligned} & F_n(x, y) - y^{n-1}L_n(x) \\ &= D_n(xy) + \frac{1}{1-x} (F_{n-1}(1, y) - xF_{n-1}(x, y)) \\ & \quad + \frac{y}{1-y} (F_{n-1}(xy, 1) - F_{n-1}(x, y)) - D_{n-1}(xy) - x^{n-2}y^{n-2}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \mathcal{F}(x, y; q) - y^{-1}\mathcal{L}(x; yq) \\ &= \mathcal{D}(xy; q) + \frac{q}{1-x} (\mathcal{F}(1, y; q) - x\mathcal{F}(x, y; q)) \\ & \quad + \frac{yq}{1-y} (\mathcal{F}(xy, 1; q) - \mathcal{F}(x, y; q)) - q\mathcal{D}(xy; q) - \frac{q^2}{1-xyq}. \end{aligned}$$

Applying (4.3), (5.2) and (5.3) gives the desired result. \square

With the assistance of the kernel method, we may deduce a functional equation satisfied by $\mathcal{F}(1, y; q)$.

Lemma 5.4. *We have*

$$\mathcal{F}(1, y; q) = \frac{q}{1-y+y^2q} \mathcal{F}(1, 1; q) + \frac{q(1-y)(1-q-2yq+2yq^2+y^2q^2)}{(1-q)(1-2yq)(1-y+y^2q)}. \quad (5.8)$$

Proof. We multiply both sides of (5.4) by $(1-x)(1-y)$. Then

$$\begin{aligned} & ((1-y+yq) - x(1-y-q+2yq)) \mathcal{F}(x, y; q) \\ &= q(1-y) \mathcal{F}(1, y; q) + \frac{q(1-q)(1-x)}{1-xyq} \mathcal{F}(1, 1; q) \\ & \quad + \frac{q(1-x)(1-y)(1-q-2yq+2yq^2+y^2q^2)}{(1-2yq)(1-xyq)}. \end{aligned}$$

We treat the kernel polynomial as a function in x and solve

$$(1-y+yq) - x(1-y-q+2yq) = 0$$

so that

$$x = \frac{1-y+yq}{1-y-q+2yq}.$$

Substituting the above into

$$0 = q(1-y) \mathcal{F}(1, y; q) + \frac{q(1-q)(1-x)}{1-xyq} \mathcal{F}(1, 1; q)$$

$$+ \frac{q(1-x)(1-y)(1-q-2yq+2yq^2+y^2q^2)}{(1-2yq)(1-xyq)},$$

we arrive at (5.8) after simplification. \square

Finally, we are ready to complete the proof of Theorem 1.1.

Proof of Theorem 1.1. It is known that (cf. [18, A279561])

$$1 + \sum_{n \geq 1} \left(1 + \sum_{i=1}^{n-1} \binom{2i}{i-1} \right) q^n = \frac{1-4q+(1-2q)\sqrt{1-4q}}{2(1-q)(1-4q)}. \quad (5.9)$$

We then rewrite (5.8) as

$$(1-y+y^2q)\mathcal{F}(1,y;q) = q\mathcal{F}(1,1;q) + \frac{q(1-y)(1-q-2yq+2yq^2+y^2q^2)}{(1-q)(1-2yq)}.$$

We treat the kernel polynomial as a function in y and solve

$$1-y+y^2q=0.$$

Then

$$y_{1,2} = \frac{1 \mp \sqrt{1-4q}}{2q}.$$

We choose the solution

$$y_1 = \frac{1 - \sqrt{1-4q}}{2q}$$

since $y_1 = 1 + q + O(q^2)$ as $q \rightarrow 0$. Substituting $y = y_1$ into

$$0 = q\mathcal{F}(1,1;q) + \frac{q(1-y)(1-q-2yq+2yq^2+y^2q^2)}{(1-q)(1-2yq)},$$

we find that

$$\begin{aligned} \mathcal{F}(1,1;q) &= \frac{-(1-2q)(1-4q) + (1-2q)\sqrt{1-4q}}{2(1-q)(1-4q)} \\ &= \frac{1-4q + (1-2q)\sqrt{1-4q}}{2(1-q)(1-4q)} - 1. \end{aligned} \quad (5.10)$$

This implies that for $n \geq 1$,

$$1 + \sum_{i=1}^{n-1} \binom{2i}{i-1} = \sum_{\ell=0}^{n-1} \sum_{k=0}^{\ell} f_n(k, \ell) = |\mathbf{I}_n(0012)|.$$

Therefore, Conjecture 1.1 is true. \square

References

1. J. S. Auli, Pattern avoidance in inversion sequences, Thesis (Ph.D.)—Dartmouth College. 2020. 160 pp.
2. J. S. Auli and S. Elizalde, Consecutive patterns in inversion sequences, *Discrete Math. Theor. Comput. Sci.* **21** (2019), no. 2, Paper No. 6, 22 pp.
3. J. S. Auli and S. Elizalde, Consecutive patterns in inversion sequences II: Avoiding patterns of relations, *J. Integer Seq.* **22** (2019), no. 7, Art. 19.7.5, 37 pp.
4. J. S. Auli and S. Elizalde, Wilf equivalences between vincular patterns in inversion sequences, *Appl. Math. Comput.* **388** (2021), 125514, 15 pp.
5. N. R. Beaton, M. Bouvel, V. Guerrini, and S. Rinaldi, Enumerating five families of pattern-avoiding inversion sequences; and introducing the powered Catalan numbers, *Theoret. Comput. Sci.* **777** (2019), 69–92.

6. W. Cao, E. Y. Jin, and Z. Lin, Enumeration of inversion sequences avoiding triples of relations, *Discrete Appl. Math.* **260** (2019), 86–97.
7. S. Corteel, M. Martinez, C. D. Savage, and M. Weselcouch, Patterns in inversion sequences I, *Discrete Math. Theor. Comput. Sci.* **18** (2016), no. 2, Paper No. 2, 21 pp.
8. D. Kim and Z. Lin, Refined restricted inversion sequences, *Sém. Lothar. Combin.* **78B** (2017), Art. 52, 12 pp.
9. D. E. Knuth, *The Art of Computer Programming, Vol. 1: Fundamental Algorithms*, Addison-Wesley, 1973, Third edition, 1997.
10. Z. Lin, Restricted inversion sequences and enhanced 3-noncrossing partitions, *European J. Combin.* **70** (2018), 202–211.
11. Z. Lin, Patterns of relation triples in inversion and ascent sequences, *Theoret. Comput. Sci.* **804** (2020), 115–125.
12. Z. Lin and S. Fu, On $\underline{120}$ -avoiding inversion and ascent sequences, *European J. Combin.* **93** (2021), 103282, 12 pp.
13. Z. Lin and D. Kim, A sextuple equidistribution arising in pattern avoidance, *J. Combin. Theory Ser. A* **155** (2018), 267–286.
14. Z. Lin and S. H. F. Yan, Vincular patterns in inversion sequences, *Appl. Math. Comput.* **364** (2020), 124672, 17 pp.
15. T. Mansour and M. Shattuck, Pattern avoidance in inversion sequences, *Pure Math. Appl. (P.U.M.A.)* **25** (2015), no. 2, 157–176.
16. M. Martinez and C. D. Savage, Patterns in inversion sequences II: inversion sequences avoiding triples of relations, *J. Integer Seq.* **21** (2018), no. 2, Art. 18.2.2, 44 pp.
17. H. Prodinger, The kernel method: a collection of examples, *Sém. Lothar. Combin.* **50** (2003/04), Art. B50f, 19 pp.
18. N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences; <http://oeis.org>.
19. C. Yan and Z. Lin, Inversion sequences avoiding pairs of patterns, *Discrete Math. Theor. Comput. Sci.* **22** (2020), no. 1, Paper No. 23, 35 pp.
20. S. H. F. Yan, Bijections for inversion sequences, ascent sequences and 3-nonnesting set partitions, *Appl. Math. Comput.* **325** (2018), 24–30.

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA
 E-mail address: shanechern@psu.edu; chenxiaohang92@gmail.com