

Hankel determinants and Jacobi continued fractions for q -Euler numbers

Shane Chern

陈小航

Universität Wien (维也纳大学)

chenxiaohang92@gmail.com

@ Chongqing Normal University

Chongqing

Jul 12, 2024

(Joint work with Lin Jiu 酒霖)

Introduction



Lin Jiu (Duke Kunshan)

Introduction



Lin Jiu (Duke Kunshan)



Karl Dilcher (Dalhousie)

- *J. Math. Anal. Appl.* **497** (2021), no. 1, Paper No. 124855, 18 pp.
- *Int. J. Number Theory* **18** (2022), no. 2, 331–359.
- *Contrib. Discrete Math.* **18** (2023), no. 2, 146–175.

International Journal of Number Theory
Vol. 18, No. 2 (2022) 331–359
© World Scientific Publishing Company
DOI: 10.1142/S179304212250021X



J. Math. Anal. Appl. 497 (2021) 124855



Orthogonal polynomials and Hankel determinants for certain Bernoulli and Euler polynomials

Karl Dilcher^a, Lin Jiu^{a,b,*}

^a Department of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia, B3H 4R2, Canada

^b Za Changshi Center for Mathematics and Computational Sciences, Duke Kunshan University, Kunshan, Suzhou, Jiangsu Province, 215316, P. R. China



ARTICLE INFO

Article history:
Received 2 July 2020
Available online 14 December 2020
Submitted by B.C. Berndt

Keywords:
Bernoulli polynomial
Euler polynomial
Hankel determinant
Orthogonal polynomial
Continued fraction
Polygamma function

ABSTRACT

Using continued fraction expansions of certain polygamma functions as a main tool, we find orthogonal polynomials with respect to the odd-index Bernoulli polynomials $B_{2k+1}(x)$ and the Euler polynomials $E_{2k+1}(x)$, for $v = 0, 1, 2$. In the process we also determine the corresponding Jacobi continued fractions (or J -fractions) and Hankel determinants. In all these cases the Hankel determinants are polynomials in x which factor completely over the rationals.

© 2020 Elsevier Inc. All rights reserved.

Hankel determinants of sequences related to Bernoulli and Euler polynomials

Karl Dilcher

Department of Mathematics and Statistics
Dalhousie University, Halifax, Nova Scotia, Canada, B3H 4R2
dilcher@mathstat.dal.ca

Lin Jiu*

Za Changshi Center for Mathematics and Computational Sciences
Duke Kunshan University
Kunshan, Suzhou, Jiangsu Province 215316, P. R. China
lin.jiu@dukekunshan.edu.cn



Contributions to Discrete Mathematics

Volume 18, Number 2, Pages 146–175
ISSN 1715-0468

HANKEL DETERMINANTS OF SHIFTED SEQUENCES OF BERNOULLI AND EULER NUMBERS

KARL DILCHER AND LIN JIU*

ABSTRACT. Hankel determinants of sequences related to Bernoulli and Euler numbers have been studied before, and numerous identities are known. However, when a sequence is shifted by one unit, the situation often changes significantly. In this paper we use classical orthogonal polynomials and related methods to prove a general result concerning Hankel determinants for shifted sequences. We then apply this result to obtain new Hankel determinant evaluations for a total of 14 sequences related to Bernoulli and Euler numbers, one of which concerns Euler polynomials.

Introduction

- A *Hankel matrix* $(M_{i,j})$ is a square matrix with constant skew diagonals, i.e., $M_{i,j} = M_{i',j'}$ whenever $i + j = i' + j'$:

$$\begin{pmatrix} s_0 & s_1 & s_2 & \cdots & s_n \\ s_1 & s_2 & s_3 & \cdots & s_{n+1} \\ s_2 & s_3 & s_4 & \cdots & s_{n+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ s_n & s_{n+1} & s_{n+2} & \cdots & s_{2n} \end{pmatrix}$$

- Letting $\{s_n\}_{n \geq 0}$ be a sequence in a field \mathbb{K} , one may define its associated Hankel matrices by $(s_{i+j})_{0 \leq i,j \leq n}$.
- We are interested in the **determinant** of these matrices.

Introduction

Example 1:

- Bernoulli numbers $B_n = (1, -\frac{1}{2}, \frac{1}{6}, 0, -\frac{1}{30}, 0, \frac{1}{42}, 0, -\frac{1}{30}, 0, \frac{5}{66}, \dots)$:

$$\sum_{n \geq 0} B_n \frac{t^n}{n!} = \frac{t}{e^t - 1}.$$

Hankel determinants:

$$\det(1) = 1, \det \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{6} \end{pmatrix} = -\frac{1}{12}, \det \begin{pmatrix} 1 & -\frac{1}{2} & \frac{1}{6} \\ -\frac{1}{2} & \frac{1}{6} & 0 \\ \frac{1}{6} & 0 & -\frac{1}{30} \end{pmatrix} = -\frac{1}{540}, \dots$$

$$1, -\frac{1}{12}, -\frac{1}{540}, \frac{1}{42000}, \frac{1}{3215625}, -\frac{4}{623959875}, \dots$$

Al-Salam and Carlitz (1959):

$$\det_{0 \leq i, j \leq n} (B_{i+j}) = (-1)^{\binom{n+1}{2}} \prod_{k=1}^n \frac{(k!)^6}{(2k)!(2k+1)!}.$$

Example 2:

- Euler numbers $E_n = (1, 0, -1, 0, 5, 0, -61, 0, 1385, 0, -50521, \dots)$:

$$\sum_{n \geq 0} E_n \frac{t^n}{n!} = \frac{2}{e^t + e^{-t}}.$$

Hankel determinants:

$$1, -1, -4, 144, 82944, -1194393600, \dots$$

Al-Salam and Carlitz (1959):

$$\det_{0 \leq i, j \leq n} (E_{i+j}) = (-1)^{\binom{n+1}{2}} \prod_{k=1}^n (k!)^2.$$

The World of q

- q -integers: For $m \in \mathbb{Z}$,

$$[m]_q := \frac{1 - q^m}{1 - q}.$$

- q -factorials: For $M \in \mathbb{N}$,

$$[M]_q! := \prod_{m=1}^M [m]_q.$$

- q -Pochhammer symbols: For $N \in \mathbb{N} \cup \{\infty\}$,

$$(A; q)_N := \prod_{k=0}^{N-1} (1 - Aq^k),$$

$$(A, B, \dots, C; q)_N := (A; q)_N (B; q)_N \cdots (C; q)_N.$$

- q -Bernoulli numbers β_n (Carlitz, 1948):

$$\beta_n := \frac{1}{(1-q)^n} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{k+1}{[k+1]_q}.$$

Alternatively, they can be recursively defined by $\beta_0 = 1$ and for $n \geq 1$,

$$\sum_{k=0}^n \binom{n}{k} q^{k+1} \beta_k - \beta_n = \begin{cases} 1, & \text{if } n = 1, \\ 0, & \text{if } n \geq 2. \end{cases}$$

Nombres de q -Bernoulli–Carlitz et fractions continues

[F Chapoton](#), [J Zeng](#) - Journal de théorie des nombres de Bordeaux, 2017 - numdam.org

Carlitz a introduit vers 1950 des q -analogues des nombres de Bernoulli. On obtient une représentation de ces q -analogues (ainsi que de variantes décalées) comme moments de ...

☆ Save 📄 Cite Cited by 7 Related articles 🔗

The World of q

• Hankel determinants for q -Bernoulli numbers:

$$\det_{0 \leq i, j \leq n} (\beta_{i+j}) = (-1)^{\binom{n+1}{2}} q^{\binom{n+1}{3}} \prod_{k=1}^n \frac{([k]_q!)^6}{[2k]_q! [2k+1]_q!}.$$

THE STARTING POINT

1. THE CARLITZ β

Carlitz [1, 2] generated the Bernoulli numbers to the sequence β_n , by the recurrence:

$$\sum_{k=0}^n \binom{n}{k} \beta_k q^{k+1} - \beta_m = \begin{cases} 1, & m = 1; \\ 0, & m > 1, \end{cases}$$

with also the value $\beta_0 = 1$.

Definition 1. The q -bracket is defined by

$$[x]_q := \frac{1 - q^x}{1 - q},$$

for all $x \in \mathbb{R}$ and $q > 0$. The q -factorial is then defined by

$$[k]_q! := [k]_q [k-1]_q \cdots [1]_q.$$

Conjecture 2.

$$H_n(\beta_k) = (-1)^{\binom{n+1}{2}} q^{\frac{(n-1)n(n+1)}{6}} \frac{\prod_{k=1}^n [k]_q^{6(n+1-k)}}{\prod_{k=1}^{2n+1} [k]_q^{2n+2-k}}.$$

From Lin, Mar 14, 2023

From Shane on Apr 04, 2023.

Throughout, let us use Carlitz's q -Bernoulli numbers β_k given by $\beta_0 = 1$ and for $n \geq 1$,

$$\sum_{k=0}^n \binom{n}{k} \beta_k q^{k+1} - \beta_n = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases}$$

“Need to Prove” 1.

$$\sum_{k \geq 0} \beta_k x^k = \frac{\beta_0}{1 + a_0 x - \frac{b_1 x^2}{1 + a_1 x - \frac{b_2 x^2}{1 + a_2 x - \cdots}}}, \quad (1)$$

where

$$a_n = \frac{1}{(1 + q^n)(1 + q^{n+1})} \left(-\frac{(1 - q^n)(1 - q^{n+1})}{1 - q} + 2q^n \right), \quad (2)$$

$$b_n = -\frac{q^{n-1}(1 - q^n)^6}{(1 - q)^2(1 - q^{2n-1})(1 - q^{2n})^2(1 - q^{2n+1})}. \quad (3)$$

From Shane, Apr 04, 2023

Nombres de q -Bernoulli–Carlitz et fractions continues

[F Chapoton](#), [J Zeng](#)

Journal de théorie des nombres de Bordeaux, 2017 · [numdam.org](#)

Resume

Carlitz a introduit vers 1950 des q -analogues des nombres de Bernoulli. On obtient une représentation de ces q -analogues (ainsi que de variantes décalées) comme moments de certains polynômes orthogonaux. Ceci donne aussi des factorisations des déterminants de Hankel des nombres de q -Bernoulli, ainsi que des fractions continues pour leurs séries génératrices. Certains de ces résultats sont des q -analogues d'énoncés connus pour les nombres de Bernoulli, mais d'autres sont sans version classique.

Abstract. q -Bernoulli–Carlitz Numbers and continuous fractions.

[numdam.org](#)

SHOW LESS ^

☆ Save  Cite Cited by 6 Related articles All 11 versions 

Théorème 4.2. *On a, pour les matrices d'indices $0 \leq i, j \leq n - 1$,*

$$(4.7) \quad \det (\beta_{i+j})_{i,j} = (-1)^{\binom{n}{2}} q^{\binom{n}{3}} \prod_{i=1}^{n-1} \frac{[i]_q!^6}{[2i]_q! [2i+1]_q!},$$

The World of q

- q -Euler numbers ϵ_n (Carlitz, 1948):

$$\epsilon_n := \frac{1}{(1-q)^n} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{1+q}{1+q^{k+1}}.$$

Alternatively, they can be recursively defined by $\epsilon_0 = 1$ and for $n \geq 1$,

$$\sum_{k=0}^n \binom{n}{k} q^{k+1} \epsilon_k + \epsilon_n = 0.$$

- At the $q \rightarrow 1$ limit, ϵ_n reduces to

$$1, -\frac{1}{2}, 0, \frac{1}{4}, 0, -\frac{1}{2}, 0, \frac{17}{8}, 0, -\frac{31}{2}, \dots,$$

which is identical to $E_n(0)$ rather than $2^n E_n(\frac{1}{2})$, the Euler numbers E_n . Here the *Euler polynomials* $E_n(x)$ are given by

$$\sum_{n \geq 0} E_n(x) \frac{t^n}{n!} = \frac{2e^{xt}}{e^t + 1}.$$

Theorem (C.-Jiu, 2023)

$$\begin{aligned} & \det_{0 \leq i, j \leq n} (\epsilon_{i+j}) \\ &= \frac{(-1)^{\binom{n+1}{2}} q^{\frac{1}{4} \binom{2n+2}{3}}}{(1-q)^{n(n+1)}} \prod_{k=1}^n \frac{(q^2, q^2; q^2)_k}{(-q, -q^2, -q^2, -q^3; q^2)_k}, \end{aligned}$$

$$\begin{aligned} & \det_{0 \leq i, j \leq n} (\epsilon_{i+j+1}) \\ &= \frac{(-1)^{\binom{n+2}{2}} q^{\frac{1}{4} \binom{2n+4}{3}}}{(1-q)^{n(n+1)} (1+q^2)^{n+1}} \prod_{k=1}^n \frac{(q^2, q^4; q^2)_k}{(-q^2, -q^3, -q^3, -q^4; q^2)_k}, \end{aligned}$$


$$\begin{aligned} & \det_{0 \leq i, j \leq n} (\epsilon_{i+j+2}) \\ &= \frac{(-1)^{\binom{n+2}{2}} q^{\frac{1}{4} \binom{2n+4}{3}} (1+q)^n (1 - (-1)^n q^{(n+2)^2})}{(1-q)^{n(n+1)} (1+q^2)^{2(n+1)} (1+q^3)^{n+1}} \prod_{k=1}^n \frac{(q^4, q^4; q^2)_k}{(-q^3, -q^4, -q^4, -q^5; q^2)_k}. \end{aligned}$$

- C. Krattenthaler, Advanced determinant calculus, *Sém. Lothar. Combin.* **42** (1999), Art. B42q, 67 pp.
- C. Krattenthaler, Advanced determinant calculus: a complement, *Linear Algebra Appl.* **411** (2005), 68–166.



ELSEVIER

Available online at www.sciencedirect.com



Linear Algebra and its Applications 411 (2005) 68–166

LINEAR ALGEBRA
AND ITS
APPLICATIONS

www.elsevier.com/locate/laa

ADVANCED DETERMINANT CALCULUS

C. KRATTENTHALER[†]

Institut für Mathematik der Universität Wien,
Strudlhofgasse 4, A-1090 Wien, Austria.

E-mail: kratt@pap.univie.ac.at

WWW: <http://radon.mat.univie.ac.at/People/kratt>

*Dedicated to the pioneer of determinant evaluations (among many other things),
George Andrews*

ABSTRACT. The purpose of this article is threefold. First, it provides the reader with a few useful and efficient tools which should enable her/him to evaluate nontrivial determinants for the case such a determinant should appear in her/his research. Second, it lists a number of such determinants that have been already evaluated, together with explanations which tell in which contexts they have appeared. Third, it points out references where further such determinant evaluations can be found.

Advanced determinant calculus: A complement[☆]

C. Krattenthaler

*Institut Camille Jordan, Université Claude Bernard Lyon-I, 21, avenue Claude Bernard,
F-69622 Villeurbanne Cedex, France*

Received 16 March 2005; accepted 6 June 2005

Submitted by R.A. Brualdi

Abstract

This is a complement to my previous article “Advanced Determinant Calculus” [C. Krattenthaler, Advanced determinant calculus, *Séminaire Lotharingien Combin.* 42 (1999) (“The Andrews Festschrift”), Article B42q, 67 pp.]. In the present article, I share with the reader my experience of applying the methods described in the previous article in order to solve a particular problem from number theory [G. Almkvist, C. Krattenthaler, J. Petersson, Some new formulas for π , *Experiment. Math.* 12 (2003) 441–456]. Moreover, I add a list of determinant evaluations which I consider as interesting, which have been found since the appearance of the previous article, or which I failed to mention there, including several conjectures and open problems.

© 2005 Elsevier Inc. All rights reserved.

Jacobi continued fractions (a.k.a. J-fractions):

Fact (Heilermann)

Let $\{\mu_k\}_{k \geq 0}$ be a sequence such that its generating function $\sum_{k \geq 0} \mu_k x^k$ has the *J-fraction* expression

$$\sum_{k \geq 0} \mu_k x^k = \frac{\mu_0}{1 + a_0 x - \frac{b_1 x^2}{1 + a_1 x - \frac{b_2 x^2}{1 + a_2 x - \ddots}}}}.$$

Then for $n \geq 0$,

$$\det_{0 \leq i, j \leq n} (\mu_{i+j}) = \mu_0^{n+1} b_1^n b_2^{n-1} \cdots b_{n-1}^2 b_n.$$

Jacobi continued fractions (a.k.a. J-fractions):

Fact (Heilermann (continued...))

Further, define $\{p_n(z)\}_{n \geq 0}$ a family of polynomials given by a three-term recursive relation for $n \geq 1$,

$$p_{n+1}(z) = (a_n + z)p_n(z) - b_n p_{n-1}(z),$$

with initial conditions $p_0(z) = 1$ and $p_1(z) = a_0 + z$. Then

$$\det_{0 \leq i, j \leq n} (\mu_{i+j+1}) = \det_{0 \leq i, j \leq n} (\mu_{i+j}) \cdot (-1)^{n+1} p_{n+1}(0).$$

Orthogonal polynomials: A family of polynomials $\{p_n(z)\}_{n \geq 0}$ with $p_n(z)$ of degree n is called *orthogonal* if there is a linear functional L on the space of polynomials in z such that $L(p_m(z)p_n(z)) = \delta_{m,n}\sigma_n$ where $\delta_{m,n}$ is the Kronecker delta and $\{\sigma_n\}_{n \geq 0}$ is a fixed nonzero sequence.

Orthogonal polynomials \Leftrightarrow Three-term recurrences:

Fact (Favard, Stieltjes)

Let $\{p_n(z)\}_{n \geq 0}$ be a family of monic polynomials with $p_n(z)$ of degree n . Then they are orthogonal if and only if there exist sequences $\{a_n\}_{n \geq 0}$ and $\{b_n\}_{n \geq 1}$ with $b_n \neq 0$ such that $p_0(z) = 1$, $p_1(z) = a_0 + z$, and for $n \geq 1$,

$$p_{n+1}(z) = (a_n + z)p_n(z) - b_n p_{n-1}(z).$$

Orthogonal polynomials \Rightarrow J-fractions:

Fact

Let L be an associated linear functional for a family of orthogonal monic polynomials $\{p_n(z)\}_{n \geq 0}$ with $p_n(z)$ of degree n . Then

$$\sum_{k \geq 0} L(z^k) x^k = \frac{L(z^0)}{1 + a_0 x - \frac{b_1 x^2}{1 + a_1 x - \frac{b_2 x^2}{1 + a_2 x - \ddots}}},$$

where $\{a_n\}_{n \geq 0}$ and $\{b_n\}_{n \geq 1}$ are given by the associated three-term recurrence.

How to evaluate Hankel determinants:

- STEP 1: Guess the J -fraction expression
- STEP 2: Formulate the associated orthogonal polynomials
- STEP 3: Determine a suitable linear functional
- STEP 4: Check orthogonality under this linear functional

STEP 1: Guess the J -fraction expression.

A

B

$$\text{Out[69]} = \left\{ \frac{q}{1+q^2}, \frac{-1-q+2q^3+q^4+q^5}{(1+q^2)(1+q^4)}, \frac{-1-q-q^2-q^3+2q^5+q^6+q^7+q^8+q^9}{(1+q^2)(1+q^4)(1-q^2+q^4)}, \right. \\ \left. \frac{-1-q-q^2-q^3-q^4-q^5+2q^7+q^8+q^9+q^{10}+q^{11}+q^{12}+q^{13}}{(1+q^2)(1-q^2+q^4)(1+q^8)}, \right. \\ \left. \frac{-1-q-q^2-q^3-q^4-q^5-q^6-q^7+2q^9+q^{10}+q^{11}+q^{12}+q^{13}+q^{14}+q^{15}+q^{16}+q^{17}}{(1+q^2)(1+q^8)(1-q^2+q^4-q^6+q^8)} \right\}$$

$$\text{Out[70]} = \left\{ -\frac{q}{(1+q^2)^2(1-q+q^2)}, -\frac{q^3(1+q^2)^2}{(1-q+q^2)(1+q^4)^2(1-q+q^2-q^3+q^4)}, \right. \\ \left. -\frac{q^5(1-q+q^2)^2(1+q+q^2)^2}{(1+q^2)^2(1-q^2+q^4)^2(1-q+q^2-q^3+q^4)(1-q+q^2-q^3+q^4-q^5+q^6)}, \right. \\ \left. -\frac{q^7(1+q^2)^2(1+q^4)^2}{(1-q+q^2)(1-q^3+q^6)(1-q+q^2-q^3+q^4-q^5+q^6)(1+q^8)^2}, \right. \\ \left. -\left(\frac{q^9(1-q+q^2-q^3+q^4)^2(1+q+q^2+q^3+q^4)^2}{(1-q^2+q^4-q^6+q^8)^2(1-q+q^2-q^3+q^4-q^5+q^6-q^7+q^8-q^9+q^{10})} \right) \right\}$$

STEP 1: Guess the J -fraction expression.

Guess(!)

Let $\ell \in \{0, 1\}$. Then


$$\sum_{k \geq 0} \epsilon_{k+\ell} x^k \stackrel{?}{=} \frac{\epsilon_\ell}{1 + a_{\ell,0}x - \frac{b_{\ell,1}x^2}{1 + a_{\ell,1}x - \frac{b_{\ell,2}x^2}{1 + a_{\ell,2}x - \dots}}},$$

where

$$a_{\ell,n} = \frac{q^{2n+\ell}(1+q)(1+q^\ell)}{(1-q)(1+q^{2n+\ell})(1+q^{2n+\ell+2})} - \frac{1}{1-q},$$

$$b_{\ell,n} = -\frac{q^{2n+2\ell-1}(1-q^{2n})(1-q^{2n+2\ell})}{(1-q)^2(1+q^{2n+\ell-1})(1+q^{2n+\ell})^2(1+q^{2n+\ell+1})}.$$

STEP 2: Formulate the associated orthogonal polynomials.



R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric orthogonal polynomials and their q -analogues*, Springer-Verlag, Berlin, 2010.

STEP 2: Formulate the associated orthogonal polynomials.

☞ *Big q -Jacobi polynomials:*

$$\mathcal{J}_{\ell,n}(z) := {}_3\phi_2 \left(\begin{matrix} q^{-n}, -q^{n+\ell+1}, z \\ q^{\ell+1}, 0 \end{matrix}; q, q \right).$$

Theorem

Define

$$\mathcal{P}_{\ell,n}(z) := \frac{(-1)^n (q^{\ell+1}; q)_n}{q^n (1-q)^n (-q^{n+\ell+1}; q)_n} {}_3\phi_2 \left(\begin{matrix} q^{-n}, -q^{n+\ell+1}, q(1-(1-q)z) \\ q^{\ell+1}, 0 \end{matrix}; q, q \right).$$

We have $\mathcal{P}_{\ell,0}(z) = 1$, $\mathcal{P}_{\ell,1}(z) = a_{\ell,0} + z$, and for $n \geq 1$,

$$\mathcal{P}_{\ell,n+1}(z) = (a_{\ell,n} + z)\mathcal{P}_{\ell,n}(z) - b_{\ell,n}\mathcal{P}_{\ell,n-1}(z).$$

STEP 3: Determine a suitable linear functional.

Want to study the linear function Φ with

$$\Phi(z^n) = \epsilon_n.$$

☞ Choose a “nice” basis for the vector space of polynomials over $\mathbb{Q}(q)$:

$$\left\{ \begin{bmatrix} n, z \\ n \end{bmatrix}_q \right\}_{n \geq 0},$$

where (cf. Chapoton–Essouabri, 2015)

$$\begin{bmatrix} m, z \\ n \end{bmatrix}_q := \frac{1}{[n]_q!} \prod_{k=m-n+1}^m ([k]_q + q^k z).$$

STEP 3: Determine a suitable linear functional.

Let Φ be the linear functional on $\mathbb{Q}(q)[z]$ given by

$$\Phi \left(\begin{bmatrix} n, z \\ n \end{bmatrix}_q \right) := \frac{1}{(-q^2; q)_n} \quad (n \geq 0).$$

Lemma

For $0 \leq m \leq n$,

$$\Phi \left(\begin{bmatrix} m, z \\ n \end{bmatrix}_q \right) = \frac{(-1)^{n-m} q^{n-m}}{(-q^2; q)_n}.$$

Lemma

For $n \geq 0$,

$$\Phi \left(\begin{bmatrix} n+1, z \\ n \end{bmatrix}_q \right) = \frac{1+q}{q} - \frac{1}{q(-q^2; q)_n}.$$

STEP 3: Determine a suitable linear functional.

Theorem

For any $P(z) \in \mathbb{Q}(q)[z]$,

$$q\Phi(P(1 + qz)) + \Phi(P(z)) = (1 + q)P(0).$$

Theorem

For $n \geq 0$,

$$\Phi(z^n) = \epsilon_n.$$

Proof. We first notice that $\Phi(z^0) = \Phi(1) = 1 = \epsilon_0$. Now for $n \geq 1$, we apply the above theorem with $P(z) = z^n$, and derive that

$$q\Phi((1 + qz)^n) + \Phi(z^n) = 0,$$

namely,

$$\sum_{k=0}^n \binom{n}{k} q^{k+1} \Phi(z^k) + \Phi(z^n) = 0.$$

This recursive relation for $\Phi(z^n)$ is identical to that for ϵ_n !!!

STEP 4: Check orthogonality under this linear functional.

Let $\ell \in \{0, 1\}$. We define two linear functionals Φ_ℓ on $\mathbb{Q}(q)[z]$ by

$$\Phi_\ell(z^n) := \Phi(z^{n+\ell}) \quad (n \geq 0).$$

Theorem

Let $\ell \in \{0, 1\}$. The family of monic polynomials $\{\mathcal{P}_{\ell,n}(z)\}_{n \geq 0}$ is orthogonal under the linear functional Φ_ℓ .

☞ Show that for $\ell \in \{0, 1\}$, the identity $\Phi_\ell(\mathcal{P}_{\ell,n}(z)) = 0$ holds whenever $n \geq 1$:

$$\begin{aligned} \mathcal{P}_{0,n}(z) &= \frac{(-1)^n (q; q)_n}{q^n (1-q)^n (-q^{n+1}; q)_n} \sum_{k=0}^n \frac{q^k (q^{-n}, -q^{n+1}; q)_k}{(q; q)_k} \begin{bmatrix} k, z \\ k \end{bmatrix}_q, \\ z \cdot \mathcal{P}_{1,n}(z) &= \frac{(-1)^n (q^2; q)_n}{q^n (1-q)^n (-q^{n+2}; q)_n} \sum_{k=0}^n \frac{q^k (q^{-n}, -q^{n+2}; q)_k}{(q; q)_k} \begin{bmatrix} k, z \\ k+1 \end{bmatrix}_q. \end{aligned}$$

Question A: Families of q -Hankel determinants?

Note that

$$\mathcal{P}_{\ell,n}(z) = \frac{(-1)^n}{q^n(1-q)^n} \tilde{\mathcal{J}}_{\ell,n}((q^2 - q)z + q),$$

where $\tilde{\mathcal{J}}_{\ell,n}(z)$ are normalizations of the big q -Jacobi $\mathcal{J}_{\ell,n}(z)$ as monic polynomials:

$$\tilde{\mathcal{J}}_{\ell,n}(z) := \frac{(q^{\ell+1}; q)_n}{(-q^{n+\ell+1}; q)_n} \mathcal{J}_{\ell,n}(z),$$

Theorem (C.-Jiu, 2023)

For each nonnegative integer ℓ , define a sequence $\{\xi_{\ell,n}\}_{n \geq 0}$ by

$$\xi_{\ell,n} := \frac{q^{(\ell+1)n}(-q; q)_n}{(-q^{\ell+2}; q)_n}.$$

Then

$$\det_{0 \leq i,j \leq n} (\xi_{\ell,i+j}) = (-1)^{\binom{n+1}{2}} q^{2\binom{n+2}{3} + (2\ell+1)\binom{n+1}{2}} \prod_{k=1}^n \frac{(q^2, q^{2\ell+2}; q^2)_k}{(-q^{\ell+1}, -q^{\ell+2}, -q^{\ell+2}, -q^{\ell+3}; q^2)_k}.$$

Question B: Bernoulli/Euler polynomials?

- Bernoulli polynomials $B_n(x)$ and Euler polynomials $E_n(x)$:

$$\sum_{n \geq 0} B_n(x) \frac{t^n}{n!} = \frac{te^{xt}}{e^t - 1} \quad \text{and} \quad \sum_{n \geq 0} E_n(x) \frac{t^n}{n!} = \frac{2e^{xt}}{e^t + 1}.$$

In particular,

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k x^{n-k} \quad \text{and} \quad E_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{E_k}{2^k} x^{n-k}.$$

- **Invariance of Hankel determinants under the binomial transform:** For every $n \geq 0$, define

$$s_n(x) := \sum_{k=0}^n \binom{n}{k} s_k x^{n-k}.$$

Then

$$\det_{0 \leq i, j \leq n} (s_{i+j}) = \det_{0 \leq i, j \leq n} (s_{i+j}(x)).$$

Question B: Bernoulli/Euler polynomials?

- Trivial determinant evaluations(!):

$$\det_{0 \leq i, j \leq n} (B_{i+j}(x)) = (-1)^{\binom{n+1}{2}} \prod_{k=1}^n \frac{(k!)^6}{(2k)!(2k+1)!},$$

$$\det_{0 \leq i, j \leq n} (E_{i+j}(x)) = \left(-\frac{1}{4}\right)^{\binom{n+1}{2}} \prod_{k=1}^n (k!)^2,$$

- Dilcher–Jiu (2021):**

$$\det_{0 \leq i, j \leq n} (B_{2(i+j)}(\frac{x+1}{2})) = ???,$$

$$\det_{0 \leq i, j \leq n} (E_{2(i+j)}(\frac{x+1}{2})) = (-1)^{\binom{n+1}{2}} \prod_{k=1}^n \left(\frac{k^2(x^2 - (2k-1)^2)}{4} \right)^{n-k+1},$$

$$\det_{0 \leq i, j \leq n} (B_{2(i+j)+1}(\frac{x+1}{2})) = (-1)^{\binom{n+1}{2}} \left(\frac{x}{2}\right)^{n+1} \prod_{k=1}^n \left(\frac{k^4(x^2 - k^2)}{4(2k-1)(2k+1)} \right)^{n-k+1},$$

$$\det_{0 \leq i, j \leq n} (E_{2(i+j)+1}(\frac{x+1}{2})) = (-1)^{\binom{n+1}{2}} \left(\frac{x}{2}\right)^{n+1} \prod_{k=1}^n \left(\frac{k^2(x^2 - 4k^2)}{4} \right)^{n-k+1}.$$

Question B: Bernoulli/Euler polynomials?

arXiv > math > arXiv:2401.17220

Search... All fields  Search

[Help](#) | [Advanced Search](#)

Mathematics > Number Theory

[Submitted on 30 Jan 2024]

Leading coefficient in the Hankel determinants related to binomial and q -binomial transforms

Shane Chern, Lin Jiu, Shuhan Li, Liuquan Wang

It is a standard result that the Hankel determinants for a sequence stay invariant after performing the binomial transform on this sequence. In this work, we extend the scenario to q -binomial transforms and study the behavior of the leading coefficient in such Hankel determinants. We also investigate the leading coefficient in the Hankel determinants for even-indexed Bernoulli polynomials with recourse to a curious binomial transform. In particular, the degrees of these Hankel determinants share the same nature as those in one of the q -binomial cases.

Subjects: **Number Theory (math.NT)**; Combinatorics (math.CO)

MSC classes: Primary 11C20, Secondary 11B68, 33D45

Cite as: arXiv:2401.17220 [math.NT]
(or arXiv:2401.17220v1 [math.NT] for this version)
<https://doi.org/10.48550/arXiv.2401.17220> 

Submission history

From: Lin Jiu [[view email](#)]

[v1] Tue, 30 Jan 2024 18:05:37 UTC (15 KB)

Access Paper:

- [Download PDF](#)
 - [Other Formats](#)
- [view license](#)

Current browse context:

math.NT

[< prev](#) | [next >](#)
[new](#) | [recent](#) | [2401](#)

Change to browse by:

[math](#)
[math.CO](#)

References & Citations

- [NASA ADS](#)
- [Google Scholar](#)
- [Semantic Scholar](#)

Export BibTeX Citation

Bookmark



Question B: Bernoulli/Euler polynomials?

Theorem (C.-Jiu-Li-Wang, 2024)

For every $n \geq 0$, $\det_{0 \leq i, j \leq n} (B_{2(i+j)}(\frac{x+1}{2}))$ is a polynomial in x of degree $n(n+1)$ with leading coefficient

$$[x^{n(n+1)}] \det_{0 \leq i, j \leq n} (B_{2(i+j)}(\frac{x+1}{2})) = (-1)^{\binom{n+1}{2}} \prod_{j=1}^n \frac{(j!)^6}{(2j)!(2j+1)!}.$$

- It is notable that the Hankel determinant $\det_{0 \leq i, j \leq n} (B_{2(i+j)}(\frac{x+1}{2}))$ is a linear combination of the terms

$$B_{2j_0}(\frac{1+x}{2}) B_{2(1+j_1)}(\frac{1+x}{2}) \cdots B_{2(n+j_n)}(\frac{1+x}{2})$$

with j_0, j_1, \dots, j_n a permutation of $0, 1, \dots, n$. Here, each term is of degree $\sum_{i=0}^n 2(i+j_i) = 2n(n+1)$. However, our result states that the degree of the Hankel determinant is $n(n+1)$, which is only **half** of the above terms, thereby indicating abundant cancelations of higher powers of x in this determinant expansion.

Question C: q -Binomial transform?

- Consider the q -binomial transform

$$\tilde{\alpha}_n(x) := \sum_{k=0}^n q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q \alpha_k x^{n-k}.$$

- We *no longer* have invariance of Hankel determinants under this q -binomial transform.
- There is a “**half** the degree” phenomenon.

Theorem (C.–Jiu–Li–Wang, 2024)

For every $n \geq 0$, $\det_{0 \leq i, j \leq n} (\tilde{\alpha}_{i+j}(x))$ is a polynomial in x of degree $\frac{n(n+1)}{2}$ with leading coefficient

$$\left[x^{\frac{n(n+1)}{2}} \right] \det_{0 \leq i, j \leq n} (\tilde{\alpha}_{i+j}(x)) = \alpha_0 \alpha_1 \cdots \alpha_n (-1)^{\binom{n+1}{2}} q^{2\binom{n+1}{3}} \prod_{j=1}^n (1 - q^j)^{n+1-j}.$$

Question C: q -Binomial transform?

- Special choices of α ?

Theorem (C.–Jiu–Li–Wang, 2024)

Choose the sequence $\alpha_k^{u,v} := q^{-\binom{k}{2}}(u; q)_k v^k$ with u and v indeterminates so as to let

$$\tilde{\alpha}_k^{u,v}(x) := \sum_{\ell=0}^k \begin{bmatrix} k \\ \ell \end{bmatrix}_q (u; q)_\ell v^\ell x^{k-\ell} = \sum_{\ell=0}^k \begin{bmatrix} k \\ \ell \end{bmatrix}_q (u; q)_{k-\ell} v^{k-\ell} x^\ell.$$

Then

$$\det_{0 \leq i, j \leq n} (\tilde{\alpha}_{i+j}^{u,v}(x)) = v^{\binom{n+1}{2}} q^{\binom{n+1}{3}} \prod_{j=1}^n (uvq^{j-1} - x)^{n+1-j} (u, q; q)_{n+1-j}.$$

Thank You!