APEX INTERNATIONAL ACADEMY

CBSE Senior Secondary Affiliation No. 1930482 | Academic Excellence & Research Division

OFFICIAL ACADEMIC SOLUTION & EVALUATION DOSSIER

VERIFIED & GRADED

ASSIGNMENT: Human Digestive System Overview

SUBJECT: General Science (SCI101)

CLASS & SECTION: Class 5 - Section A

ACADEMIC YEAR: 2025 - 2026

MARKS EVALUATED: 94 / 100 (Grade A)

TEACHER DIRECTIVE & PROBLEM FOCUS:
Chart the path of food from mouth to stomach with annotations of enzymes, secretions, and villi absorption.

1. Executive Overview & Problem Context

The human gastrointestinal tract is an eight-meter continuous muscular tube coordinating mechanical digestion and enzymatic chemical hydrolysis. Ingested macromolecular carbohydrates, proteins, and lipids are disassembled into absorbable monosaccharides, amino acids, and fatty acids to fuel cellular cellular respiration.

2. Core Analysis, Solutions & Systematic Working

- **Oral Cavity: Mastication & Salivary Amylase:**

  Mechanical crushing by incisors and molars mixes food with saliva secreted by parotid, submandibular, and sublingual glands. Salivary Amylase (Ptyalin) initiates carbohydrate hydrolysis, converting cooked starches into maltose disaccharides at pH ~6.8, forming a soft lubricated bolus.

- **Esophagus & Peristaltic Propulsion:**

  The bolus triggers the swallowing reflex, closing the cartilaginous epiglottis over the trachea. Rhythmic involuntary contractions of circular and longitudinal smooth muscles (peristalsis) propel food past the lower esophageal sphincter into the stomach.

- **Gastric Digestion in the Stomach:**

  Gastric mucosa secretes 2-3 liters of gastric juice daily: Hydrochloric Acid (HCl, creating pH 1.5-2.0 to denature proteins and neutralize pathogens), Pepsinogen (cleaved by acid into active Pepsin to hydrolyze proteins into peptides), and protective alkaline mucus preventing gastric autodigestion.

- **Small Intestine: Enzymatic Breakdown & Villi Absorption:**

  In the Duodenum, acidic chyme is neutralized by alkaline sodium bicarbonate from the Pancreas. Liver bile salts emulsify dietary lipids. Pancreatic Lipase, Trypsin, and Chymotrypsin complete digestion. Millions of microscopic finger-like projections called Villi and Microvilli amplify luminal surface area by 600-fold to maximize nutrient absorption into capillary and lacteal bloodstreams.

APEX INTERNATIONAL ACADEMY

CBSE Senior Secondary Affiliation No. 1930482 | Academic Excellence & Research Division

OFFICIAL ACADEMIC SOLUTION & EVALUATION DOSSIER

VERIFIED & GRADED

3. Detailed Data Matrix & Systematic Findings

Digestive Organ	Secreted Fluid / Enzyme	Target Substrate	End Product Formed
Salivary Glands	Salivary Amylase (Ptyalin)	Complex Starch	Maltose & Dextrins
Stomach Lining	Hydrochloric Acid + Pepsin	Dietary Proteins	Peptides & Polypeptides
Liver / Gallbladder	Bile Salts & Pigments	Large Fat Globules	Emulsified Fat Droplets
Exocrine Pancreas	Lipase, Trypsin, Amylase	Proteins, Fats, Starch	Fatty acids, Amino acids
Small Intestine Villi	Erepsin, Maltase, Lactase	Oligosaccharides & Peptides	Glucose, Galactose, Amino acids

4. Student Conclusions & Reflective Synthesis

Digestion represents an integrated biological process where mechanical maceration and enzymatic hydrolysis convert ingested nourishment into bioavailable molecular precursors for metabolic energy and cellular maintenance.

5. Official Faculty Evaluation & Grading Rubric

ASSESSMENT CRITERIA BREAKDOWN

Conceptual Clarity & Factual Accuracy: 28 / 30

Analytical Depth & Research Methodology: 28 / 30

Problem Solving & Structured Working: 18 / 20

Presentation, Organization & Neatness: 14 / 20

FACULTY FEEDBACK REMARK:

"Outstanding physiological depth! Biochemical enzyme mapping and anatomical organ diagrams are clear and accurate."

APEX EVALUATION DIVISION

VERIFIED & ARCHIVED