Correlations with inclination within tertiles of z, M*, SFR, DM_host

	low tertile	mid tertile	high tertile	all	pooled	partial τ
z vs i, binned by z (N = 55)	+0.14 (0.57) MC +0.12	+0.36 (0.14) MC +0.32	-0.24 (0.35) MC -0.21	-0.02 (0.86) MC -0.04	+0.09 (0.54)	n/a
M* vs i, binned by z (N = 51)	+0.10 (0.72) MC +0.08	+0.21 (0.41) MC +0.14	+0.15 (0.57) MC +0.20	+0.14 (0.31) MC +0.13	+0.15 (0.33)	+0.11 (0.21) MC +0.09
SFR vs i, binned by z (N = 46)	-0.39 (0.14) MC -0.35	+0.38 (0.19) MC +0.31	-0.12 (0.65) MC -0.19	-0.12 (0.42) MC -0.14	-0.07 (0.68)	-0.06 (0.56) MC -0.09
DM_host vs i, binned by z (N = 55)	+0.16 (0.50) MC +0.14	-0.19 (0.44) MC -0.13	+0.02 (0.95) MC -0.05	+0.06 (0.64) MC +0.02	-0.00 (0.99)	+0.05 (0.64) MC +0.01
z vs i, binned by M* (N = 51)	-0.05 (0.86) MC -0.06	-0.06 (0.83) MC -0.10	-0.24 (0.34) MC -0.10	-0.04 (0.76) MC -0.05	-0.12 (0.45)	-0.05 (0.61) MC -0.05
M* vs i, binned by M* (N = 52)	+0.06 (0.81) MC -0.10	+0.18 (0.47) MC +0.16	-0.16 (0.52) MC -0.04	+0.11 (0.43) MC +0.10	+0.03 (0.86)	n/a
SFR vs i, binned by M* (N = 46)	**-0.64 (0.013)** **MC -0.58**	+0.24 (0.37) MC +0.13	-0.11 (0.69) MC -0.03	-0.12 (0.42) MC -0.14	-0.18 (0.29)	-0.08 (0.36) MC -0.11
DM_host vs i, binned by M* (N = 51)	-0.06 (0.82) MC -0.02	+0.28 (0.29) MC +0.29	+0.07 (0.78) MC -0.07	+0.12 (0.39) MC +0.06	+0.09 (0.55)	+0.09 (0.38) MC +0.05
z vs i, binned by SFR (N = 46)	-0.05 (0.84) MC -0.16	+0.03 (0.91) MC +0.14	-0.03 (0.92) MC -0.04	-0.07 (0.65) MC -0.07	-0.02 (0.92)	-0.03 (0.75) MC -0.02
M* vs i, binned by SFR (N = 46)	-0.08 (0.77) MC -0.13	-0.14 (0.62) MC +0.05	+0.32 (0.24) MC +0.32	+0.10 (0.52) MC +0.09	+0.03 (0.84)	+0.09 (0.38) MC +0.08
SFR vs i, binned by SFR (N = 46)	-0.44 (0.087) MC -0.27	+0.10 (0.72) MC +0.04	-0.29 (0.29) MC -0.35	-0.12 (0.42) MC -0.14	-0.23 (0.17)	n/a
DM_host vs i, binned by SFR (N = 46)	+0.03 (0.91) MC +0.10	+0.37 (0.18) MC +0.11	+0.07 (0.81) MC -0.02	+0.12 (0.42) MC +0.07	+0.16 (0.35)	+0.10 (0.28) MC +0.06
z vs i, binned by DM_host (N = 55)	+0.05 (0.83) MC +0.01	-0.11 (0.67) MC +0.02	-0.14 (0.58) MC -0.19	-0.02 (0.86) MC -0.04	-0.06 (0.67)	-0.02 (0.86) MC -0.02
M* vs i, binned by DM_host (N = 51)	-0.06 (0.83) MC +0.05	**+0.53 (0.027)** **MC +0.39**	-0.04 (0.89) MC -0.01	+0.14 (0.31) MC +0.13	+0.17 (0.29)	+0.10 (0.27) MC +0.09
SFR vs i, binned by DM_host (N = 46)	-0.41 (0.13) MC -0.16	+0.18 (0.51) MC -0.13	-0.35 (0.20) MC -0.25	-0.12 (0.42) MC -0.14	-0.19 (0.25)	-0.09 (0.44) MC -0.10
DM_host vs i, binned by DM_host (N = 55)	-0.08 (0.76) MC -0.01	+0.28 (0.24) MC +0.04	-0.22 (0.39) MC -0.19	+0.06 (0.64) MC +0.02	+0.00 (0.98)	n/a

Cells: Spearman r_S (p) for the tertiles, all hosts and the Fisher-pooled within-tertile value; Kendall partial τ given the binning quantity.
MC: median over 2,000 draws through the measurement errors with tertiles re-formed. Bold: $p < 0.05$. Black frame: Benjamini–Hochberg $q < 0.05$ over the tertiles of the grid.